

Crop Report – September 9, 2026



[Crop Pest Report](#) [Crop Weather Report](#) [Reporting Area Map](#) [Seasonal Reports](#) [Weekly Weather Maps](#)

Inside this Report...

Upcoming Events
Weather Report
Commodity Reports
Cereals
Oilseeds
Pulses & Soybeans
Forages & Livestock
Regional Comments
Southwest
Northwest
Central
Eastern
Interlake

Upcoming Events

- [CropTalk Webinar](#): 9:00 AM, September 9, 2026. Fall Harvest Weather Update – Alison Sass.
- [Special Crops Field Workshops](#): Sept. 9 (Beausejour), Sept. 17 (Melita).

Weather Report

- Another week of scattered extreme weather brought heavy rains to some regions. Thunderstorms early in the week brought large rain accumulations and strong winds to the western regions with Manitou seeing a maximum windspeed of 90 km/hr and Minitonas accumulating 37 mm of rain on August 31. Severe storms on September 6 and 7 brought strong winds and heavy rains to parts of the Central and Interlake regions. Portage la Prairie received 86.9 mm in just one hour on September 6. Other areas that saw intense hourly accumulations include Stonewall, Selkirk, Woodlands, and Elie. Powerful winds caused significant damage throughout the City of Winnipeg and west of Elie.
- Climate normals for total accumulated precipitation from May 1 to September 7 range from 230 mm to 356 mm and are based on 30-year historical data. Since May 1, large areas of the Northwest, Southwest, and Interlake regions have accumulated more than 350 mm of precipitation. Seasonal accumulations are above normal for the majority of the Northwest and Southwest regions. Large areas of the Northwest, Southwest, and Interlake regions have seen accumulations over 120% of normal. A few localized areas of the Interlake, Central and Eastern regions have seen accumulations below 90% of the 30-year average.
- Total Accumulation of Growing Degree Days shows the accumulated growing degree days (GDD) for the period of May 1 – September 7, 2026. GDD accumulations ranged between 1281 and 1760 GDD so far this growing season. Most of southern Manitoba has accumulated more than 1500 GDD.

To find interactive soil temperature/moisture and air temperature information see Agri-Maps Current Weather [viewer](#)

Table 1. Range of measurements of eight-day accumulated precipitation in Manitoba's Agricultural Regions. (August 31 – September 7, 2026).

Region	Wettest location last seven days	Driest location last seven days
Central	Portage la Prairie (112.7 mm)	Plumas (13.5 mm)
Eastern	Gardenton (77.4 mm)	Dugald (28.9 mm)
Interlake	Stonewall (98.2 mm)	Lake Francis (18.3 mm)
Northwest	Minitonas (67.6 mm)	Roblin (19.0 mm)
Southwest	Souris (87.3 mm)	Kola (19.8 mm)

Table 2: Percentage of Harvest Completion by Crop and Region to September 8, 2026.

Crop	Southwest	Northwest	Central	Eastern	Interlake	MB AVG
Winter Wheat	95%	80%	100%	99%	100%	95%
Fall Rye	95%	80%	100%	99%	100%	95%
Spring Wheat	25%	20%	90%	90%	70%	54%
Barley	40%	20%	85%	90%	75%	58%
Oats	30%	0%	80%	85%	75%	51%
Field Pea	90%	15%	100%	100%	98%	82%
Canola	5%	2%	30%	20%	20%	15%
Regional AVG	14%	8%	46%	43%	36%	28%

Crops still unharvested, or negligible acres displayed as – or omitted from this table.

Commodity Reports

Note: All yields are reported in comparison to 10-year (2015-2024) averages by region based on MASC published data. For reference, please refer to the [regional average tables by crop](#).

Cereals

- Winter cereal harvest is complete across the province, with average yields.
- Spring cereal harvest is more than 20% complete in all regions, with yield reports of average to above average with good quality.
- Majority of corn is in the early dent stage.

Oilseeds

- Canola harvest has started in all regions. A significant number of acres have been swathed this year.
- The majority of sunflowers range from R7 to R8 stage.

Pulses & Soybeans

- Majority of soybeans range from late R6 to R7 stage.
- Pea harvest is nearly complete, with the majority of yields reported as being average and good quality.

Forages & Livestock

Forages

- Pastures remain in good condition, with recent rainfall and warm temperatures supporting strong fall regrowth and extending grazing potential into the fall.
- Heavy and frequent rainfall has delayed forage and cereal harvest in areas across the province.
- Most producers are wrapping up hay production for the year and are focusing on hauling hay and straw.
- Second-cut hay harvest is nearing completion, while some producers continue working to put up slough hay where conditions allow.
- Straw baling has been delayed in some areas by wet conditions.
- Cereal silage harvest is nearly complete, with producers reporting above-average yields. Corn silage harvest has begun in some areas but has been temporarily delayed by recent rainfall.
- Adequate moisture has resulted in good regrowth for grazing following the harvest of polycrop fields for silage.

Livestock

- Cattle generally remain in good condition on pasture.
- Recent rainfall has improved pasture conditions in areas where forage supplies were beginning to decline. Some producers in the interlake region have begun supplementing cattle out on pasture.
- A few producers have begun moving cattle onto fall grazing areas and have begun weaning.
- Cases of foot rot and pinkeye continue to gradually decline.
- Recent rainfall has replenished dugouts, with water supplies generally full or around 90% capacity.
- Manure application is increasing as more fields become available for spreading.

Regional Comments

Southwest

There was a good start to the week, as most producers started harvesting and grain was dry by Saturday. Heavy rains on Sunday in the early afternoon put a stop to all harvesting, as most areas received rain, with some areas reporting hail. Several areas received over 2 inches of rain. There are now a lot of lodged crops, making harvest even more difficult than it already was. Cereal grain quality has already started to deteriorate, with some producers reporting sprouting. There is also severe lodging in most canola crops, primarily due to a mixture of disease, heavy rains, and strong winds.

Winter wheat and fall rye are about 90–95% complete, with reports of average yields and good quality.

Peas are about 90–95% complete, with average yields and good quality. The remaining crop will likely be affected by the recent rains, and quality is expected to be downgraded.

Spring wheat harvest is about 20–25% complete, with average to above-average yields and good quality so far. However, the recent rains are likely to change that once producers can get back into the fields.

Barley is about 40% complete, with average to above-average yields. Quality has started to decline due to the recent rains, and following the latest rainfall, producers are already reporting concerns with sprouting.

Canola harvest is minimal, at less than 10% complete. Reports indicate average yields and good quality so far. Several fields are being swathed, mostly because of severe lodging caused by disease. Several fields are also being sprayed.

Soybeans are in the late R6 to early R7 stage. Some fields are showing changes in leaf color, with some pods starting to turn. The crop is taking advantage of the recent moisture.

Corn and sunflowers continue to develop as the crops take advantage of the moisture and good growing conditions.

Cereal silage is pretty much complete, and some corn silage has begun.

Northwest

Wet conditions early in the week slowed harvest progress in some areas. Warmer temperatures through the week improved field conditions, allowing progress, particularly in areas that received less precipitation. Unsettled weather returned to the region late in the weekend, with heavy rainfall in some areas again slowing harvest progress.

As harvest slowly progresses, many challenges from the season still exist due to infrastructure damage from spring flooding. While major roads have been repaired, there are still roads that are washed out and fields with washouts and cuts.

Fall rye and winter wheat harvest continues and is approximately 95% complete.

Majority of the field pea crop has moved into R7, and harvest continues where crops have reached maturity. Approximately 50% of the crop is harvested. Reported yields have been mixed, ranging from below average to average.

Spring wheat is mostly in the hard dough stage with some later seeded fields behind. Pre-harvest glyphosate applications continue as stages are reached. Approximately 20% of the spring wheat is harvested.

Canola fields continue to range in terms of stages, with most of the crop nearing harvest maturity. Desiccation and swathing is underway in fields that have reached the appropriate stage. Fields that experienced excess moisture have patches of delayed growth that are still flowering. A small start to canola harvest has begun in some early seeded fields.

Majority of the soybean crop is in the R6-R7 stage and looking good where excess moisture hasn't delayed growth.

Central

The region was very dry this past week; however, many locations received enough rainfall Sunday evening to benefit long-season crops, especially corn and soybeans.

Overall, the dry conditions have allowed producers to make good progress with harvest. Cool nights and heavy dews have occasionally delayed harvest and resulted in much of the harvested crop requiring additional drying.

Winter wheat and fall rye harvest is complete. Spring cereal harvest continues, with spring wheat yield and protein percentages ranging from average to good. Barley yields are reported as slightly above average. Oat test weights, however, are lower than typical, likely due to lower soil moisture during critical development stages.

Soybeans are at the R6 – R7 growth stages. The crop would still benefit from additional rainfall. In areas where soil moisture is lowest, leaves are already yellowing and dropping due to a combination of limited rainfall and hot temperatures. Many plants are producing predominantly two-seeded pods, with very few four-seeded pods. Pod abortion has also occurred in the upper canopy in many fields. The soybean crop is now past the stages where soybean aphid is a concern, and populations appear to be naturally declining, likely due to the abundance of beneficial insects in fields. White mold (*Sclerotinia*) is showing up in some soybean fields. This is primarily occurring in fields with the most crop growth, which are likely also among those with the highest yield potential.

Almost all of the pea crop has been harvested, with the remaining fields expected to be completed over the coming days. Quality has been good, while yields have been average.

Canola harvest and swathing continue. Late-season diseases, including sclerotinia and verticillium stripe, are noticeable. More canola is being swathed this year than in recent years, likely due to increasing levels of verticillium stripe. Pre-harvest sprouting has also been an issue in some fields.

Corn ranges from the late milk to early dent stages. Although the crop would benefit from additional rainfall, the majority of the corn crop appears to have fared relatively well despite the dry conditions experienced over the past few weeks.

Sunflowers are at the R7 to R8 growth stages.

Many producers are mowing down patches of weeds, particularly kochia and amaranth species suspected of herbicide resistance. Producers are also taking advantage of soybean fields that are turning color to scout for potential waterhemp and Palmer amaranth plants, which can be more easily identified and rouged at this stage. Producers are also scouting recently harvested cereal, pea, and canola fields for new waterhemp emergence and regrowth. Amaranth species can emerge in late August and early September and may produce seed before freeze-up.

Eastern

The region saw rainfall this last week ranging from 29 mm to 64 mm. Rainfall was sufficient to bring harvesting activities to a halt over the weekend. Producers are hoping for sunny and windy weather to dry things up quickly and resume harvest.

Winter wheat/fall rye harvest is finishing up in the region and estimated at 99% complete. Yield reports have been slightly below average.

Overall, the spring wheat harvest is estimated at 90% complete. All spring cereal yields have been reported as average at best. Field drains and lower drowned out areas of fields from the spring's heavy rainfall continue to bring yield numbers down.

Most corn is in the dent stage of kernel development. The crop is continuing to grow well as starch accumulates in the kernel. On average the milk line is at the 50% mark. Warm weather and recent rainfall will continue to help in the crop development as starch continues to accumulate in the kernels.

Canola harvest became more general across the region this past week as producers finished up their cereals. An estimate of 15% of acres are complete. Yields are below average to average (field dependent). The lower yields are thought to be due to the hot dry weather during the flowering period.

Soybeans are at the R6 to early R7 stage. Many fields have leaves that are continuing to change color. Overall, fields are looking good. Producers in the region are thankful for the rain we received as it will help with the pod filling stage and increase yields.

Interlake

The region received significant rainfall over the weekend, delaying harvest. South Interlake areas recorded the highest rainfall. The Stonewall area received 98.2mm, Selkirk 78.1mm, Woodlands 66.7mm and Petersfield 47.7mm. There are reports of lodging and hail damage from the storms.

Variable growing conditions this season have resulted in a wide range of yields.

Winter wheat and fall rye harvest is 100% complete, with average to above-average yields where field conditions were favorable. Flood-affected areas had below-average yields.

Spring wheat harvest is about 70% complete, with average to below-average yields. Quality remains #1, although ergot has been reported. Oat and barley harvest is 75% complete, with average to slightly above-average yields. Most oats have adequate weight, though some are light. Silage and grain corn are developing well, with most fields at the dent stage.

Soybeans are rapidly turning in most fields and have reached the R7 stage, although some plants are short. Yellow-brown leaves are increasingly visible, with most advanced fields beginning to drop leaves. Moisture stress is evident in some Southern Interlake fields, while recent rains are supporting pod fill.

Pea harvest is about 98% complete in most areas, with slightly above-average yields.

Canola harvest is 20% complete, with yields ranging from below average to average based on field conditions. Sunflowers are at R7, and most advanced fields are entering R8.

Rain is increasing weed pressure, with more waterhemp and redroot pigweed visible. Grasshoppers are being monitored, with no major concerns.