

Crop Report – July 7, 2026



[Crop Pest Report](#) [Crop Weather Report](#) [Reporting Area Map](#) [Seasonal Reports](#) [Weekly Weather Maps](#)

Inside this Report...

Upcoming Events
Weather Report
Commodity Reports
Cereals
Oilseeds
Pulses & Soybeans
Forages & Livestock
Regional Comments
Southwest
Northwest
Central
Eastern
Interlake
Agronomy Topics
Assessing Hail Damage in Cereal Crops



Flooded road near Roblin
Photo credit: Elizabeth Nernberg

Upcoming Events

- [CropTalk Webinar](#): Diversification Centre Update, Wednesday, July 8 at 9 am
- [Factsheet on Assessing the Impact of Excess Moisture in Crops](#)

Weather Report

- Variable precipitation occurred over the past 7 days across agro-Manitoba. Once again, rains and isolated thunderstorms brought hail, strong winds, intense rain, and possible tornadoes swept through the majority of the province. The largest accumulations occurred in the Southwest near Boissevain and Souris, as well as the Northwest region in the already water-logged areas of San Clara, Ethelbert, Dauphin, Minnedosa, and Russell. Overland flooding and overflowing rivers and streams have caused many communities, including the City of Brandon, to declare local states of emergency. Weekly accumulations ranged from 9 (Sprague) to 188 mm (Boissevain). A tornado was confirmed in the Whyte Ridge neighborhood of Winnipeg on June 29.
- Climate normals for total accumulated precipitation from May 1 to July 5 range from 131 mm to 211 mm (Table 2) and are based on 30-year historical data. Since May 1, areas of the Northwest and Interlake regions have accumulated 220 mm of precipitation. Frequent precipitation events have brought seasonal accumulations well above normal for the entire Northwest and Interlake regions. Areas of the Central and Southwest region also have above-normal accumulations due to recent rainfall. Areas of the Northwest have accumulated more than 200% of normal precipitation. A few localized areas of the Central, and Eastern regions have seen accumulations under 80% of the 30-year average.
- Total Accumulation of Growing Degree Days shows the accumulated growing degree days (GDD) for the period of May 1 – July 5, 2026. GDD accumulations ranged between 497 and 731 GDD so far this growing season. Most of Southern Manitoba has accumulated more than 600 GDD.

To find interactive soil temperature/moisture and air temperature information see Agri-Maps Current Weather [viewer](#)

Range of measurements of seven-day accumulated precipitation in Manitoba's Agricultural Regions (June 29-July 5, 2026)

Region	Wettest location last seven days	Driest location last seven days
Central	Plumas (104.7 mm)	Morris (11.9 mm)
Eastern	Pinawa (93.7 mm)	Sprague (9.3 mm)
Interlake	Moosehorn (80.9 mm)	Clarkleigh (20.1 mm)
Northwest	Ashville (146.2 mm)	The Pas (46.8 mm)
Southwest	Boissevain (187.5 mm)	Waskada (13.0 mm)

Commodity Reports

Cereals

- Winter wheat and fall rye are both flowering, with the crop generally looking very promising, with strong yield potential. Many producers have already completed fungicide applications.
- Spring cereals are mostly between the tillering and stem elongation stages, with the earliest fields reaching early heading. Herbicide applications are largely complete, and producers are beginning to focus on fungicide decisions for fusarium head blight. Some cereal disease symptoms, including suspected tan spot, continue to be observed, with increased scouting recommended as crop canopies develop.
- Corn is generally between the V6 and V8 stages. Crop stands remain good across most of the region, although areas affected by excess moisture continue to show uneven growth and nutrient stress symptoms

Oilseeds

- Canola development has continued rapidly and is generally between the 6-leaf and flowering stages, with the earliest fields at 40-50% flowering. Many producers are scouting using a sweep net for cabbage seed pod weevil which has been observed feeding in large numbers in some fields on buds.
- Sunflowers have reached V8 stage.

Pulses & Soybeans

- Soybeans are at the 3rd to 4th trifoliolate stage. The soybeans have yellowed from the excess water, but most fields are beginning to grow out of stressed conditions.
- Peas are progressing well and have reached canopy closure in most fields. Peas are at the 10–12 node stage to just beginning to flower.

Forages & Livestock

Forages

- Excessive rainfall and saturated field conditions continue to delay forage harvest across much of Manitoba, with many producers waiting for extended periods of dry weather before resuming operations.
- Some producers took advantage of short weather windows to cut hay despite the risk of additional rainfall, as forage quality continues to decline with advancing maturity.
- First-cut harvest is progressing across the province, with dairy operations in some areas already preparing for or beginning second-cut silage, while beef producers continue first-cut harvest.
- Standing water and flooding have inundated hay land in several regions, delaying harvest, reducing forage quality, and in some cases preventing field access altogether.
- Hayfields in the Northwest Region, particularly around Dauphin, Ethelbert, and Ste. Rose, remain too wet for harvest and will require considerable drying before equipment can enter.
- Most established alfalfa, grass hay stands, and pastures have generally tolerated the excess moisture with limited stand damage; however, localized flooding has caused significant losses in low-lying areas.
- Forage yields are expected to be adequate overall, but forage quality may be reduced due to delayed harvest and challenging drying conditions. Many producers are opting to produce silage rather than dry hay because of persistent wet weather.
- Annual feed crop conditions remain variable across regions, with excess moisture negatively affecting growth in some areas. Some producers are considering seeding greenfeed on flooded acres if field conditions improve.
- Corn continues to progress well in many areas, although localized hail damage, particularly in the Boissevain–Killarney area, has resulted in severe crop losses in some fields.
- Herbicide and pesticide applications remain behind schedule due to saturated fields, resulting in increased weed and insect pressure in annual feed crops.

Livestock

- Pasture forage supplies remain abundant, with strong regrowth following grazing and adequate feed available across most regions.
- Water supplies for livestock are excellent, with dugouts, sloughs, and other water sources full or overflowing following persistent rainfall.
- Excess moisture on pastures continues to create management challenges, including muddy conditions, soil degradation, hoof damage in high-traffic areas, and increased insect pressure.
- Cases of foot rot are increasing in several regions due to prolonged wet conditions, and producers are encouraged to closely monitor cattle for signs of lameness and treat affected animals promptly.
- Significant fly pressure is being reported in many areas, adding to livestock management concerns.
- Flooding along rivers and waterways has forced some producers to relocate cattle or provide supplemental feed where grazing access has been reduced by standing water.
- In the Northwest Region, washed-out roads and river crossings have limited access to livestock, making routine welfare checks more difficult and delaying the movement of bulls to pasture for some operations.
- Highway closures in some areas temporarily disrupted feeder cattle deliveries for producers with pre-sold livestock, creating additional management pressure where feed supplies were becoming limited.
- Cattle marketings remain seasonally light, with several auction marts operating on reduced summer schedules or temporarily closed. Market conditions continue to remain favourable despite lower sale volumes.

Regional Comments

Southwest

The Southwest region is experiencing very wet conditions following storms over the weekend. Most areas south of Highway #1 received 2–7 inches of rain, on top of the heavy rainfall from the previous week. Some crops are completely submerged, while others are partially under water.

Significant hail was reported near Cromer, Dand, Boissevain, Killarney, Wawanesa, and north of Hartney. Minor hail, approximately the size of a loonie, was reported near Reston and Brandon. Boissevain received golf ball-sized hail, while Killarney experienced baseball-sized hail. Moderate lodging in cereal crops has also been observed in the Melita area.

Overall, most of the Southwest region has good crop potential if conditions dry out soon. However, some flooded areas are unlikely to recover, and those crops will be lost. Despite these losses, conditions are not as severe as those reported in areas to the north and east.

Most pea fields have filled in and are just beginning to flower. They are approaching the stage for fungicide application if required; however, disease pressure remains low, and growers may be able to delay application. Winter wheat and fall rye are headed and continue to look good, with low disease levels reported so far. Producers were able to spray most of these acres before the major rainfall events began.

Spring-seeded cereals are at the flag leaf to early heading stage. Producers are attempting to apply fungicides, but many fields remain too wet, resulting in rutting during ground applications. Spray planes are heavily booked, which may affect the timing needed for optimal disease control. Leaf disease pressure remains relatively low at this time, but the current warm and humid conditions could increase disease development. Spring wheat crop condition is at 50% good and 50% average.

Most canola fields are at the full cabbage to early bolting stage, with the earliest fields approaching 30% flowering. Fungicide application timing is becoming critical for effective disease control. Higher cabbage seedpod weevil populations have been noted in the southern portion of the region, although there have been few reports of spraying. Weed control applications are still ongoing in some later-seeded canola fields.

Soybean crops have generally handled the wet conditions well, although some iron deficiency chlorosis (IDC) is beginning to appear. Most soybean fields are currently at the V3 to early flowering growth stage.

Corn and sunflower crops are progressing at an average pace. They continue to grow, and warmer temperatures would be beneficial for further development.

Northwest

An intense weather event moved in the Northwest Region to start the week. Rain began Monday and continued into Tuesday bringing significant amounts across the entire region. The large amounts across the region overwhelmed ditches/drains and rivers very quickly, causing significant overland flooding and damage to infrastructure. In addition to the entire Northwest region receiving damage, the Swan Valley area was affected for a second time. Water levels have since receded; however many fields are still affected by overland flooding and damage continues to be assessed.

Infrastructure damage was significant in this storm, with all previous repairs in the Swan Valley region washed away again. Much more additional damage was received across the entire region, making commuting impossible, as well as accessing fields.

Although some fields received timely herbicide applications, there are many fields that have missed the application window due to unfavourable conditions, including wet fields, rain and high winds.

Fall rye and winter wheat are headed out and continue towards flowering stage.

Field peas continue in the vegetative stage and most advanced are now into the R1 stage. Areas that have been continuously wet are showing signs of excess moisture.

Crop stages continue to range in spring cereals, from tillering to flag leaf stage. Once again, areas under continuous excess moisture are showing symptoms of stress.

Canola fields also range in stages, from 4 leaf to the earliest fields now in early flower. While some crops look healthy, some fields under excess moisture are thin and starting to stress bolt. Herbicide applications may be challenging due to excess moisture.

Soybean fields range from the second to fourth trifoliate stage and some fields are showing signs of IDC.

Central

The region continued to receive localized storms over the past week, with all areas receiving rain but volumes varying greatly. Producers have mostly wrapped up in-crop herbicide applications as more crops approach canopy closure. Only a very small proportion of farms are expected to make a second pass of herbicides. Producers are focusing on fungicide decisions, and insect monitoring. Many producers are relying on aerial applicators, especially due to the wet field conditions this year.

Corn is progressing quickly and is generally between the V6 – V8 stages. Overall, the crop is looking good and healthy.

Soybeans are from V6, with most of the crop at or near R1. Crop development has accelerated with warmer temperatures. Soybeans are tolerating the wet conditions relatively well, although areas with prolonged saturation show reduced growth. IDC is noticeably visible in some fields, but the crop is mostly growing out of these early season issues.

Peas are flowering or very close to flowering and some producers have begun fungicide applications. Fall rye is at the soft dough stage, and winter wheat is at the late anthesis stage. Many producers have already completed fungicide applications.

The most advanced spring wheat is in anthesis, with some of the earliest crops just beginning to head. Producers are focusing on disease monitoring and fungicide decisions. Spraying for Fusarium Head Blight will begin in the next few days. There have been very few reports of leaf diseases to date, but producers continue to monitor for issues.

Canola development has continued rapidly and is generally between bolting stage to 50% flowering. Spraying for sclerotinia has begun and will become more common over the coming week. Many producers are expected to make fungicide applications given the recent rainfall events.

Many producers are scouting using a sweep net for cabbage seed pod weevil which has been observed feeding in large numbers in some fields on buds, with some insecticide spraying taking place. Cabbage seed pod weevil is typically more numerous in the earliest flowering fields as the weevil is attracted to yellow so moves into those fields.

Low numbers of grasshopper nymphs are being spotted in field margins. Numbers are typically lower in wetter years, but producers should continue to monitor for pest damage.

Eastern

The Eastern region saw rainfall this last week ranging from 9.3 mm to 93.7 mm. Field conditions remain wet as showers continue to move through the region. Producers are hopeful for a few good drying days before fungicide applications in both wheat and canola get fully underway.

Herbicide applications are wrapping up with a few second pass applications on corn remaining. Wheat has moved into anthesis, and fungicide applications are planned this week. Canola is in full bloom, with fungicide applications also planned for this week.

Fall rye fields have moved into the soft dough stage and are looking good. The earliest wheat fields are in full anthesis and are looking very good other than the field drain areas where crop loss has occurred. FHB fungicide applications are planned for this week.

Most corn is at the V7 to V8 stage. Final second pass herbicide applications are currently being completed. The crop looks good for the most part and has taken advantage of some of the warmer temperatures we have received.

Canola is at the bolting to 50% flower stage due to the long seeding window this spring. With the wet humid conditions, many producers are planning for a fungicide application to protect against sclerotinia; applications will begin this week.

Soybeans are at the first flower (R1) stage. Many fields have yellowed from the excess water, but most are outgrowing the stressed conditions. Sunflowers are delayed due to the wet conditions but are looking good for the most part.

Cabbage seed pod weevil continues to be a concern with assessments being made if economic thresholds are met to justify an insecticide application. Producers will assess on a field-by-field basis.

Interlake

Rainfall was significant this past week, with many areas experiencing saturation and flooding in low-lying spots. Many areas received substantial amounts of rain, along with some reports of hail. This consistent precipitation has led to saturated fields and flooding in low-lying areas across much of the South Interlake. The wet conditions are increasing the risk of root diseases in pulse crops and delaying herbicide applications. The cool and wet weather has also slowed crop development, complicating haying operations for farmers.

Winter wheat and fall rye are beginning to flower, demonstrating strong stands. While conditions for winter cereal and fall rye remain generally good, areas affected by excess moisture are exhibiting signs of stress and hail damage.

Spring cereals are advancing into the tillering and stem elongation stages, with the first fields showing visible flag leaves. Most herbicide applications are complete, and producers are now concentrating on disease monitoring and fungicide operations. Spring wheat ranges from good to excellent in some areas, while others are rated as poor to fair. Spring cereal crops on drier ground are thriving, with about a quarter rated as excellent.

Corn is progressing well, reaching the V3 and V5 stages. Crop stands are excellent in some locations, while others impacted by excessive moisture are struggling. There are signs of uneven growth and nutrient stress in waterlogged fields.

Canola is making good progress, currently at the 5-leaf and bolting stages, with the earliest fields already flowering. Many producers are actively scouting for cabbage seedpod weevil using sweep nets, as they are being observed in large numbers feeding on buds, prompting anticipated insecticide applications. Oilseed crops in the flood affected southern regions are generally in poor condition, while those in northern areas have largely maintained normal development, although cool, wet weather has caused some delays.

Sunflowers have reached the V5 stage.

Soybeans are currently in the 2nd to 4th trifoliolate stage, with the first round of herbicide applications mostly complete, leaving just a few second applications remaining. Some soybean fields are showing yellowing leaves due to excess moisture, while others are showing resilience. Peas are progressing well and are approaching canopy closure in most fields, currently at the 10-node stage. Among pulse crops, about 60% are in good condition, with the rest poor and showing signs of moisture stress.

Across the region, excessive moisture and persistent weed growth are the main concerns. Reports indicate moderate to severe damage from insects and strong winds. Flea beetles continue to be a problem, causing moderate damage, while minor issues from cutworms and cabbage seedpod weevils have also been noted.

Agronomy Topics

Assessing Hail Damage in Cereal Crops

Hail has affected crops in several areas of Manitoba over the last week. The amount of loss expected from a hail event depends on the severity of hail, crop type, and the growth stage of the crop. Assessments of damage should occur a few days after a hail event.

Wheat: Hail damage during jointing or in the boot stage is difficult to assess. Spikes can still pollinate and fill, and regrowth from new tillers can occur. The more advanced the wheat is at the time of hail the greater the yield loss. The greatest yield reduction from hail occurs in the milk stage.

Table 1. Grain yield reduction in spring wheat with 100 percent of stems bent (Derived from Busch, 1975)

Growth Stage	Yield Reduction (%)
Boot (Zadoks 45)	28-39%
Anthesis (Zadoks 65)	15-60%
Milk (Zadoka 75)	30-70%
Soft dough (Zadoks 83)	16-55%
Hard dough (Zadoka 87)	3-47%

Source: Hail Damage in Small Grain by Joe Lauer, Corn Agronomist, University of Wisconsin, May 18, 2007.

Oats and Barley: Oats and barley will tiller and recover better from hail than wheat, especially prior to the boot stage. Grower experience has demonstrated that barley, hailed severely in the boot stage, has recovered to produce 70-80% of normal yield. Crop hailed prior to the boot stage should be left if stems or green tissue remains.

Corn: Leaf loss without damage to the growing point has a small impact on yield. Check corn for damage to the stem, and percent leaf defoliation. Between the V6 stage and tasseling percent yield loss is estimated to range from 0-34% depending on percent defoliation. For more information: [How to assess corn after a hail event](#).