

Crop Report – September 22, 2026



[Crop Pest Report](#) [Crop Weather Report](#) [Reporting Area Map](#) [Seasonal Reports](#) [Weekly Weather Maps](#)

Inside this Report...

Upcoming Events
Weather Report
Commodity Reports
Cereals
Oilseeds
Pulses & Soybeans
Forages & Livestock
Regional Comments
Southwest
Northwest
Central
Eastern
Interlake

Upcoming Events & News

- Get the print version of **Seed Manitoba** delivered directly to your door! Sign up for the print edition and/or the Seed Manitoba newsletter here: <https://seedmb.ca/subscribe/>

Weather Report

- Rain this week brought variable precipitation accumulations across Manitoba. Most regions received more than 8 mm of rain. Parts of the Southwest, Northwest, and areas around the southern edge of Lake Winnipeg received more than 30 mm. The highest accumulation of precipitation in the past seven days was at Miniota with 77.8 mm.
- Climate normals for total accumulated precipitation from May 1 to September 20 range from 245 mm to 380 mm and are based on 30-year historical data. Since May 1, large areas of the Northwest and Southwest regions have accumulated more than 380 mm of precipitation. Seasonal accumulations are above normal for the majority of the Northwest and Southwest regions. Large areas of the Northwest, Southwest, and Interlake regions have seen accumulations over 130% of normal. A few localized areas of the Interlake, Central and Eastern regions have seen accumulations below 90% of the 30-year average.
- Total Accumulation of Growing Degree Days shows the accumulated growing degree days (GDD) for the period of May 1 – September 20, 2026. GDD accumulations ranged between 1349 and 1889 GDD so far this growing season. Most of southern Manitoba has accumulated more than 1600 GDD.

To find interactive soil temperature/moisture and air temperature information see Agri-Maps Current Weather [viewer](#)

Table 1. Range of measurements of seven-day accumulated precipitation in Manitoba's Agricultural Regions. (September 14 – 20, 2026).

Region	Wettest location last seven days	Driest location last seven days
Central	Plumas (41.6 mm)	Morden (9.1 mm)
Eastern	Lac Du Bonnet (34.6 mm)	Sprague Lake (12.8 mm)
Interlake	Lake Francis (40.9 mm)	Taylor's Point (20.1 mm)
Northwest	Drifting River (43.9 mm)	The Pas (1.6 mm)
Southwest	Miniota (77.8 mm)	Glenboro (22.5 mm)

Table 2: Percentage of Harvest Completion by Crop and Region to September 21, 2026.

Crop	Southwest	Northwest	Central	Eastern	Interlake	MB AVG
<i>Winter Wheat</i>	100%	100%	100%	100%	100%	100%
<i>Fall Rye</i>	100%	100%	100%	100%	100%	100%
<i>Spring Wheat</i>	45%	50%	98%	95%	92%	71%
<i>Barley</i>	70%	30%	98%	90%	92%	76%
<i>Oats</i>	50%	15%	96%	95%	90%	66%
<i>Field Pea</i>	100%	75%	100%	100%	100%	96%
<i>Canola</i>	20%	10%	75%	75%	45%	43%
<i>Soybean</i>	-	-	1%	1%	-	-
<i>Dry Bean</i>	-	-	20%	-	-	16%
<i>Silage Corn</i>	15%	-	10%	-	15%	10%
<i>Potatoes</i>	30%	-	30%	-	-	30%
Regional AVG	27%	24%	64%	62%	56%	43%

Crops still unharvested, or negligible acres displayed as – or omitted from this table.

Commodity Reports

Note: All yields are reported in comparison to 10-year (2015-2024) averages by region based on MASC published data. For reference, please refer to the [regional average tables by crop](#).

Cereals

- Winter cereal harvest is complete across the province, with yields ranging from slightly below average to above average.
- Spring cereal harvest is nearing completion in the Central, Eastern and Interlake regions. The majority of spring cereals are yielding average, but yields range from slightly below average to above average across the province. Quality had been good in earlier harvested crops, but downgrading due to sprouting and weathering is occurring in recently harvested fields.
- Corn is in the dent stage. Silage corn harvest has started.
- Winter cereals are growing well with the recent moisture.

Oilseeds

- Canola harvest is ongoing with yields reported as below average to average. Strong winds and rain have caused lodging in some fields. Sprouting, pod shatter and pod drop have been an issue in some areas.
- Sunflowers are in the R8 to R9 stage.
- Flax is at stage 12, with stem colour changing and leaves starting to drop. Preharvest herbicides have been applied.

Pulses & Soybeans

- Majority of soybeans range from R7 to R8. Harvest has started on limited acres in the Central and Eastern regions.
- Harvest has started on the earliest edible bean fields.
- Pea harvest is nearly complete, with the majority of yields reported as being average and good quality.

Forages & Livestock

Forage

- Frequent rainfall continues to dictate forage harvest progress, with field conditions limiting equipment access and delaying remaining hay and silage operations, particularly for slough hay and remaining second-cut forage.
- Third and fourth cut forage harvest is underway on dairy operations where conditions permit. Frequent rainfall and less-than-ideal harvest conditions are affecting forage quality in some operations.
- Producers are encouraged to sample stored forage, as it is harvested, to determine feed quality and ensure rations can be matched to livestock nutrient requirements.
- Straw is being baled and hauled home as it becomes available.
- Corn silage harvest has begun in areas where field conditions are suitable, with producers expecting average to above-average yields in most areas; however, in areas where plants were stressed, the yield is expected to be reduced. Other producers are waiting for fields to dry enough to support equipment traffic, while additional rainfall in the forecast may delay harvest progress.
- Pasture regrowth has been exceptional due to frequent rainfall, allowing producers to extend the grazing season further into the fall. Forage growth is beginning to slow as daylight hours shorten and temperatures cool.
- Dugouts remain near full capacity, with frequent rainfall providing continued recharge and maintaining good water quality. Many dugouts are approximately 100 per cent full and have been refreshed by recent rainfall.

Livestock

- Wet corral conditions are creating challenges for producers preparing to wean calves and for operations expecting incoming backgrounding cattle. Producers are monitoring pen conditions and preparing to minimize mud-related issues.
- Saturated pasture conditions are resulting in increased hoof traffic and some soil disturbance in areas receiving heavier use.
- Manure spreading continues where corrals are dry enough to access and field conditions allow.
- Most insect pressure has subsided, and reported cases of foot rot have decreased significantly.

Regional Comments

Southwest

Little to no harvest progress this past week due to rain and spotty showers. Water laying in fields makes travel difficult. Cool evenings and foggy mornings have resulted in dewy morning conditions. Producers with access to driers spent much of last week drying grain and trying to create room for wet grain, so that when they can get back in the field, they have places to store it. Some producers tested fields over the weekend, but fields and grain samples were very wet. In many areas standing and swathed crops are sprouting. Most areas are reporting severe lodging in all crops still to be harvested.

Winter cereal and field pea harvest is complete, with reports of average yields and good quality.

Spring wheat harvest is 40 to 45% complete, with average to above-average yields. Quality has been good in earlier-harvested crops, but downgrading is occurring in more recently harvested fields.

Barley harvest is approximately 70% complete, with average to above-average yields. Quality has started to decrease due to recent rains, and producers are reporting concerns about sprouting. Recently, harvested crops were being downgraded due to sprouting, colour, and low test weight, with much of the crop being graded as feed.

Oat harvest is approximately 50% complete. Oat harvest is limited as many fields are still green, and wet conditions are making producers reluctant to swath.

Canola harvest remains limited, at less than 25% complete. Reports indicate average yields and good quality. Several fields are being swathed, mostly due to severe lodging caused by disease. Several fields are also being sprayed. Harvest conditions are difficult due to lodging and wet fields.

Soybeans are in the early R8 stage, depending on the variety and seeding date. Some fields are at 95% maturity, while others remain green and are just starting to change colour.

Corn is at dent stage. Sunflowers are at R7 to R8.

Northwest

Precipitation at the beginning of the week held up harvest progress for most of the region. Low areas continue to hold water, creating challenging harvest conditions. Harvest was able to resume toward the end of the week and over the weekend. Heavy dews continue to narrow the harvest window. Sprouting is occurring in standing and swathed crops. While some post harvest operations are occurring as crops are harvested, some are waiting for drier conditions.

Desiccation and swathing of canola continues as stages are reached.

Winter cereal seeding continues, and with recent moisture and heat, crops are growing nicely.

Approximately 75% of the field pea crops is harvested. The remainder of the crop is ready to harvest or near harvest. Reported yields have been mixed, ranging from below average to average.

Spring wheat harvest is approximately 50% complete, and yields range from slightly below average to above average. Pre-harvest glyphosate applications continue in later spring wheat as stages are reached.

Canola fields continue to range in stages, with the majority of the crop nearing maturity. Desiccation and swathing is underway in fields that have reached the appropriate stage. Fields that experienced excess moisture have patches of delayed growth that are behind. Canola harvest is approximately 10% complete, with below average to average yields being reported.

Majority of soybeans are in the late R6 to R7 stage and are looking good where excess moisture didn't delay growth this spring. Stagey conditions are evident in some areas that experienced excess moisture through the season.

Central

Harvest progression varies considerably across the region. The area surrounding Portage la Prairie has been impacted by excess rainfall in September and much of the spring cereal and canola crop remains in the field. Delays in harvest has led to several issues.

In most of the region, spring cereal harvest is almost at completion. Much of the harvested grain requires additional drying. Many of the later harvested cereal crops have lodging issues, and quality declines, including pre harvest sprouting issues. Across the entire region, spring wheat yields and protein percentages range from average to good. Barley yields are slightly above average. Oat test weights, however, are lower than typical, likely due to lower soil moisture during critical development stages.

A small number of producers began soybean harvest early last week. Fields are at R7 (beginning maturity) through to R8. Due to dry and warm conditions earlier in the season, many plants have predominantly two-seeded pods, with very few four-seeded pods. Pod abortion has also occurred in the upper canopy in many fields. Many fields have areas of soybeans maturing at different rates, and this has made decisions regarding harvest timing difficult.

Pea harvest is complete. Quality has been good, while yields have been average, with a few fields that experienced more optimized soil moisture conditions showing above-average yields.

Edible beans are being cut and pre-harvest herbicide applications are being applied. The earliest fields have been harvested.

Canola harvest and swathing continue, with approximately 65% of the crop now harvested. Pre-harvest sprouting has been an issue in some fields. In general, yields have been lower than in recent years, likely a combination of early-season excess moisture and mid to later season dryness, hot and dry conditions at flowering, cabbage seed pod weevil, and verticillium stripe. Pod shattering and drop has been a common issue. Many fields have a strong carpet of volunteer canola.

Flax is at stage 12 (seed ripe), with stem colour changing and leaves dropping. Pre-harvest herbicides have been applied, and the earliest fields may begin to be harvested in the coming week if conditions allow. Overall, flax crops are good, aside from areas that had waterlogging early in the season, and there are a few patches within fields which will likely yield poorly.

Sunflowers are late R8 (back of the head fully yellow, bracts remaining green), with the crop likely to approach physical maturity in the next week (R9, bracts becoming yellow and brown).

Corn is at R5 (dent), with milk lines progressing 75% down the kernels. Corn silage harvest has begun, although high plant moisture and kernel integrity are concerns for this year's crop. The corn crop appears to have fared relatively well despite the dry conditions experienced earlier in the season. Blackbirds are feeding on cobs.

Other fall activities are well underway, including baling, harrowing, cultivating, and ditching. Fall cover crop seeding has finished. September 15th is typically considered the last opportunity for establishment.

Eastern

Rainfall across the region over the past week continued to delay harvest. Very little harvesting happened over the last week as field conditions remained wet with few good drying days.

Wheat harvest is estimated at 95% complete. Oat harvest is estimated at 95% complete while barley is at 90% complete. All spring cereal yields have been reported as average at best. Damage from drowned out areas of fields earlier this spring have brought yields down.

The majority of corn is in the dent stage of kernel development; on average the milk line is at the 75% mark. The crop is continuing to grow well as starch accumulates in the kernel. Growers agree that enough moisture has fallen to finish up the crop. Producers are hopeful the frost stays away as the corn continues to accumulate starch and mature.

Canola harvest is where the most progress was made across the region this past week. Harvest is estimated at 75% complete, with some farms finished. Yields are below average to average. Hot and dry weather during the flowering period is being blamed for the average to below average yields along with the water damaged areas from earlier in spring.

Soybeans are at the R7 to R8 stage. Fields this past week have had significant leaf drop as the crop continues to mature and dry down. Overall, fields are looking good. Many varieties have "seed-pod rattle" in the pods, most fields still need to dry down but harvest has started in some early maturing seed production field. Soybean harvest completion is at 1%.

Fall field work is ongoing in many areas. Many farms are able to use the harvest delay to get their stubble worked in, in preparation for next year's planting.

Interlake

Widespread rainfall over the past week has delayed harvest. Crop moisture is declining slower than expected. Excess moisture and heavy wind have caused lodging and concerns about grain quality. Yields and quality vary across the region due to differences in rainfall throughout the season. Frost in some areas has helped mature crops dry down.

Spring wheat harvest is 92% complete, with average to below average yields. Oat and barley harvest is 90% complete, with average to slightly above average yields. Strong winds and lodging have decreased yields in some later seeded fields.

Corn silage chopping continues. Yields vary with seasonal rainfall, while early reports range from average to slightly above average. Grain corn is at R5, with frost related leaf browning evident in some fields.

The majority of soybeans are at R7 to R8 and are dropping leaves. If weather permits, harvest will begin later this week. Pea harvest is nearly complete in most areas, with slightly above average yields.

Canola harvest is about 45% complete, with early yields ranging from below average to above average. Strong winds and rain reduced yields in some standing and swathed fields. Reseeded canola is performing better in areas that received timely early season rain.

Most sunflowers are at R8 to R9.

Fall rye and winter wheat have been seeded and are starting to emerge.