Assessment of walleye pollock in the eastern Bering Sea

James Ianelli, Sophia Wassermann, Abigail McCarthy, Carey McGilliard, Elizabeth Siddon, Kalei Shotwell, Grant Adams, Nathan Lauffenburger

Alaska Fisheries Science Center, NOAA Fisheries

SI-assisted development

Tools assisted data processing, code execution, figure preparation and report writing under author direction. The authors retain responsibility for the scientific content, analyses, interpretations and conclusions.

Suggested citation

Ianelli, J., S. Wassermann, A. McCarthy, C. McGilliard, E. Siddon, K. Shotwell, G. Adams, and N. Lauffenburger. 2026. 2026 Assessment of Eastern Bering Sea Walleye Pollock.

https://files.npfmc.org/SAFE/2026/EBSpollock.pdf

Contents

1 Executive summary
2 Introduction
3 Fishery and management history
4 Data
5 Analytic approach
6 Results
7 Projections and harvest recommendations
8 Risk table and ABC recommendation
9 Status determination
10 F limit
11 Ecosystem considerations
12 Data gaps and research priorities
13 Acknowledgements
14 Assessment source archive
Literature cited
Tables
Figures
Appendices
Appendix A SSC and Plan Team Requests and Responses
Appendix B Ecosystem and Socioeconomic Profile of Eastern Bering Sea Walleye Pollock
Appendix C Research and Other Noncommercial Pollock Removals
Appendix D Detailed Current-Model Estimates
Appendix E Proposed Projection Input Curves
Appendix F Model Equations
Appendix G FT-NIRS Age Prediction for EBS Pollock
Appendix H BTS Spatiotemporal Biomass Index and Abundance at Age, Including the Northern Bering Sea
Appendix I Assessment Development and Supporting Record
Appendix J Contents

1 Executive summary

The last full EBS pollock assessment was completed in 2024 (Ianelli et al. 2024). A catch report was prepared in 2025 (Alaska Fisheries Science Center, National Marine Fisheries Service 2025). Changes in this assessment are described relative to 2024.

This executive summary presents Model 26.0 (Rceattle), the base case for this assessment and its Tier 3 harvest recommendations. Model 23.0 (ADMB), accepted in 2024, provides the comparison with the previous assessment.

1.1 Changes in assessment inputs

Updated annual catch estimates from 1991 onward, including 1,312,504.58 t for 2026.

Added 2025–2026 bottom-trawl survey (BTS) indices and age compositions, the 2026 acoustic-trawl survey (ATS) index and provisional age composition, and acoustic vessel-of-opportunity (AVO) indices through 2026. Historical BTS estimates were also revised.

Added fishery age compositions for 2024–2025, with year-specific conversions to traditional microscopic ages and likelihood sample sizes of 578 and 530, respectively.

Updated fishery, population and spawning weight-at-age schedules. The 2026 ATS ages use the BTS age–length key; FT-NIRS ages from the ATS samples are expected in 2027.

Sampling and conversion methods are described in Section 4.

1.2 Changes in assessment methods

The changes from Model 23.0 to Model 26.0 are:

Implemented the age-structured assessment in Rceattle in place of ADMB. The development comparisons are documented in Appendix I (Ianelli et al. 2026).

Estimated weight at age in a separate random-effects model and used the resulting schedules as fixed inputs to the assessment. Model 23.0 estimates cohort and year effects on weight at age within the assessment. The Model 26.0 uncertainty estimates therefore condition on the estimated weights (Section 5.2).

Natural-mortality and catchability assumptions retain the previous assessment's treatment.

1.3 Summary of results

Estimated total biomass in 2026 is 12.162 million t (approximate 95% interval 9.534–15.515 million t). Female spawning biomass (SSB) is 3.057 million t (2.489–3.754 million t). The largest fishing mortality rate among ages is 0.3 per year. These estimates and ranges condition on the base model and fixed biological inputs (Section 6.1.1).

Future catches are assumed to equal the specified annual TAC of 1.375 million t in 2027 and the same amount in 2028. The 2028 value is a planning assumption. Projections advance the fitted population using mean 2023–2025 fishery selectivity, fixed biological inputs and recruitment equal to the fitted 1978–2023 mean (Section 7).

The advice table compares the 2025–2026 recommendations from the 2024 assessment with this assessment's 2027–2028 recommendations. Projected female spawning biomass and ages-3+ biomass use Scenario 2 with the assumed catches above.

The final specifications set the 2026 and 2027 Bering Sea subarea pollock OFL at 2,496,000 t and ABC at 2,036,000 t; specified annual TAC is 1,375,000 t. These are the adopted specifications, separate from the recommendations from this assessment.

A risk table was evaluated, and all categories were determined to be Level 1. This supported recommending the maximum ABC (Section 8).

Model 26.0 advice from revised fits				
Quantity	As estimated or specified in 2024 for:		Primary baseline Model 26.0 for:	
	2025	2026	2027	2028
Natural mortality (annual rate, ages 3+)	0.300	0.300	0.300	0.300
Tier and sub-tier	3a	3a	3a	3a
Projected total biomass (ages 3+, t)	8,526,000	8,075,000	11,270,000	12,521,000
Projected female spawning biomass (t)	3,118,000	3,342,000	3,326,000	3,670,000
B100% (t)	5,902,000	5,902,000	6,234,000	6,234,000
B40% (t)	2,361,000	2,361,000	2,493,000	2,493,000
B35% (t)	2,066,000	2,066,000	2,182,000	2,182,000
F_OFL (annual rate)	0.513	0.513	0.688	0.688
Maximum F_ABC (annual rate)	0.394	0.394	0.534	0.534
Recommended F_ABC (annual rate)	0.394	0.394	0.534	0.534
OFL (t)	2,957,000	2,496,000	2,746,000	2,870,000
Maximum ABC (t)	2,417,000	2,036,000	2,222,000	2,319,000
Recommended ABC (t)	2,417,000	2,036,000	2,222,000	2,319,000
Historical source: December 2024 SAFE, Executive Summary, Tier 3 version, p. 4; B40% is printed in section 7.2, p. 35.				
2027–2028 projections assume catch of 1,375,000 t in each year, mean 2023–2025 selectivity and fixed biological inputs. Total biomass covers ages 3+ at the start of the year; spawning biomass is female biomass at spawning. F_OFL and maximum F_ABC are the fully selected F35% and F40% rates. Biomass and catch are rounded to the nearest 1,000 t.				
Spawning biomass exceeds B40% in both years, placing the stock in Tier 3a. All risk categories are Level 1, supporting the maximum ABC. Recruitment is fixed at the fitted 1978–2023 mean; these projections show point estimates. Adopted catch specifications are reported separately.				

Model 26.0 stock-status summary. The 2025–2026 columns report status for 2025 and 2026; historical determinations are from the 2024 assessment.				
Status	As determined in 2024 for:		As determined this year for:	
	2023	2024	2025	2026
Overfishing	No	n/a	No	n/a
Overfished	n/a	No	n/a	No
Approaching overfished	n/a	No	n/a	No
Historical determinations are from the December 2024 EBS pollock SAFE, Executive Summary, p. 4.				
The 2025–2026 entries are assessment conclusions, separate from final NMFS status determinations and SSC/Council review. Overfishing for the most recent complete year compares the updated 2025 catch of 1,326,215.61 t with the specified 2025 Bering Sea subarea OFL of 2,957,000 t. The 2026 overfished determination uses each model's estimated 2026 female spawning biomass; approaching status uses its Scenario 7 projections through 2028 (Section 9). Model 26.1.2 is the Early-late fishery split comparison. n/a denotes a year/status combination outside this reporting layout.				

1.4 SSC and Plan Team requests

The September 2026 Plan Team requested the following work:

Present Model 26.0 for management use and Model 23.0 for continuity; both are compared in this assessment.

Examine age-1 BTS selectivity and the influence of BTS variability on retrospective patterns. Additional selectivity and data-influence tests are outside the scope of this assessment.

Extend the simulation self-test and examine retrospective patterns in time-varying selectivity. The completed simulation experiment and its limitations are described in Section 6.3.1.

Present retrospective results for Models 23.0 and 26.0 side by side. Matched runs were completed and compared; both models showed similar spawning-biomass retrospective patterns (Appendix A).

Document age-composition weighting and evaluate residual patterns. Selected sample sizes, one-step-ahead residuals and Francis comparisons are presented in the assessment.

Continue development of the Ecosystem and Socioeconomic Profile; we developed a draft using AI to compile and coordinate information from disparate sources (Appendix B).

Earlier SSC requests address model development, uncertainty, survey treatment and composition weighting. The individual requests and responses are presented in Appendix A.

2 Introduction

2.1 General

Walleye pollock (*Gadus chalcogrammus*; hereafter referred to as pollock) are broadly distributed throughout the North Pacific with the largest concentrations found in the eastern Bering Sea (EBS). Also known as Alaska pollock, this species continues to play important roles ecologically and economically.

Changes in assessment inputs and methods since 2024

The 2026 assessment updates the accepted December 2024 assessment; the 2025 document was a catch report. Data changes include revised annual catches from 1991 onward, fishery age compositions for 2024-2025, BTS indices and age compositions for 2025-2026, the 2026 ATS index and provisional age composition, and AVO indices through 2026. Historical BTS estimates were also revised. Spatiotemporal estimation of survey abundance continues the approach adopted in 2024; the current biomass-index and abundance-at-age products use sdmTMB and tinyVAST, respectively.

Fishery age compositions incorporate year-specific conversions from FT-NIRS ages to the traditional microscopic-age scale. Fishery, population, and spawning weight-at-age schedules have also been updated. The provisional 2026 ATS age composition applies the BTS age-length key to ATS lengths; FT-NIRS ages from the ATS samples are expected in 2027. Section 4 and Appendices G-H describe these data treatments and their uncertainty.

The main assessment-method change is the transition from the ADMB implementation of Model 23.0 to the Rceattle implementation of Model 26.0. Weight at age is estimated in a separate random-effects model and supplied as fixed schedules, whereas Model 23.0 estimates cohort and year effects on weight at age within the assessment. Model 26.0 uncertainty therefore conditions on the estimated weights. Natural-mortality and catchability assumptions retain the previous treatment. The configuration comparisons, diagnostics, and sensitivities evaluate the effects of the data updates and the implementation change (Sections 5-6 and Appendix I).

2.2 Review of Life History

In the EBS pollock generally spawn during March-May and in relatively localized regions during specific periods (Bailey (2000)). Generally spawning begins nearshore north of Unimak Island in March and April and later near the Pribilof Islands (Bacheler et al. (2010)). Females spawn in batches with up to 10 batches of eggs per female per year (during the peak spawning period). Eggs and larvae of EBS pollock are planktonic for a period of about 90 days and appear to be sensitive to environmental conditions. These conditions likely affect their dispersal into favorable areas (for subsequent separation from predators) and also affect general food requirements for over-wintering survival (Gann et al. (2015), Heintz et al. (2013), Hunt Jr. et al. (2011), Ciannelli et al. (2004)). Duffy-Anderson et al. (2016) provide a review of the early life history of EBS pollock.

Throughout their range juvenile pollock feed on a variety of planktonic crustaceans, including calanoid copepods and euphausiids. In the EBS shelf region, one-year-old pollock are found throughout the water column, but also commonly occur in the NMFS bottom trawl survey. Ages 2 and 3 year old pollock are rarely caught in summer bottom trawl survey gear and are more common in the midwater zone as detected by mid-water acoustic trawl surveys. Younger pollock are generally found in the more northern parts of the survey area and appear to move to the southeast as they age (Buckley et al. (2009)). Euphausiids, principally *Thysanoessa inermis* and *T. raschii*, are among the most important prey items for pollock in the Bering Sea (Livingston (1991); Lang et al. (2000); Brodeur et al. (2002); Ciannelli et al. (2004); Lang et al. (2005)). Pollock diets become more piscivorous with age, and cannibalism has been commonly observed in this region. However, Buckley et al. (2015) showed spatial patterns of pollock foraging varies by size of predators. For example, the northern part of the shelf region between the 100 and 200 m isobaths (closest to the shelf break) tends to be more piscivorous than pollock found in more near-shore shallow areas.

2.3 Stock structure

Stock structure for EBS pollock was evaluated in Ianelli et al. (2015). In that review past work on genetics (e.g., Bailey et al. (1999), Canino et al. (2005)) provided insight on genetic differentiation. The investigation also compared synchrony in year-classes and growth patterns by region. Pollock samples from areas including Zhemchug Canyon, Japan, Prince William Sound, Bogoslof, Shelikof, and the northern Bering Sea were processed and results presented in Ianelli et al. (2021). Genetics research at the AFSC (Ianelli et al. 2023) indicated that genetic groups mostly aligned with current stock-management delineations. the Aleutian Islands and Bogoslof “stocks” showed subtle differentiation from the Gulf of Alaska management group.

3 Fishery and management history

Directed foreign fishing began in 1964. US vessels entered the fishery in 1980, and it became fully domestic by 1988. The domestic observer program formally began in 1990; later monitoring developments include electronic monitoring on participating catcher vessels and biological sampling during offloading (Ianelli et al. 2024). The directed fishery primarily uses pelagic trawl gear, with inshore, catcher/processor and mothership sectors. The historical catch and adopted harvest-limit series is continued through 2026 in Table 1.

Updated reported catches are 1,311,261 t in 2024 (2025 catch report) and 1,326,193 t in 2025 (National Marine Fisheries Service Alaska Region 2026b). The two 2026 rows distinguish an official catch of 1,297,206 t through October 17 (National Marine Fisheries Service Alaska Region 2026a) from the annual catch estimate of 1,312,504.58 t used in this assessment.

For 2026, adopted ABC and OFL are 2,036,000 and 2,496,000 t. The initial Bering Sea subarea TAC of 1,375,000 t became 1,389,500 t after inseason reallocations; the August allocation table specifies 139,400 t for the Community Development Quota (CDQ) directed fishing allowance and 46,000 t incidental catch allowance. CDQ directed catch and the non-Bogoslof incidental account are each included once in the reported Bering Sea subarea total; the latter already includes incidental CDQ catch. Bogoslof and Aleutian Islands accounts remain separate (National Marine Fisheries Service 2026a, 2026c).

Continuing 2026 measures allocate 45% of Bering Sea subarea directed allowances to the A season (January 20–June 10) and 55% to the B season (June 10–November 1). The non-CDQ directed allowance is divided 50%, 40% and 10% among the inshore, catcher/processor and mothership sectors. The Steller sea lion conservation area limit before noon April 1 is 28% of each sector's annual directed allowance (National Marine Fisheries Service 2026a). These regulatory seasons differ from the week-ending month groups used in several historical fishery displays. An additional 2026 spatial action closed the Winter Herring Savings Area (58–60° N, 172–175° W) to directed pollock trawling, including CDQ, from noon September 30, 2026 to noon March 1, 2027 (National Marine Fisheries Service 2026b).

Management has combined area restrictions, allocations, monitoring and incentives to avoid prohibited species. Selected historical milestones are summarized in Table 2; Council recommendations, voluntary programs and regulatory implementation retain their separate dates. The voluntary rolling-hotspot programs began for chum salmon in 2001 and Chinook salmon in 2002. The Council selected Amendment 84 in 2005, followed by exempted fishing in 2006 and implementing regulations in 2007 (National Marine Fisheries Service 2007, 2023).

Amendment 91's rule became effective on September 29, 2010, and its Chinook bycatch program operated in the 2011 fishery. It combined a 60,000-fish overall prohibited species catch (PSC) limit with incentive plan agreements (IPAs) and sector shares of a 47,591-fish performance standard. A sector retains the higher allocation if it exceeds its share of the standard in at most two of seven consecutive years. Exceeding the standard in three of those years triggers the lower allocation in all subsequent years. Amendment 110 followed Council final action in April 2015 and a final rule effective July 11, 2016. It strengthened Chinook and chum avoidance requirements and linked the following year's Chinook limits to the State of Alaska's index for the Kuskokwim, Unalakleet and Upper Yukon systems. At an index of 250,000 fish or less, the PSC limit and performance standard fall to **45,000 and 33,318 fish**, respectively. The index first informed the 2017 fishing year. The pollock fishery uses the **45% / 55% pollock seasonal allocations** described above (National Marine Fisheries Service 2010, 2016).

Salmon accounting and genetic stock identification serve different purposes. The Bering Sea pollock program requires census counts of salmon in catches, with observer verification at at-sea processors or during offloading. Participating catcher vessels may use compliance electronic monitoring (EM), while observers at shoreside processors collect biological samples. EM verifies handling and discard compliance. Genetic origin is determined from biological samples. Representative biological samples, rather than every counted fish, are genetically analyzed to estimate stock composition. These are fishery-dependent records, separate from the assessment's fishery-independent survey indices (National Marine Fisheries Service 2010). Compliance monitoring and shoreside sampling are described by NOAA Fisheries (National Marine Fisheries Service Alaska Regional Office 2025). The genetic analysis of the 2015 Chinook salmon bycatch illustrates this subsampling approach (Guthrie et al. 2017).

3.1 Fishery conditions

The A-season fishery targets pre-spawning pollock and supports roe production. Fishing has historically concentrated north and west of Unimak Island and along the shelf toward the Pribilof Islands, with access affected by ice, weather and fish distribution. The summer–fall B-season fishery follows changes in fish availability across the shelf. Salmon-avoidance measures and vessel choices also influence where fishing occurs. The assessment authors report good fishing conditions in the 2026 A season; later in the B season fishing moved farther north and pollock of desirable sizes were less available.

Catch by area and season is summarized in Table 3. The week-ending extract records 1,267,543 tons through October 17, 2026: 604,041 tons in the A-season grouping and 663,501 tons in the B-season grouping. These seasonal groupings follow the reporting weeks described in the caption.

The annual catch-accounting extract summarizes retained and discarded catch in Table 4. The 2026 total is 1,312,505 tons, including 8,997 tons discarded (0.7%). The annual and seasonal extracts have different coverage and update schedules. Only the 1998 records included catch without an area; this catch is allocated between NW and SE in proportion to reported catch within each disposition.

Nominal pollock catch per tow-hour is generally higher in the A season than in the B season, with A-season peaks in 2019 and 2023 (Figure 1). Species-specific patterns differ: A-season Chinook rates peak in 2006–2007, whereas the chum salmon series has its largest B-season rates in 2005 and 2017. Herring and sablefish each have a pronounced peak in 2020, in the A and B seasons respectively. These are nominal rates for available pollock-dominated EBS hauls, calculated from summed catch and valid tow-hours with panel-specific units. Qualifying CV records are absent in 2025–2026 under the selection filter, sablefish has a reporting gap in 2014, and 2026 B season is partial through September 18.

Estimated catch numbers for both sexes decline to a trough in 2009, rise again during the 2010s and reach their largest annual totals in 2021 (Figure 2). B-season numbers generally exceed A-season numbers within each sex, and female annual catch numbers decline from 1,075.8 million fish in 2023 to 916.7 million in 2025, while male catch numbers decline from 1,218.9 million to 952.6 million. Catch numbers are expressed in thousands of fish. The 2024–2025 estimates use FT-NIRS ages and remain provisional.

Cumulative catch through April stays near 40% of annual specified TAC over much of 2000–2017 and near 44% from 2018 onward (Figure 3). In 2026 the cumulative shares are 6.69%, 23.14%, 39.42% and 43.72% at the end of January, February, March and April, respectively. Monthly totals are based on weekly catch reports and the specified annual TAC before quota transfers. These observations cover the first four months of the year; the A season continues through June 10.

A-season roe production exceeds B-season production throughout the 2000–2026 series (Figure 4). A-season production peaks in 2006 and again in 2019, declines through 2021 and rises from 2022 to 2026. The latest A-season value is 19,092.9 t of roe. The 2025 B-season total is 1,795.8 t; the 2026 total of 406.4 t covers June–October and represents a partial-season estimate. The series combines offshore Bering Sea and inshore Bering Sea and Aleutian Islands (BSAI) records in the original January–May and June–December groups.

The broader BSAI fishery's catch, vessel participation and market context are summarized in the ESP socioeconomic tables. Those statistics retain their BSAI, national-trade and global-catch scopes and their latest available observation years.

3.1.1 Recent spatial catch patterns

The seasonal maps compare 2024–2026 using the available catch-location records (Figure 5; Figure 6). A-season maps cover January 20–June 9. B-season maps use June 10–September 18 in every year, matching the 2026 data cutoff. Vertical bar heights show recorded catch tons. The rotated, elevated view separates neighboring columns. Each season uses a fixed vertical scale across its three years, with a 10,000-t reference column in each panel. The vertical scales differ between seasons; compare catch amounts using the reference columns.

A-season fishing remained concentrated on the southeastern shelf, with more catch extending westward in 2025. The share of available observed catch west of 170°W increased from 15% in 2024 to 39% in 2025, then returned to 15% in 2026. The catch-weighted mean latitude changed from 56.07°N to 56.43°N and 56.19°N, respectively. The 2026 pattern therefore shifted back toward the southeastern grounds relative to 2025.

The B-season distribution extended farther north and west in 2025–2026 than in 2024. The share of recorded catch north of 58°N increased from 54% in 2024 to 75% in 2025 and 79% in 2026. The share west of 170°W rose from 76% to about 92% in both later years. Catch-weighted mean latitude was 57.66°N in 2024, 58.56°N in 2025 and 58.52°N in 2026. Thus, the recent northward shift persisted in 2026, while the catch-weighted mean longitude moved eastward relative to 2025, from 174.65°W to 173.26°W.

These summaries describe the available observed catch, including catcher-processors, motherships and observed catcher vessels. The extract omits electronic-monitoring catcher-vessel records, so changes in fleet coverage can also affect the mapped patterns. The maps show catch concentration; the updated fleet-dispersion index below measures the separation among observed tows.

3.1.2 Fleet dispersion

Fleet dispersion describes how widely separated fishing locations are within a day (Figure 7). Following the 2024 assessment, we calculate the great-circle distance between each unique pair of tows made on the same day, separately for catcher-processors and observed catcher vessels. Distance sums and pair counts are then pooled by year and season. Days with more tow pairs contribute more to the seasonal average. Larger values indicate a more dispersed pattern of observed fishing locations; smaller values indicate greater concentration.

A-season dispersion increased from 47.8 nautical miles in 2024 to 68.8 in 2025, then declined to 36.5 in 2026. This change is consistent with the 2025 westward extension and the more concentrated southeastern pattern in 2026 shown in the catch maps. For B-season records through September 18 in each year, dispersion declined from 78.5 nautical miles in 2024 to 67.8 in 2025 and 56.5 in 2026. Thus, the observed B-season fishing locations became more concentrated even as much of the catch remained farther north than in 2024. A shift in fishing location and a change in spatial spread describe different aspects of fleet behavior.

The figure retains the historical day-of-year boundary at day 161, which approximates the seasonal transition. Dashed lines provide a common September 18 cutoff to support comparisons with the partial 2026 B season. The full series uses consistent duration calculations and vessel-type selections. Catcher-vessel coverage is incomplete because electronic-monitoring records are absent. Changes in the mix of observed vessels, bycatch avoidance and fishing conditions can influence this index, so it should be interpreted together with catch distribution and catch rates.

3.1.3 Seasonal changes in fish size

Following the approach described in Section 3.1.1 of the 2024 assessment, we summarize the mean weight of pollock within each source record and weight that record by its pollock catch. Mean fish weight is the recorded pollock weight divided by the corresponding fish count. Grouping these record means into weight bins shows how the sizes encountered by the fishery change over time (Figure 8). These distributions describe mean fish weights for each observer record.

The dominant A-season mean-weight bin increased from 550 g in 2023 to 600 g in 2024–2025 and 700 g in 2026. For the B season, the dominant bin increased from 450 g in 2023 to 550 g in 2024 and 700 g in 2025, then shifted to 450 g in the available 2026 records. This recent contrast is consistent with the reported difficulty finding larger pollock during the 2026 B season. Changes in these distributions can reflect growth, recruitment, the areas fished and the mixture of vessels represented in the observations.

Weekly displays show how size patterns develop within the A-season and B-season windows (Figure 9; Figure 10). The source extract extends through September 18, 2026, so the B-season results describe the available portion of the season. Catcher-vessel observations exclude electronic-monitoring records and represent only part of the shoreside fleet. Annual panels use a source-week boundary at week 24; the weekly panels cover weeks 2–15 and 25–40. These plotting windows differ from the regulatory season dates. Ridge heights are scaled to reveal distribution shape; catch totals should be compared using the underlying bin totals.

4 Data

Fishery catch data begin in 1964, whereas the BTS began in 1982 and the ATS in 1994 for the purposes of this assessment (see the table below). Available fishery age compositions end in 2025, one year before the requested 2026 catch and survey updates. Survey gaps and provisional age products retain their qualifications below.

Prohibited-species bycatch is summarized in Table 54. Chinook catches increased from 8,054 fish in 2024 to 19,825 in 2025, and non-Chinook catches increased from 35,152 to 151,457 fish. Halibut mortality rose from 53 to 66 t. Preliminary 2026 catches through October 24 were 15,153 Chinook, 149,468 non-Chinook, and 56 t of halibut mortality. Other bycatch summaries appear in Tables 55-57.

Source	Type	Years
Fishery	Catch biomass	1964–2025; 2026 year to date and projected annual input*
Fishery	Catch age composition	1964–2025†
Fishery	Historical trawl CPUE	1965–1976
Bottom-trawl survey	Biomass index and age-specific proportions	1982–2019, 2021–2026†
Acoustic-trawl survey	Biomass index and age-specific proportions	1994, 1996, 1997, 1999, 2000, 2002, 2004, 2006–2010, 2012, 2014, 2016, 2018, 2020, 2022, 2024, 2026†
Acoustic vessels of opportunity (AVO)	Relative backscatter index	2006–2019, 2021–2026†

**Annual catch inputs from 1991 onward use the catch estimates. Its 2026 total is 1,312,504.58 t; the within-year coverage cutoff is unspecified. †Available update observations are shown; the reference baseline ends in 2024 (fishery ages 2023); Model 26.0 uses the updated observations and selected provisional schedules, with source review retained, including the revised EBS/NBS BTS estimates. The 2020 acoustic survey used USV transects; age-specific proportions are unavailable. Provisional 2026 ATS ages use the BTS age–length key (ALK) applied to ATS population length-frequency estimates. FT-NIRS ages for the 2026 ATS samples will be available in 2027.*

FT-NIRS age estimation and survey developments are documented under age compositions, the FT-NIRS appendix, and the ATS, AVO and BTS methods sections. The assessment retains the fixed natural-mortality schedule described in Other data. Spatial catch and bycatch information helps evaluate changing distributions and guide avoidance. The assessment retains the established stock boundaries.

Model 26.0 uses observations through 2026 where available, an end-year forecast for 2026 catch, and fishery ages through 2025 (Table I1). The register identifies each source and observation year: fishery ages 2024–2025, annual survey products 2025–2026, and the scheduled 2026 ATS observation. Historical revisions, carried weights and forecast schedules have separate record types. It also distinguishes intake into the completed Model 23.0 increments, Model 23.2 bridge and selected Model 26.0 fit. Scientific acceptance remains a separate status.

4.1 Fishery catch and age compositions

Annual catch rose from about 0.175 million t in 1964 to the series maximum of 1.875 million t in 1972 (Figure 11). Catch subsequently fluctuated, fell to 0.811 and 0.810 million t in 2009 and 2010, respectively, and recovered to 1.326 million t in 2025. The 2026 records total 1.268 million t through October 17 and represent a partial year. Records before 1991 provide annual totals; later records distinguish the A-season and northwest and southeast B-season catches.

Fishery catch-at-age estimates for 1964–2025 are summarized in Table 5. Ages 15 and older are pooled as 15+. Where complete annual catch numbers are unavailable, we calculate numbers at age from the assessment proportions, annual fishery weight at age and catch biomass.

The annual assessment catch includes retained and discarded commercial pollock across gears and targets in the Bering Sea, with CDQ counted once and the separately reported Bogoslof and Aleutian Islands catches excluded. The catch update uses the catch estimates for 1991–2026, with pre-1991 model inputs retained. The revised totals are 1,311,260.74 t in 2024, 1,326,215.61 t in 2025 and 1,312,504.58 t in 2026. The 2026 estimate is the annual input selected for this assessment. Explicit Bogoslof (518) and outside-stock area codes are excluded. The 1998 records with blank area codes remain included under their Bering Sea classification. Noncommercial removals remain separately documented in Appendix C.

We estimate fishery age compositions from biological samples collected by at-sea observers and during shore-side offloading. The sampling records include fish measured for length frequencies, paired weight–length measurements, and age estimates (Table 6; Table 7; Table 8). The processing keeps recent unsexed fish separate rather than assigning them to males or females. Within each sex and sampling stratum, we expand length frequencies to the associated catch biomass using mean weight at length. Age–length keys allocate the expanded numbers to ages; we then sum the numbers

across strata and sexes and normalize them to give annual proportions (Kimura 1989; Barbeaux et al. 2005; Ianelli et al. 2024). Earlier assessments provide the pre-1991 age compositions. For the 2024–2025 update, the processing strata are January–May across Bering Sea areas, followed by June–December split into northwest (NMFS reporting areas 520 and above) and southeast (areas below 520). The 2024 assessment describes January–June and July–December groups; the current sampling groups use the calendar periods defined above. The new length and age records cover all 12 months. Unsupported age–length cells leave approximately 0.495% and 0.358% of expansion-input biomass unallocated in 2024 and 2025, respectively; these are sampling-support limits, rather than missing months or deductions from assessment catch.

The linked bootstrap represents variation among sampling units and among length and age observations within those units. Its composition precision estimates inform the selected likelihood sample sizes: 443, 578 and 530 for 2023, 2024 and 2025, respectively. They represent effective information for fitting the annual proportions, rather than numbers of fish measured or aged (Section 4.3.1). The model fits annual sex-combined proportions at ages 1–14 and 15+. Fishery age observations end in 2025.

The 2024–2025 age-estimate data are from pelagic trawl fisheries. For 2025, FT-NIRS predictions from model a00-d3 are expressed on an independent traditional microscopic-ageing (TMA) basis using 668 paired estimates (Appendix G). Within each FT-NIRS age, the paired observations define the proportions assigned to each tester age. These probabilities redistribute the catch-expanded numbers among ages while preserving their total. Source age 13 lacks paired tester observations; its conditional probabilities are the midpoint of the age-12 and age-14 rows. The 2025 source and tester ages range from 2 to 15, and the assessment fits ages 1–14 and 15+. The likelihood uses the selected input sample size of 530. Uncertainty in the conversion key and the age-13 interpolation is additional to the sampling uncertainty represented by that input weight.

The 2024 fishery age composition is expressed on the independent TMA basis using 693 paired reader–tester observations. For source age 13, the key pools TMA-minus-FT-NIRS age offsets from the same year's source ages 9–12 (34 pairs) and adds those offsets to age 13. Empirical key rows retain their observed probabilities. The conversion preserves total numbers; the selected likelihood input remains 578. Key uncertainty is additional to the sampling precision represented by that input weight.

The 2024 fishery weights use the same year-specific conversion and pooled-age rule, with the documented 100% CV for the 15+ weight observation. The 2025 weights use the same tester-age key and age-13 midpoint as the 2025 age composition. Appendix G describes these year-specific methods. The fishery age-age keys are separate from the BTS key.

4.2 Survey data

4.2.1 Bottom-trawl survey

The AFSC bottom-trawl survey provides indices of pollock abundance and population age structure, complementing the acoustic-trawl survey's sampling of the midwater component. Historical and recent sampling effort is summarized in Table 9. Annual surveys began in 1979, with standardized gear and methods from 1982 (Ianelli et al. 2024). The 2025 and 2026 observations extend the 42 survey years available for the 2024 assessment to 44 observed years during 1982–2026. The survey was absent in 2020; that year remains a gap in the observation series. The biomass estimates and assessment CVs are listed in Table 10. Availability to bottom-trawl gear varies with age and vertical distribution. Ages 2–3 are often relatively scarce in the catches because juveniles tend to occupy the water column (Monnahan et al. 2021; O'Leary et al. 2022).

Bottom temperature appears to influence the distribution of pollock available to the bottom-trawl survey. The AKsf survey-mapping example (Ianelli 2026) places annual CPUE distributions alongside the 0, 2 and 4 °C bottom-temperature contours for 2018, 2019 and 2021–2026 (Figure 12). High CPUE is concentrated along parts of the western and southern shelf, with several dense patches near the 2 °C contour in 2022, 2024 and 2026. The more extensive warm-water conditions in 2018–2019 contrast with the broader areas of cold bottom water in later panels, while the locations and extent of the pollock concentrations also change.

These patterns suggest that temperature helps structure near-bottom habitat and survey availability, with concentrations often near the margins of colder shelf water. Depth, prey, age composition and vertical movements can also contribute to the pattern. The maps show spatial associations between survey distributions and temperature. Other environmental and biological factors may also contribute. They use locally interpolated, density-dependent-corrected CPUE within survey coverage. The example's correction version and adoption status are unspecified, so its CPUE surface serves as a descriptive illustration alongside the assessment indices.

The standard EBS survey expanded northwest in 1987. Sampling in the separate NBS region has varied among years, while pollock distribution has shifted between regions. Model-based estimates integrate station observations, including zero catches, across the EBS and NBS and account for spatial and temporal variation (O'Leary et al. 2020). The biomass index uses sdmTMB and abundance at age uses tinyVAST, replacing the earlier VAST implementation. The complete selected abundance-at-age series incorporates historical revisions as well as the recent observations (Table 11). The regional coverage, model specifications, annual covariance and remaining production-source qualifications are described in Appendix H. Estimates for unsampled years are model predictions.

The BTS length distributions show pronounced changes in the sizes represented in the survey (Figure 13). Peaks among small fish and their progression toward larger lengths illustrate cohort growth, while the relative contribution of small and large fish varies among years. These distributions complement the age compositions by showing changes in size directly.

The BTS age compositions show changes in the ages represented in the survey and the progression of cohorts through time (Figure 14). Highlighted bars follow the 1992, 2012 and 2018 year classes through successive ages. The 2012 year class progresses from age 1 in 2013 to age 10 in 2022, while the 2018 year class progresses from age 1 in 2019 to age 8 in 2026. The blank 2020 row marks the gap when the survey was absent.

The age compositions combined traditional age readings and FT-NIRS estimates. We converted the 2026 FT-NIRS estimates to traditional ages using 307 screened paired readings (Appendix G). The conversion preserved total survey abundance and pooled ages 15 and older into the 15+ group.

Spatiotemporal estimation of BTS weight at age, adopted in 2022, replaces the earlier conversion of mean length to weight (Indivero et al. 2023). The full BTS-derived weight schedules used for population biomass are presented in Table 12; the compact biological-input summary retains the recent fishery and spawning schedules and fixed biological parameters. We applied the screened 2026 age key consistently to BTS weight-at-age means and their delta-method standard errors, retaining the joint covariance treatment. We used the same key for the provisional ATS abundance and biomass-derived weights, including the weights used to predict the 2026 AVO index. We refitted the standalone random-effects weight-at-age model with the revised BTS means and standard errors under the existing equations, likelihood and source-covariance treatment, preserving fishery observations and the 2025 conversion key. These uncertainty calculations exclude empirical age-key uncertainty. The conversion, weight calculations and assessment use are described in Appendix G.

The missing 2020 survey and subsequent 2021 observation affect interpretation of the time series. Retrospective comparisons describe changes in population estimates as observations accumulate (Section 6.1.5); they do not isolate the influence of these two survey years.

4.2.2 Acoustic-trawl survey

The 2026 ATS ages are provisional: we apply the BTS age–length key (ALK) to ATS population length-frequency estimates. FT-NIRS ages from the 2026 ATS samples will be available in 2027.

Historical and acoustic-trawl sampling effort is summarized in Table 13. The 2026 collections retain their regional scope, while unavailable direct age counts remain NA.

Acoustic-trawl surveys (ATS) are typically conducted every other year and estimate the off-bottom component of the pollock stock, complementing the annual BTS index of near-bottom pollock.

Estimated pollock biomass for the EBS shelf has averaged about 3.202 million t since 1994, when the revised time series begins (Table 14). The revised estimates include the water column down to 0.5 m above the seafloor; the historical

midwater index extended to 3 m above the seafloor. Low recruitment during the relatively warm early 2000s was subsequently reflected in lower survey biomass during the cold period of 2006–2012 (Honkalehto and McCarthy 2015). With growth of the strong 2012 year class, biomass exceeded 4 million t in 2014 and 2016, another warm period.

The 2026 regional estimates included the Steller sea lion conservation area (SCA) and the Northern Extension (Table 15).

Historical abundance at age

The historical series showed strong pulses of age-1 fish in 1997, 2007, 2009, 2014 and 2024 (Table 16). The provisional 2026 estimates were largest at ages 2 and 3: 3.73 and 3.96 billion fish, respectively, compared with 2.03 billion age-1 fish.

The ATS age compositions show pulses of young fish and the progression of cohorts through successive surveys (Figure 15). The highlighted 2018 year class appears at age 4 in 2022, age 6 in 2024 and age 8 in 2026. Ages 2 and 3 dominate the provisional 2026 composition. The 2026 estimates use the BTS age–length key applied to ATS population length frequencies; FT-NIRS ages from the 2026 ATS samples are expected in 2027. Blank years represent gaps in the age-composition series.

2026 survey design and results

The 2026 ATS covered the EBS shelf from approximately 60 m depth to the shelf break between 30 May and 4 August. Core-area transects were spaced at 20 nmi; selected transects extended into the Northern Extension at 40 nmi spacing. Survey timing was similar to 2022, with a longer duration in 2026 because of special projects. The area east of 170° W was surveyed from 30 May to 25 June, and the area west of 170° W from 26 June to 4 August.

The preliminary 2026 core-area estimate was 12.7 billion fish and 3.682 million t of biomass (Table 17). Biomass increased by 28% relative to the 2024 estimate of 2.875 million t (11.4 billion fish). The 2026 biomass estimate was 4% below the 2022 estimate of 3.834 million t (9.67 billion fish) and 16% above the 1994–2024 survey mean of 3.176 million t. These are survey estimates; assessment intake and adoption remain separate decisions.

Provisional 2026 ATS age compositions over supported lengths were estimated by applying the FT-NIRS-based BTS age–length key to ATS abundance-at-length (Appendix G). Midwater pollock were concentrated between the 100 m contour and the shelf break, particularly northwest of Unimak Pass and between Zhemchug and Pervenets canyons. Length distributions suggested that presumed age-1 fish (<20 cm fork length) formed dense schools on the northwestern shelf, with smaller concentrations east of the Pribilof Islands and near Unimak Pass. Presumed age-3+ fish accounted for most biomass, with 74% of their biomass west of 170° W.

Provisional abundance at age from the 2026 BTS keys

We estimated provisional ages by matching ATS lengths to the 2026 BTS FT-NIRS age–length key, then converting those ages to traditional ages using paired readings. Ages 2 and 3 dominated the resulting distribution (Table 18; Figure 16). Fish beyond the key's length range represented less than 0.001% of core-area abundance. The technical appendix describes the keys, calibration and allocation assumptions.

These ATS age estimates are provisional and will be updated with ATS age-composition data in 2027. Assessment comparison and final model selection remain separate from this derivation.

Provisional biomass and mean weight at age

We applied the same age keys to biomass and divided biomass by abundance to estimate provisional mean weights at age. The biomass allocation and mean weights are presented in Table 19, Table 20 and Figure 17. These estimates share the provisional status of the age compositions.

Survey uncertainty and assessment treatment

We estimated ATS biomass sampling error with the established one-dimensional geostatistical method, which accounted for spatial patterns along survey transects (Petitgas 1993; Walline 2007; Williamson and Traynor 1996). The 2026

sampling error was 3.4% of estimated biomass, compared with a historical average of 4.4%. Complete core-area coverage at 20 nmi transect spacing likely contributed to the lower error.

For assessment use, we increased the survey errors to allow for additional uncertainty, including target strength and trawl selectivity. We scaled the assessment coefficients of variation to average 20% over 1994–2026 and assigned a value of 45% to 2000 (Table 14). We retained the established geostatistical method because the approach described by Urmy et al. (2025) was unavailable for production use in 2026.

The assessment used 20 biomass observations through 2026. Table 21 describes the separate treatment of survey age compositions.

Spatial and historical patterns

The Northern Extension contained an additional 0.078 million t beyond the 3.682 million t core-area estimate (Table 15).

The ATS index used in the assessment covered the water column down to 0.5 m above the seafloor, while the historical heatmap showed biomass from near the surface to 3 m above the bottom (Figure 18). Biomass was low during 2008–2012, especially in 2008–2009, and increased in 2014–2016. The maps also showed shifts in the strength and location of concentrations along the western and southern shelf.

Biomass time series and observation errors

The series contained 20 survey estimates between 1994 and 2026. We plotted approximate 95% sampling intervals around each biomass estimate using the survey standard errors (Figure 19; Table 14). Gaps in the annual series marked years without a survey estimate. These intervals described survey sampling variability; the assessment used larger errors to allow for additional uncertainty.

Biomass varied strongly along the survey tracks. Concentrations were highest on parts of the western and southern shelf and generally lower in the far north (Figure 20). The map included the Northern Extension and showed GMT contours of mean biomass per sampling interval (tons per EDSU). The map uses approximately 10-nautical-mile cell means, with the smoothed surface displayed within 30 km of sampled cell centers.

The western region contained the greatest biomass and abundance, followed by the eastern region and the SCA (Table 15).

4.2.3 Acoustic vessel-of-opportunity index

The acoustic vessel-of-opportunity (AVO) index provides a relative-abundance observation for age-1 and older midwater pollock within the U.S. EEZ (Honkalehto et al. 2011; Lauffenburger et al. 2025). The 2026 observations were collected at 38 kHz from 29 May to 5 August aboard the summer bottom-trawl charter vessels. The index covers the water column from 3 m above the seabed to 16 m below the surface.

Annual index

Relative AVO backscatter is lowest at 0.299 in 2009 and highest at 1.55 in 2015 (Table 22; Figure 21). It is 0.863 in 2024, 1.07 in 2025 and 1.12 in 2026, with sampling CVs of 2.72%, 5.26% and 4.55%, respectively. The 2025 and 2026 sampling confidence intervals overlap. The display series covers 2009–2026, excluding 2020, when no bottom-trawl survey took place. Each annual area-integrated estimate is divided by the mean of the 17 estimates to give a mean-one series. The table also retains total pollock backscatter, in millions of square metres, with sampling errors and scaled confidence intervals. This is an acoustic backscattering cross section, rather than geographic area (Lauffenburger et al. 2025).

For the nine paired survey years in the 2009–2026 display period, AVO total pollock backscatter and revised ATS biomass have a squared Pearson correlation of 0.89. This association is descriptive: AVO covers 3 m above the seabed to 16 m below the surface, while the revised ATS index extends from 0.5 m above the seabed to 16 m below the surface. The vertical domains differ. The 2026 AVO report gives 0.90 using an ATS 2026 estimate of 3.62 million t; the comparison here uses the revised ATS estimate of 3.682 million t (AFSC-RACE MACE 2026).

Spatial distribution and sampling coverage

Strong backscatter occurs along western EBS tracks in the recent maps, with weaker returns interspersed along central and southeastern tracks (Figure 22). The annual panels use a common map extent and color scale while showing differences in track coverage.

The longer series also contains alternating strong and weak returns along the sampled shelf (Figure 22). The year 2020 is omitted because AVO observations are unavailable. Survey coverage and processing vary among years. Inferences about population redistribution require comparable sampling and integration across years.

The number of observed survey intervals ranges from 5,557 in 2009 to 12,229 in 2024 (Table 23). Recorded-zero shares fall from 56.3% in 2009 to 7.8% in 2015, vary subsequently, and are 14.9%, 5.0% and 14.2% in 2024–2026, respectively.

The maps summarize observed pollock backscatter in approximately 20-nautical-mile cells, including recorded zeros. GMT gridding and contours show the mean backscatter in m² nmi^{-2} on a common color scale. Surfaces extend up to 45 km from sampled cell centers; more distant areas remain masked. The annual AVO index integrates backscatter over the survey area and is reported in million m². Survey coverage and processing vary among years, so comparisons of spatial patterns depend on comparable sampling.

Assessment use and observation uncertainty

The AVO assessment index is 2.01 million m² in 2024, 2.50 million m² in 2025 and 2.60 million m² in 2026 (Table 24). Its 20 observations cover 2006–2026, excluding 2020. The assessment uses the newly processed 2011 and 2013 estimates and their sampling errors (AFSC-RACE MACE 2026). The 2006–2008 estimates retain the legacy rescaling described in the 2024 assessment (Ianelli et al. 2024). Mean-one display normalization remains separate from the assessment inputs in million square metres.

Survey sampling errors describe one-dimensional geostatistical variability. The assessment applies additional observation-error inflation to an arithmetic mean CV of 23% across its 20 observations. The two newly processed years use the existing modern-year error inflation rule, followed by common recalibration of all 20 CVs to that mean. Assessment errors differ from sampling errors and from the ATS mean assessment CV of 20%.

For the Normal index likelihood, arithmetic observation SEs equal the retained assessment index multiplied by its selected CV. For the display series, let I_t denote the original index, $\acute{I}$ its 17-year mean and $R_t = I_t / \acute{I}$. The same constant rescales its index, SEs and confidence limits, leaving the CV unchanged:

$$SE(R_t) = R_t CV_t, [L_t, U_t] = \frac{I_t \pm 1.96\, I_t CV_t}{\acute{I}}.$$

The source 95% interval is represented by the half-width $1.96\, I_t CV_t$, checked for every observed year. The displayed intervals describe survey sampling uncertainty with normalization treated as a change of units; assessment error inflation is a separate model input.

4.3 Other data

Model 26.0 fitted the 12 historical CPUE observations from 1965–1976 using fishery selectivity and fishery weights at age. We retained the observations and errors from the reference fit through 2024.

Fixed biological assumptions

We retained the 2024 biological assumptions (Ianelli et al. 2024). Natural mortality was 0.9 yr^{-1} at age 1, 0.45 yr^{-1} at age 2 and 0.3 yr^{-1} at ages 3–15+. Maturity at ages 1–15+ was 0, .008, .289, .641, .842, .901, .947, .963, .970, 1, 1, 1, 1, 1, 1. The model calculated April spawning biomass one quarter-year after the start of the year and applied a female fraction of 0.5 once to sex-combined abundance. We used a 15-by-15 identity matrix for ageing error, matching the archived ADMB configuration. This likelihood assumption treated ages as exact after data preparation, which used FT-NIRS age conversion and age–length keys.

All weight schedules covered ages 1–15+ for the combined sexes in kg per fish. The model used fishery weights to calculate catch biomass, BTS weights to calculate population biomass and A-season weights to calculate spawning biomass.

Fishery weights

We retained the fixed 1964–1990 schedule and used annual WtAgeRe observation means at ages 3–15+ for 1991–2025. Saved model predictions filled the missing 2025 values at ages 13–14. For 2026, we used standalone random-effects model predictions at ages 3–15+ and carried forward the 2024 weights at ages 1–2. These updates changed 442 of 915 cells over 1964–2024, all at ages 3–15+ in 1991–2024. We recorded which weights came from observations, filled missing values or forecasts. The weight-model fit used the underlying BTS survey observations and treated the age-conversion key as fixed when calculating errors. We assigned a 100% CV to the 2024 fishery weight at age 15+ to reflect its uncertainty.

Population and spawning weights

Population weights use BTS observations through 2026. Three historical values retain their previous treatment; details are in the technical supplement.

For spawning biomass, we replaced the older weights at ages 3–15+ with A-season fitted values for 2024–2025 and projections for 2026. Within the common-year comparison, this update changed the 13 values at ages 3–15+ in 2024. We retained baseline weights at ages 1–2; age-2 maturity was .008.

ATS and AVO weights

We retained each historical schedule through 2024 and carried its weights forward to 2025. After BTS-ALK and conditional TMA preparation, we calculated provisional 2026 ATS weights by dividing biomass by the corresponding abundance. For age 15+, we pooled biomass and abundance before division. Appendix G documented the supported lengths and unallocated tail. We used the 2026 ATS vector as a proxy for AVO weights, although the historical schedules differed.

Section 5.2 described the rationale for changing weight estimation. Weight-schedule uncertainty remained separate from the assessment's conditional parameter uncertainty. Native forecasts repeated the 2026 weight vectors. The proposed Tier 3 handoff used its documented fishery forecast and carried-forward weights. The projection inputs and catch recommendations are presented in the projection and harvest-advice sections.

4.3.1 Age-composition input sample sizes

The full-data Model 23.0 fit uses fishery composition N=443, 578 and 530 for 2023, 2024 and 2025, respectively (Table 21). These are the rounded continuity-scaled harmonic estimates from the corrected sampler bootstrap, applied in the completed full-data Model 23.0 fit at the author's direction. The bootstrap preserves linked sampling units within fishery strata and uses 10,000 replicates per year. ATS 2026 uses N=94, and BTS retains the selected 2026 N, shown as 131. Fishery N is fixed at 10 in each year from 1964–1977. The three composition-likelihood multipliers are 1.

Model 26.0 fits age compositions with a multinomial model. The baseline retains likelihood sample sizes from the bridge workbook. These likelihood sample sizes differ from counts of fish sampled and from diagnostic effective sample sizes. For every fishery or survey year, the final assessment must provide the raw sampling information, starting sample size, adjustment method, and final sample size used in the likelihood. The 2020 biomass estimate remains a separate fitted survey observation.

One-step-ahead (OSA) residuals compare an observation with its conditional prediction after preceding age categories have been considered. The OSA and Francis comparisons document the selected weights and observed-data residual patterns in Section 6.1.6.

4.3.2 Annual fishery weight at age

Mean weight at age connects the assessment's estimates of fish numbers to catch biomass and harvest projections. Seasonal and spatial differences in fish condition and fishery distribution affect these means. Fishery sampling is

therefore organized by season and area, with stratum estimates combined using the corresponding catch information. Completed-year observations provide the basis for estimating weights for the observation year and forecasting the weights needed for future catches.

For recent fishery samples, Fourier-transform near-infrared spectroscopy (FT-NIR) of otoliths provides predicted ages. Year-specific paired FT-NIR and traditional morphological ageing (TMA) observations align these predictions with the TMA age basis. Estimated abundance and biomass in each FT-NIR age bin are redistributed using the same conditional tester-age key. Mean weight at a target age is then converted biomass divided by converted abundance. For the 15+ group, biomass and abundance are pooled before taking their ratio. The conversion retains each source bin's mean weight when allocating it among tester ages. Its reported uncertainty conditions on the selected key; the supported ages and approved tail-allocation rules are documented in Appendix G.

The weight-model approach described in Section 14.5 of the 2024 assessment (Ianelli et al. 2024) links growth within cohorts to annual changes in condition. A smooth mean age pattern defines the typical weight increments. A cohort's weight in the previous year provides the starting point for the next age, while annual random effects allow variation in starting weight and growth. Observed mean weights and their variances inform this model, separating sampling precision from variability among years. The fitted model supplies predictions for missing observations and forecasts for future years. Model 26.0 uses the resulting weight schedules as fixed biological inputs.

The annual fishery weights in Table 25 are those used in the completed full-data Model 23.0 fit. All 453 available observation means at ages 3–15+ through 2025 agree with the verified annual WtAgeRe source. Missing 2025 ages 13 and 14 use fitted predictions; 2026 ages 3–15+ use saved current-year predictions. Ages 1–2 in 2025–2026 retain the 2024 inputs. The 1964–1990 schedule remains fixed, and the terminal age is 15+.

The 2027 and 2028 rows contain the completed fit's annual fishery-weight forecasts. Age-3 weight is 0.370 kg in both years, and 15+ weight is 1.50 and 1.48 kg, respectively. The projection uses annual fishery weights for catch and A-season weights for spawning biomass.

The technical supplement documents data sources and preparation, model configurations and run identities, numerical checks, and instructions for reproducing the report.

5 Analytic approach

5.1 Main and comparison models

Model 26.0 (Rceattle) is the adopted model for Tier 3 calculations and specifications. Models 23.0 and 23.2 provide comparisons with the previous assessment and the transition in model structure. Other sensitivities separate the early Foreign fishery (1964–1990) from the later domestic fishery (1991–2026) and vary the penalty on declining selectivity.

Model 26.1.2 is the Early-late fishery split sensitivity. The fits and retrospective comparisons hold the process standard deviations fixed at the values estimated by its marginal model. Model 26.0 uncertainty and retrospective diagnostics are described below.

5.1.1 Model 23.0 and impact of new data

As in past years, we incrementally introduced new data relative to the previous assessment while retaining the Model 23.0 formulation. We began with total catch through 2026, incorporating the revised catch estimates for 1991–2026 (2026: 1,312,504.58 t) and extending the model endpoint and parameter and projection horizons to accommodate the added years. At this first step, fishery age compositions and survey observations retained their historical coverage. We then updated weight at age and A-season spawning weights, followed by fishery catch-at-age data and their information weights.

The survey updates followed in sequence: the bottom-trawl survey (BTS), the acoustic-trawl survey (ATS), and the acoustic vessels of opportunity (AVO) index. BTS retained the 2020 survey gap and annual information weights, including the real-valued 2026 input. ATS incorporated the selected age-1 rule and provisional 2026 age information,

and AVO incorporated its annual observations and assessment-error recalibration. Historical revisions and uncertainty recalibration were included with each survey's update (Section 4.2.2; Section 4.2.3).

Each step includes all preceding updates, so its estimated effect depends on the order in which the data were added. The resulting 2026 estimates are summarized in Table 26, and the cumulative annual point estimates are shown in Figure 23. These comparisons use matched definitions of population-weight biomass, female spawning biomass, age-1 recruitment and maximum realized fishing mortality across ages. Model equations are given in the model-equations appendix.

The saved 2027 spawning-biomass and young-abundance projections from the cumulative input-update analyses are compared in Figure 24. These diagnostic projections retain the earlier 2026 catch estimate of 1,310,000 t; the 2026 estimates above use the revised catch inputs. The display starts with Catch + 2026 endpoint, followed by + Fishery age data and the subsequent survey updates. The intervening weight-at-age and A-season spawning-weight updates remain incorporated in the fishery-age and later fits. Abundance at ages 1–3 is measured at the start of 2027, before removals; spawning biomass includes mortality to spawning time. Projection settings are detailed in the technical supplement. Management projections retain their separate catch, selectivity, weight and recruitment assumptions (Section 7). The selected fishery information weights remain 443, 578 and 530 for 2023–2025. The two recent fishery compositions use their respective TMA conversions (Section 4).

The input sequence and configuration comparison answer different questions. The former retains the Model 23.0 formulation while adding observations and updating schedules. The latter compares formulations on matched observations and biology (Table 27). The historical 2024 advice is retained for reference.

Approximate distributions compare the change in spawning biomass from 2026 to 2027 (Figure 25) and the historical change in total abundance at ages 1–4 from 2025 to 2026 (Figure 26). They show the same selected cumulative sequence, followed by Models 23.2 and 26.0. Their input and projection settings are documented in the technical supplement. Each distribution retains its matching fitted-parameter covariance between years and, for abundance, among ages. The normal approximations are conditional on the model and biological and projection assumptions. They describe local estimation uncertainty and are separate from posterior inference.

Among the additions in the saved 2027 projection comparison, ATS produced the largest increase in female spawning biomass: from 2.685 to 2.975 million t, an increase of about 290,000 t (10.8%) relative to the preceding BTS step. AVO increased projected spawning biomass a further 61,000 t (2.0%), to 3.036 million t. The effects on estimated 2026 spawning biomass differed: fishery age data produced the largest increase, about 213,000 t (8.4%); ATS reduced it by about 73,000 t (2.6%), and AVO increased it by about 47,000 t (1.7%). These are successive changes within the specified sequence, with each survey contrast including its historical revisions and error calibration.

5.1.2 Model 23.2 fitted to the 2026 assessment data

Model 23.2 is a structural bridge between the ADMB continuity model and the Rceattle configuration. Its annual spawning-biomass and recruitment trajectories retain their documented scope. The established zero-variance management quantities are excluded from management comparisons. Current configuration effects use verified saved population outputs and their stated units. The comparison of Models 23.0 and 23.2 includes matching local asymptotic uncertainty ranges (Figure 27). Biomass uses common population weights, and fishing mortality is the maximum realized rate across ages. Positive annual quantities use the stated first-order log-scale intervals.

5.1.3 Model 26.0 fitted to the 2026 assessment data

Model 26.0 uses one fishery and fixed weight-at-age schedules. The updated base model fit retains its native working-likelihood, parameter bounds, selectivity forms, priors and fixed biological assumptions. We check outputs against the written inputs and saved native quantities. Population-weight biomass, female spawning biomass, age-1 recruitment and maximum fishing mortality retain distinct definitions. Their annual estimates and matching local asymptotic uncertainty are compared with Model 23.2 in Figure 28. These ranges use each model's own covariance; each interval describes uncertainty around its fitted estimate.

5.1.4 Model 26.1: Foreign and Modern domestic fisheries

The split-fishery alternatives distinguish Foreign fishing in 1964–1990 from Modern domestic fishing in 1991–2026. Model 26.1.0 supplies the fitted split-fishery comparison; Model 26.1.1 reduces the declining-selectivity penalty, and the zero-penalty case removes it. These models retain their recorded fixed annual process-SD settings. Model 26.1.2 extends that formulation by estimating the Modern annual selectivity-process SD while integrating over annual selectivity effects. The Foreign process SD remains fixed. Local uncertainty uses the verified marginal information for this alternative; its observation diagnostics condition on the fitted annual effects.

5.2 Weight-at-age treatment in the model transition

Model 23.0 estimates weight at age using cohort and year effects and their input sigmas. The Model 26.0 configuration instead uses fishery, population and spawning-weight schedules. The fishery and BTS weights retain their year-specific age conversions and observed-data masks. The 2024 fishery weights use the same annual tester-age key and pooled-age rule as the 2024 composition. The 2025 fishery uses its separate key and age-13 midpoint. Future schedules keep their forecast-year identity.

The comparison distinguishes differences in schedule means from the treatment of weight uncertainty. Weight-estimation uncertainty is outside the base Model 26.0 fit and the recruitment-only projection simulations. Population weights define total biomass; female spawning weights, maturity, female fraction and spawning timing define spawning biomass.

5.3 Estimation and numerical checks

Fitted-model results require successful native execution, the recorded gradient criterion, a positive-definite Hessian, finite parameter standard errors and independently verified input identities. These checks establish numerical usability. Scientific interpretation, model selection and final advice remain separate decisions.

We convert recent fishery age compositions before fitting; the observation ageing-error operator remains the documented identity. The fits retain the AFSC composition working score and its selected information weights. Local asymptotic standard errors use the fitted-model native covariance.

6 Results

6.1 Model 26.0 assessment

Model 26.0 provides the main estimates of stock size and the basis for the Tier 3 catch limits presented here. Results from the other model configurations, including Model 26.1.2, show how alternative assumptions affect the assessment. Additional analyses describe uncertainty in Model 26.0 estimates and how those estimates change when recent years of data are removed. Comparisons with past assessments use the data and catch advice reported in each assessment year.

6.1.1 Population estimates and uncertainty

The Model 26.0 base model estimates 2026 total biomass at **12.162 million t**, female spawning biomass at **3.057 million t**, age-1 recruitment at **36.3 billion fish**, and maximum fishing mortality across ages at **0.299 per year** (Table 28).

The local asymptotic 95% ranges in the tables use the revised fit's native covariance. The uncertainty figure is shown in Figure 29. They describe estimation uncertainty under the base model. Weight, maturity, female fraction and natural mortality remain fixed inputs. Their uncertainty and alternative model assumptions remain separate questions. Annual estimates and ranges appear in Table 29.

Regularized-posterior intervals use all 5,000 existing draws, using the final reported summaries (Table 30). The 2026 female spawning-biomass median is **2.946 million t**, with a central 95% credible interval of **2.450–3.584 million t**. Total-biomass median is **11.462 million t**, with interval **9.215–14.719 million t**. Age-1 recruitment median is **31.1518 billion fish**, with interval **12.2880–91.0616 billion fish**. These are the final reported MCMC summaries; production point estimates and asymptotic intervals retain their separate roles.

Numbers-at-age track cohorts through the population (Figure 30). This shows periods of how recruitment and changes in fishing mortality by age affect abundance relative to their mean abundance estimates (and uncertainty) propagates through time. Actual point estimates are reported in Table 31, while the 2026 abundance-at-age estimates (and asymptotic uncertainties) are shown in Table 32.

Comparison with the 2024 assessment

The 2018 year class remains a large recruitment point estimate (Figure 31). Its age-1 abundance in 2019 is **87.8 billion fish** in the Model 26.0 base model, compared with **81.1 billion** in the accepted 2024 assessment, a revision of **8.3%**. The revision reflects both updated observations and changes in model structure.

The estimate for the same 2024 female spawning biomass changes from **3.389 million t** in the saved 2024 assessment to **3.526 million t** in Model 26.0, a revision of **4.0%**. Within the trajectory, spawning biomass changes from that 2024 estimate to **3.057 million t** in 2026 (**-13.3%**). Total biomass changes from **10.767 million t** to **12.162 million t** over the same years (**13.0%**). These within-trajectory changes differ from shared-year assessment revisions.

The recruitment comparison uses calendar age-1 year and billion-fish units (Figure 31). Model 26.1.2 results are included to examine how recruitment estimates for the early period change relative to the 2024 assessment. Historical values are point estimates. Current error bars use each model's local asymptotic covariance: the base-model covariance for Model 26.0 and the marginal information for Model 26.1.2. Recent years without a historical estimate are identified. These intervals describe uncertainty around the fitted estimates.

Recruitment per unit of spawning biomass

The log recruits-per-spawner diagnostic from the 2024 assessment is updated using the Model 26.0 estimates (Figure 32). For each birth year, age-1 recruitment in the following year is divided by female spawning biomass in the birth year. Taking the natural logarithm puts proportional changes on a comparable scale: higher values indicate more recruits per unit of spawning biomass.

The series shows substantial variation among year classes. The 1978, 2008 and 2018 year classes have particularly high recruits-per-spawner values. Following the strong 2018 year class, the ratio declined through the 2022 year class; estimates for 2023–2025 are higher but remain less certain. The spawning-biomass panel generally shows lower ratios at higher biomass, with considerable variation among years at similar biomass. This pattern can reflect density dependence, environmental conditions and estimation uncertainty. Spawning biomass appears in both the horizontal axis and the ratio, which can contribute to the apparent relationship. Model 26.0 uses mean recruitment rather than an estimated Ricker curve. The figure provides a descriptive check on productivity using point estimates of the recruitment-to-biomass ratio.

Biological schedules and selectivity retain their age, sex and year definitions (Figure 33; Table 33; Table 34). Fishing mortality is also shown by age (Figure 34). Whole-year spawning exploitation compares paired spawning-biomass calculations from the base model production fit (Figure 35). For the interval from spawning in 2026 to spawning in 2027, its production-point estimate is **13.6793%**: **3.3260 million t** with fishing versus **3.8531 million t** without fishing. The branches share prior fishing history and their spawning-date starting population, with 1.375 million t annual catch assumed in 2027. The separate MCMC posterior median for this exploitation fraction is **14.5333%**, with a central 95% credible interval of **11.2075–18.3700%**. These paired-state calculations carry the revised fishery-weight schedule into 2027 and retain the fixed 1.375 million t catch assumption. The production-fit point remains separate from that posterior interval.

6.1.2 Estimated model parameters

The parameter-family summary identifies fitted parameter groups and their asymptotic uncertainty (Table 35). Complete estimates and the largest parameter correlations are retained in the detailed estimates appendix. The summary likelihood contributions are in Table 36, with the full component breakdown in Table 37. Their values follow the native working-likelihood and penalty definitions.

6.1.3 Fits to the observations

The fitted indices follow the main rises and falls in the survey series, including the low ATS biomass in 2006–2009 and its subsequent increase (Figure 36). Annual departures remain. The BTS residuals include negative departures in 2006

and 2009 and positive departures in 2014–2015, each outside the plotted 95% reference range. The AVO observation in 2015 is also higher than fitted, whereas all ATS biomass residuals fall within that range (Figure 37; Table 38). Thus, the broad survey trends are represented more closely than some individual annual observations. BTS residuals account for covariance among survey years. The retained 2020 ATS age-1 placeholder is excluded from observed-index summaries; the 2020 USV biomass estimate remains an observation.

The fishery selectivity curves show substantial changes in the ages most vulnerable to fishing, with dome-shaped patterns in many years and flatter patterns in others (Figure 38). BTS selectivity generally increases with age and approaches a plateau, while the ATS curve, shared with AVO, is relatively flat at older ages (Figure 39; Figure 40). These curves describe relative age patterns rather than absolute abundance. They include model estimates for unsampled years, including 2026 for the fishery. Recruitment varies strongly among years: the 2018 year class, entering at age 1 in 2019, has the largest positive departure from the fitted recruitment level. The latest age-1 estimates have wider uncertainty ranges (Figure 41).

The mean-age comparisons show that the model follows most of the observed changes in survey age composition and closely tracks fishery mean age in the domestic period. The clearest sustained fishery discrepancy occurs in the early years, when predicted mean ages are higher than observed (Figure 42). The observed bars show plus or minus two standard errors of mean age; they describe sampling uncertainty, rather than uncertainty in the fitted population. Fishery age observations end in 2025. The annual composition panels show how these mean ages arise from the proportions at each age and allow individual cohorts to be followed as they age (Figure 43; Figure 44; Figure 45; Figure 46).

Aggregate observed and fitted age proportions agree closely, especially for the fishery, but the residual panels retain patterns by age and year (Figure 47; Stewart and Monnahan (2025); Monnahan et al. (2026)). For example, BTS residuals include extended runs of the same sign at several ages, and the early fishery residuals also show structured departures. Averaging across years can obscure these differences. The OSA residual standard deviations are below one for all three sources (about 0.76 for the fishery, 0.86 for BTS and 0.64 for ATS), and the QQ plots are generally narrower than the standard-normal reference, although some BTS tail residuals are large. Close aggregate fits therefore need to be considered alongside the annual patterns. The afscOSA v0.0.1 displays retain the native Model 26.0 OSAs and selected likelihood sample sizes. They describe conditional observation fit under the working-score approximation; parameter uncertainty is outside their scope.

6.1.4 Profile likelihood diagnostics

A profile examines how strongly the observations and model assumptions constrain ATS survey catchability in Model 26.0. Catchability scales the survey index relative to the population available to the survey. We fixed this parameter at **17 values from 0.45 to 0.65**, including the freely estimated value of **0.51694**, and re-estimated all other active parameters. The grid uses eight equal intervals on each side of the estimated value: approximately 0.00837 below it and 0.01663 above it. Data, bounds, likelihood terms, and penalties match the Model 26.0 fit, including the updated 2011 and 2013 AVO observations (Figure 48).

The best fit remains at catchability **0.51694**. Lower or higher values increase the negative log likelihood (NLL), indicating a poorer overall fit after the other parameters have adjusted. The conventional change of 1.92 NLL units is crossed between **0.45000 and 0.45837** on the lower side and between **0.58347 and 0.60010** on the upper side. Linear interpolation between these points gives approximate crossings of **0.4576 and 0.5838**. These are diagnostic bounds from the sampled profile; the objective includes model penalties, and the crossings have been interpolated rather than directly fitted.

The component breakdown shows how the model balances the different data sources. At catchability 0.45, the total NLL increases by **2.48** units. Index likelihoods contribute 2.36 units, while the combined composition fit improves by 0.43 units. The ATS index and compositions fit less well, while BTS compositions fit better. At catchability 0.65, the total increase is **6.80** units. Compositions contribute 4.50 units, largely from the BTS, and selectivity penalties contribute 1.93 units. The combined index term changes by only 0.14 units because improved BTS index fit partly offsets poorer ATS index fit. Thus, the preferred catchability reflects agreement across surveys, fishery observations, and selectivity assumptions.

All 17 fits have maximum absolute gradients below 10^{-4} and positive-definite Hessians. Seven points required additional numerical refinement or successful optimizer confirmation. Weighted NLL components sum to the total objective at every point. Scientific evaluation should also consider sensitivity to composition weighting and selectivity assumptions.

The final ATS-q posterior median is **0.5093**, with central 95% credible interval **0.4514–0.5755**. The profile minimum is **0.51694**, with approximate profile limits **0.4576–0.5838**. Figure 49 compares the negative log ATS-q marginal density with the optimized profile NLL. The kernel density integrates to one, and each displayed curve is centered at its own minimum over ATS q. The marginal integrates over the other parameters and includes posterior regularization; the profile optimizes those parameters under the native penalized objective. Panel B fits weighted quadratic curves directly to individual-draw native NLL components and compares them with the profile components. All 5,000 draws contribute; binning is excluded. MCMC curves are centered at the ATS-q marginal mode and displayed over the central 99% of sampled q. These conditional mean curves have a different interpretation from the negative log marginal density, which has no unique additive decomposition into the native likelihood terms. The smoothers summarize the broad component trends; tail coverage is limited by the available draws.

The paired MCMC draws show the relationships among ATS q, summed abundance at ages 2–5 in 2026, age-1 abundance in 2019 (the 2018 year class), and projected 2027 female spawning biomass (Figure 50). Diagonal panels show their marginal densities. All four quantities are paired within each saved draw.

Projected 2027 female spawning biomass is most strongly associated with the summed abundance of ages 2–5 in 2026 among the quantities shown ($r = 0.84$). This contrasts sharply with the 2024 assessment, in which projected 2025 spawning biomass depended heavily on the estimated magnitude of the 2018 year class (Figure 52 of the 2024 SAFE). The 2018 year-class dependence remains in the projection, as shown by its positive correlation with 2027 spawning biomass ($r = 0.36$), but is less pronounced than the dependence on the younger fish now approaching or contributing to spawning biomass.

At each fixed ATS catchability, we also project **2027 female spawning biomass** and its conditional 95% confidence interval (Table 39). The projection uses 1,375 kt catch in 2027, mean 2023–2025 fishery selectivity, April 1 spawning, the weight schedules, and the age-15+ group. At the fitted catchability, the 2027 SSB estimate is **3.325 million t**, reproducing the projection.

These intervals propagate fitted-parameter uncertainty using the inverse fixed-q penalized Hessian and the delta method on log spawning biomass. They condition on each ATS q and the stated projection assumptions. They exclude uncertainty in ATS q, future recruitment process variation, and weight-at-age schedules. Thus, they describe uncertainty in the estimated 2027 SSB at each profile point; the recruitment-only projection bands in Section 7 describe a different source of uncertainty.

6.1.5 Retrospective patterns

Across the full nine peels ending in 2017–2025, Model 26.0 female spawning-biomass Mohn's rho is **+0.121** (Table 40). Age-1 recruitment rho is **−0.0985**, and maximum fishing-mortality rho is **−0.0636**. Each statistic averages the relative revision between a peel's terminal estimate and the full-2026 estimate for that same calendar year. The annual revisions are reported in Table 41. Peel-specific ABC and OFL remain outside this calculation.

The paired retrospective comparison shows female spawning biomass, age-1 recruitment and maximum fishing mortality for Model 26.0 and the Early-late fishery split Model 26.1.2 sensitivity (Figure 51). Each colored curve removes one through nine terminal years and is compared with its own analysis's full 2026 reference. Model 26.0's full nine-peel statistics are reported above.

The conditional Model 26.1.2 comparison has nine-peel spawning-biomass rho of **+0.0735**, recruitment rho of **−0.133**, and maximum fishing-mortality rho of **+0.00226**. Both configurations show positive average spawning-biomass revisions and negative average recruitment revisions, while their average fishing-mortality revisions differ. These full-window summaries describe the overall direction of retrospective change across all nine endpoints.

The conditional alternative fixes Foreign and Modern process SDs at 0.1235 and 0.582 and re-estimates the remaining parameters and annual selectivity effects. Its full reference uses the same fixed SDs. These curves therefore describe a conditional diagnostic rather than a full marginal-estimator retrospective. All panels show point estimates.

The Model 26.0 simulated-data and posterior-predictive retrospective experiments provide separate conditional comparisons (Section 6.3.1).

Historical retrospectives

Successive assessments provide different estimates of biomass as observations accumulate and methods change. Following Figure 62 of the 2024 SAFE, Figure 52 compares beginning-of-year age-3+ biomass from earlier assessments with the Model 26.0 series. The 2024 comparison uses the published values in the official SAFE Table 23. There was no assessment fit in 2025.

For this historical comparison, age-3+ biomass is the sum of beginning-of-year numbers at ages 3–15+ multiplied by the corresponding fishery weights at age in each year. This definition differs from total population-weight biomass reported elsewhere in the assessment.

The 2026 assessment estimates age-3+ biomass in 2023 at 11.983 million t, compared with 9.695 million t in the 2024 assessment, an increase of about 24%. The corresponding estimates for 2024 are 10.185 and 9.415 million t, a difference of about 8.2%. The Model 26.0 trajectory declines from 14.438 million t in 2022 to 8.774 million t in 2025, then rises to 13.286 million t in 2026. Earlier assessment trajectories show that both the magnitude and timing of biomass changes have been revised over time.

These historical comparisons combine the effects of new observations, revised biological inputs and changes in model structure. The retrospective peels above instead remove recent years within a specified model framework. The historical figure is therefore a record of assessment revisions, with each series retaining its original assumptions and available years.

6.1.6 Age-composition weighting comparison

The age-composition diagnostics suggest relatively minor changes to the weights used in the 2026 assessment (Figure 53; Figure 54). Given the small suggested adjustments, the 2023–2025 fishery age compositions therefore retain input information weights of 443, 578 and 530. These weights determine the influence of the compositions in model fitting. The data section reports the raw numbers of fish sampled.

Composition weighting can be evaluated more fully in future assessments, including alternatives such as the Dirichlet-multinomial distribution, which allows greater variation among age samples than the multinomial model. Such comparisons would help assess whether estimating this additional variation within the model improves the treatment of age-composition data and changes the resulting stock estimates.

6.2 Model 26.1 split-fishery alternatives

Models 26.1.0 and 26.1.1 separate the Foreign fishery through 1990 from the Modern domestic fishery beginning in 1991. Their annual selectivity standard deviations are fixed at 0.5, while the declining-age penalty is 12.5 in Model 26.1.0 and 3.125 in Model 26.1.1. The zero-penalty case provides a further comparison. The Model 26.1.2 sensitivity shown here retains the 3.125 penalty and fixes the Foreign and Modern annual process standard deviations at 0.1235 and 0.582, respectively. The remaining parameters and annual selectivity effects are estimated conditionally on these values.

The likelihood comparison separates data-likelihood contributions, penalty terms and the marginal fitting criterion (Table 42).

The projections use the revised ordinary fits and the Early-late fishery split Model 26.1.2 comparison. The alternatives retain matched observations, likelihood weights and projection assumptions (Table 43). Model 26.1.2 is the sensitivity analysis; Model 26.0 (Rceattle) is the adopted assessment model and main proposal for Tier 3 calculations and specifications. Both comparisons use the same projection-selectivity window. Adopted catch specifications remain separate.

6.2.1 Selectivity

Separating the Foreign (1964–1990) and Modern domestic (1991–2026) fisheries changes the fitted selectivity most visibly near the boundary between the two periods. Over much of the later period, Models 26.0 and 26.1.0 retain similar annual age patterns, including the peaks and declines at older ages (Figure 55). The split allows the Foreign curve to differ more freely from the Modern curve, while Model 26.0 represents the history as a single fishery.

Both models estimate a shift from 1990 to 1991 toward lower relative selectivity at younger ages and higher relative selectivity at ages 11 and older. The shift is larger in Model 26.1.0: at age 5, the change is about −0.69 on the native mean-one scale, compared with −0.31 in Model 26.0; at ages 12 and older, the increases are about 0.55 and 0.24, respectively (Figure 56). This comparison shows how the split changes the estimated transition. The fitted curves alone establish neither an abrupt change in fishing practices nor an improvement in the population estimates; the observation fits and retrospective results provide the complementary checks below.

The full-history curves are divided by each year's maximum so that their shapes can be compared. The boundary-year display instead retains native mean-one normalization, and its differences refer to that scale. These are fitted point curves. The Model 26.1.2 curves used in subsequent comparisons retain the fitted selectivity states from its fixed-SD conditional fit and remain distinct from the Model 26.1.0 comparison here.

6.2.2 Fishery age-composition fits

We compared Models 26.0 and 26.1.2 using the same 62 annual fishery age compositions and likelihood weights (Figure 57; Figure 58). Model 26.0 fitted one fishery across the full record; Model 26.1.2 separated the Foreign fishery through 1990 from the Modern domestic fishery beginning in 1991. We displayed both models over those same periods to compare their fits directly. Both models reproduced the dominant age groups in the combined compositions, while the annual plots revealed differences that the combined summaries obscured.

Model 26.1.2 fitted the Modern domestic observations more closely. It reduced the sample-weighted root-mean-square difference in age proportions from **0.693 to 0.590 percentage points** and the root-mean-square error in annual mean age from **0.123 to 0.0963 years**. Both models predicted a younger mean age than the observations showed in 1991, but Model 26.1.2 reduced that discrepancy from **0.510 to 0.339 years**. The smaller differences across ages and years showed that the split-fishery configuration improved the modern fit rather than simply matching its combined age distribution.

Model 26.0 fitted the Foreign-period observations more closely. Model 26.1.2 increased the sample-weighted age-proportion error from **2.67 to 3.43 percentage points** and the annual mean-age error from **0.655 to 0.696 years**. Both models predicted older mean ages than the early observations showed, particularly in 1964–1965 and 1970. The age-by-year panels showed persistent overprediction at several older ages and underprediction at younger ages during much of the early record. Separating the fisheries therefore improved the modern fit while leaving larger discrepancies in the Foreign period.

We calculated these comparisons from each model's fitted age proportions and held the observations and their weights constant. Model 26.1.2 used its fixed-process-SD conditional fit. The mean-age bars represented observation sampling uncertainty; the fitted curves represented point estimates. We used these results to assess where each model reproduced the fishery observations and where systematic discrepancies remained. The technical supplement retained the estimation and residual-method details.

6.3 Further model evaluations

6.3.1 Observation-error retrospective self-test

In response to Plan Team and SSC requests to investigate retrospective patterns, we developed a new simulation-based retrospective experiment for this assessment (Figure 59). Its distinctive feature is a paired design that examines revisions both at the fitted parameter estimates and across plausible parameter states from the regularized posterior. One set of ten simulated datasets uses the fitted parameters; a second set uses ten posterior parameter draws. Each dataset has its own full 2026 fit and nine optimization peels ending in 2017–2025. Revisions compare each peel's terminal estimate with its own full fit for the same calendar year. This approach extends the usual retrospective analysis by examining how

observation variation and uncertainty in the generating parameters influence the revisions, providing additional context for interpreting the assessment's retrospective pattern.

Across all nine peels ending in 2017–2025, Model 26.0 assessment Mohn's rho for female spawning biomass is **+0.121**, compared with median values of **−0.0117** for datasets generated at the fitted parameters and **−0.0574** for datasets generated from posterior draws. The assessment value exceeds every simulated rho in both sets: the fixed-parameter range is **−0.115 to +0.0335**, and the posterior-draw range is **−0.0971 to +0.116**. Thus, the positive average spawning-biomass revision in the assessment is stronger than that reproduced by these simulated datasets.

For age-1 recruitment, the assessment's nine-peel rho is **−0.0985**, within both simulated ranges. Median recruitment rho is **−0.00751** in the fixed-parameter set and **+0.0830** in the posterior-draw set. The contrast between spawning biomass and recruitment shows that the retrospective pattern depends on the population quantity considered.

The paired design explores how observation variation and uncertainty in generating parameters contribute to the full nine-peel pattern. Ten datasets per branch provide an exploratory comparison. The technical supplement describes the fixed process history, data-generation approximations and other conditioning assumptions. These experiments assess biomass and recruitment revisions. Confidence bands describing retrospective ABC or OFL patterns are beyond the scope of this work but would be interesting to evaluate.

6.3.2 Posterior diagnostics and predictive checks

The regularized posterior explores parameter uncertainty separately from the production base-model estimates (Figure 60; Figure 61; Figure 62). Four chains each retained 1,250 draws after 1,000 warmup iterations. Combined parameter diagnostics give maximum R-hat 1.024 and minimum bulk ESS 242, with zero divergences and zero maximum-treedepth events. The reported population quantities mix more consistently: maximum R-hat is 1.0013 and minimum bulk ESS is 3,914. MCMC standard errors of the 2026 spawning-biomass and biomass means are about 4,110 t and 20,600 t; their first-half versus second-half mean shifts are less than 0.03 posterior standard deviations. These diagnostics describe the saved chains and retain the recorded parameter-level qualifications.

Final posterior summaries

The posterior mean female spawning biomass in 2026 is 2.965 million t, with a central 95% credible interval of 2.450–3.584 million t (Table 44). These summaries use all 5,000 retained MCMC draws under the regularizing priors and fixed biological inputs. Full-precision results are downloadable as CSV files containing terminal stock and selectivity summaries (CSV), annual population intervals (CSV), summaries of all 1,266 parameters (CSV), and spawning-exploitation intervals (CSV).

Predictive checks ask whether the fitted model can generate data that resemble the observations (Figure 63; Figure 64). We selected 1,000 MCMC draws spanning the four chains. Each draw supplies a possible set of model parameters, from which a new set of observations is simulated. The shaded ranges therefore include both parameter uncertainty and the sampling variation assumed by the model. Points outside these ranges, or repeated patterns of differences, help identify features the model struggles to reproduce. The ranges apply to each age or year individually; some points outside them are expected even when the assumptions are appropriate.

For the indices, simulated observations use the same error assumptions used in fitting, including the relationships among BTS errors. The overall BTS mismatch measure is as large as the observed measure in only 0.2% of the simulations, indicating that this feature of the BTS data is rarely reproduced. Normal error assumptions also allow some simulated indices to be negative.

For age compositions, simulated fish are randomly assigned to age groups using each draw's predicted proportions and the sample size used in fitting, rounded to a whole number. The simulations approximate the fitting method, which adds a small constant (0.001) to age proportions. The figure compares the latest three actual samples for each source with these simulated age proportions. Age conversions, weights and other biological inputs are held fixed. These checks assess how well the model reproduces observations; the retrospective analyses address how estimates change as additional years of data become available.

7 Projections and harvest recommendations

7.1 Proposed Tier 3 configuration and reference points

The proposed fishery selectivity is the mean of the 2023–2025 annual curves, the latest three years with fishery age observations. Each age is averaged before scaling the resulting curve to a maximum of one. The separately inferred 2026 curve is distinguished from directly age-informed years (the projection-input appendix).

Fishery weights use the Model 23.0 joint weight model's 2027 forecast for ages 3–15+, with 2026 weights retained at ages 1–2. 2026 population and spawning weights remain fixed. These schedules are carried through the projection; their estimation uncertainty is outside it.

The deterministic recruitment level uses the arithmetic mean of age-1 entries in 1978–2023, corresponding to birth cohorts 1977–2022. The less precise 2024–2026 entries are excluded. Every entry in the advice tables equals this selected historical mean. The separate projection figures add future recruitment variation to illustrate a range of possible paths. Reference points and fishing rates are reported in Table 45. The Model 26.1.2 column uses the revised Early-late fishery split comparison under the same projection assumptions.

The Tier 3 reference points are spawning-potential quantities. The ABC rule reduces fishing mortality below B40% and sets it to zero below B20%. Projections start from each fitted population and use the stated catch scenarios and biological inputs. The author-rate fraction is one, subject to the risk evaluation and final review.

The previous-year OFL-equivalent fishing mortality is reported in Section 10.

7.2 Standard harvest scenarios

The proposed projections retain the standard harvest scenarios and their common biology (Table 46). The expected-catch scenario assumes 1.375 million t in each of 2027 and 2028; the 2028 catch is a planning assumption. Short-term outputs are reported in Table 47.

The catch and female spawning-biomass panels compare Alternative 2 with recent-mean-F Alternative 3 (Figure 65; Figure 66). Black lines show the means of 1,000 future-recruitment simulations, ribbons show their 5th–95th percentiles, and ten colored trajectories illustrate individual outcomes. Each color follows the same recruitment realization across both alternatives and figures. Recruitment follows the established inverse-Gaussian method, using the arithmetic and harmonic means of the fitted 1978–2023 age-1 entries. Initial age-1 abundance in 2027 equals the historical arithmetic mean; each subsequent entry varies among simulations. The widening bands show how future recruitment variation accumulates as new cohorts enter the fishery and spawning population. Alternative 2 catch paths coincide in 2027 and 2028 because those catches are prescribed.

The projection tables retain all seven scenarios with recruitment fixed at the historical arithmetic mean (Table 48; Table 49; Table 50). The historical phase plot places the fitted 1981–2026 trajectory and deterministic Alternative 2 projections for 2027–2028 on common axes (Figure 67). The horizontal axis is female spawning biomass divided by B35%; the vertical axis is maximum fishing mortality at age divided by F35%. Both denominators are the current primary projection reference points (B35% = 2.182 million t; F35% = 0.688 per year), held constant across the trajectory. This common benchmark shows changes in the fitted stock and fishing mortality through time. The 2024 assessment used annually varying reference points; the historical positions here therefore also reflect the choice of a common current benchmark. F35% is a spawning-per-recruit proxy for FMSY, and B35% is the corresponding spawning-biomass reference point. Solid lines mark ratios of one, and the dashed vertical line marks half B35%. Two-digit year labels trace the trajectory, with the two projection years boxed and shaded yellow. The fitted 2026 position and both projected positions are above B35% and below F35%. The plotted points describe the fitted and deterministic projected states; uncertainty is reported in the accompanying projection figures and decision tables.

Both calculations include the age-15+ group once and retain three spawning-biomass iterations in the Tier 3 harvest rule. The figure ribbons describe future recruitment variation with a fixed fitted starting state and biological and selectivity schedules; assessment parameter uncertainty is outside their scope. Management history remains separate in Table 1 and Table 2.

7.2.1 Decision table

As in past assessments, we present estimated consequences and probabilities for a range of future catches and the subsequent fishing effort implied by each choice (Table 51; Table 52). The decision table compares eight possible 2027 catches, from 10 to 2,540 kt, using Model 26.0 with the updated AVO data. The indicator definitions identify the quantities, comparison benchmarks and years used in the calculations. Together, the tables show how catch choices affect fishing mortality, future spawning biomass and age structure, and how likely each choice is to meet the specified benchmarks.

Each catch column represents a different fishing strategy. Fishing mortality is calculated to achieve that catch in 2027, then held constant through 2031. Catches in subsequent years therefore change with the projected abundance and age composition of the stock. The effort indicator gives the probability that mean age-specific fishing mortality in 2027 exceeds its 2026 value. Interpreting it as relative effort assumes that catchability and the age pattern of fishery selectivity remain fixed. It provides a comparison among strategies rather than a forecast of vessel numbers, fishing days or operating costs.

Readers can compare columns to examine the trade-off between the initial catch and subsequent stock conditions. The probability rows summarize the chance of crossing the stated biomass, fishing-mortality or age-structure benchmarks. These are conditional comparisons that support catch decisions alongside the standard harvest projections and risk table.

Entries are approximate chances out of 100. They propagate uncertainty in the fitted population, selectivity, estimated mean recruitment and comparison benchmarks. Future recruitment is held at the estimated 1978–2023 mean, and biological schedules are fixed. These probabilities describe uncertainty in the assessment estimates under each catch choice. Additional year-to-year recruitment variation and uncertainty in weights and maturity are outside these calculations. Values displayed as 0 or 100 reflect rounding.

The 1,270-kt column is retained for comparison with the earlier decision-table analysis. The expected-catch scenario elsewhere in this section assumes 1,375 kt in 2027 and 2028. The decision tables provide a broader comparison of catch choices; they use local asymptotic uncertainty rather than MCMC draws.

7.2.2 Reference point sensitivity to fishery selectivity specification

Fishery selectivity determines how fishing mortality is distributed across ages and therefore affects both the reference fishing rates and the catches obtained at those rates. The central projection uses the age-specific arithmetic mean of the raw 2023–2025 fishery selectivity curves, rescaled so that its maximum is one. This recent three-year average smooths annual variation while retaining the age pattern from the latest years with fishery age observations. Averaging precedes rescaling because normalizing each annual curve first would give a different average. Figure 68 introduces this recent-average specification and contrasts it with individual annual estimates. The inferred 2026 curve is shown separately because fishery age observations end in 2025. It provides a terminal-year comparison, while the management projections retain the 2023–2025 average.

The recent curves show that changes in fishery recruitment involve particular ages rather than a uniform shift toward younger fish. The first ascending half-maximum crossing was near age 6 in 2021–2022, compared with approximately 4.5, 5.0 and 5.2 in 2023–2025. Relative selectivity at age 5 was especially high in 2023 and then declined in 2024–2025. Selectivity at ages 1–3 remained low and varied independently. These differences matter because a curve that exposes age-5 fish more strongly can produce a different catch and spawning impact from one that concentrates fishing at older ages, even when both have the same maximum selectivity.

The full annual comparisons show which parts of each curve differ from the projection assumption (Figure 69). The repeated 2023–2025 reference makes shifts in the ascending limb, peak and older-age tail visible across 2007–2026. For example, the 2019 curve peaks near age 7 and declines substantially at older ages, whereas several earlier curves retain high selectivity across the older ages. Thus, the annual panels provide the age-specific explanation for the management differences in the next figure. The older curves represent alternative fishing patterns for sensitivity analysis. Their relevance as forecasts depends on evidence about current and future fishery conditions.

The selected recent-average curve gives relatively conservative catch ceilings compared with most annual alternatives (Figure 70). Fourteen of the twenty annual curves produce a higher maximum ABC in 2027, and fifteen do so in 2028.

Relative to the matching recent-average baseline, annual-curve changes in maximum ABC range from −10.1% to +21.4% in 2027 and from −7.5% to +10.0% in 2028. This is a comparison of catch ceilings under the specified projection conditions. It supports the recent average as a relatively cautious central assumption, while also identifying plausible lower outcomes. The lowest annual-curve result remains a conditional sensitivity case, and the curves carry equal status as comparisons rather than assigned probabilities.

The ranking changes between projection years. The 2018 curve gives the lowest maximum ABC in 2027, while the 2019 curve gives the lowest in 2028 despite increasing the 2027 maximum ABC. The latest observation-informed curve, 2025, raises the 2027 ceiling by about 0.9% but lowers the 2028 ceiling by about 4.5%. These reversals arise because the full age pattern affects catches immediately and changes the ages surviving into the following year. The comparisons hold the fitted starting state, recruitment draws, biological schedules and assumed catch of 1.375 million t in each year constant. F40% and F35% are recalculated for each selectivity curve to preserve their spawning-per-recruit targets. The spread among curves is a sensitivity range conditional on these assumptions.

The younger-age summaries explain why a single description such as “younger selection” is insufficient (Figure 71). They separate exposure of ages 1–3, selectivity at ages 4 and 5, the first ascending half-maximum age, and changes across the full curve. For example, from 2022 to 2023 the half-maximum age falls from about 6.0 to 4.5 years, while the curve-weighted mean age changes only from 10.09 to 10.00 years. Conversely, from 2023 to 2025 the half-maximum age rises while the curve-weighted mean age falls. The ascending limb and older-age tail therefore move in different directions. This figure connects the detailed annual panels to the reference-point results: the age at which fish begin entering the fishery and the relative exposure of older fish carry different information. Its mean-age measure weights selectivity equally across model ages 1–15, with age 15 representing the plus group; population abundance and catch composition are excluded from that shape index.

The reference-rate comparison makes the dependence on age selection explicit (Figure 72). Under fixed biology, each curve has its own fully selected F35% rate, the fishing-mortality scalar that reduces lifetime spawning output per recruit to 35% of the unfished level. Later exposure can allow more spawning before capture and therefore support a higher fully selected rate at the same spawning-per-recruit target. However, the full curve, including its older-age tail, determines this response. The high 2019 F35% illustrates how a reference scalar can rise sharply under a particular age pattern. Such an increase describes the rate permitted under that pattern; it provides no independent evidence of increased stock productivity or a larger sustainable catch.

Two distinctions are essential when interpreting an FMSY proxy against selected age. First, F35% is a spawning-per-recruit proxy for FMSY, whereas FMSY itself depends on the equilibrium yield and recruitment relationship. The historical FMSY comparison and this F35% diagnostic therefore represent different reference quantities. Second, fully selected F is a scalar: fishing mortality at each age equals that scalar multiplied by selectivity at that age. Comparisons among years require the same selectivity normalization and the corresponding age-specific mortality pattern. The scatterplot uses a selectivity-weighted age index over ages 1–10, whereas its spawning-per-recruit calculation includes ages 1–14 and 15+, fixed 2026 natural mortality, maturity and spawning weights, and quarter-year spawning timing. Its age index also differs from the ages 1–15 index in the preceding figure and from the abundance-weighted age of selected fish. It is a compact descriptor of curve shape, and a common mean selected age can accompany different reference rates. The complete selectivity curve and resulting catch and spawning effects provide the appropriate basis for interpreting the FMSY proxy and comparing harvest specifications.

8 Risk table and ABC recommendation

The author risk evaluation retains all four categories at **Level 1 – Normal** (Table 53). The ecosystem and fishery observations are summarized below. Working recommendations remain subject to author scientific evaluation and Plan Team, SSC and Council review. The Level 1 evaluation supports no reduction from maximum permissible ABC.

Table 53: 2026 risk-category scores and supporting summaries. Assessment-related and fishery-informed stock evaluations are Level 1 – Normal; population dynamics and ecosystem evaluations draw on the synthesis by Elizabeth Siddon and Kalei Shotwell.

Considerations			
Assessment-related	Population Dynamics	Ecosystem	Fishery-informed Stock
Level 1 – Normal	Level 1 – Normal	Level 1 – Normal	Level 1 – Normal
Model developments improved fitting, with minor implications for stock status and projected ABC/OFL.	Egg and larval abundance was low; later prey conditions were favorable and cannibalism was low.	Cool conditions, abundant prey and improving adult condition indicated favorable ecosystem conditions.	A-season fishing appeared good; later B-season fishing was farther north than normal, with limited desirable sizes.

8.1 Assessment-related considerations

Model 26.0 (Rceattle) is the adopted assessment model and main proposal for Tier 3 calculations and specifications. The alternative models examine how the assessment responds to changes in its structure and treatment of annual selectivity variation. Comparisons with Models 23.0 and 23.2 connect the current assessment to the previous formulation and help distinguish the effects of updated observations from changes in model structure. The split-fishery alternatives test whether separate selectivity patterns for the Foreign fishery in 1964–1990 and the Modern domestic fishery from 1991 onward better represent the fishing history. Further comparisons vary the constraint on declining selectivity and the treatment of year-to-year changes in selectivity.

Model 26.1.2 provides the principal sensitivity to these choices. Its underlying marginal formulation estimates the variability of Modern fishery selectivity while accounting for annual selectivity effects. The conditional comparison presented here fixes the Foreign and Modern process standard deviations at their selected full-data values and estimates the remaining parameters and annual selectivity effects. This configuration fits the Modern fishery age compositions more closely than Model 26.0, while larger discrepancies remain in the Foreign period. The result identifies a potential benefit of separating the fisheries, alongside a trade-off in how the model represents the earlier observations.

The Model 26.1.2 retrospective analysis asks whether estimates change as the most recent observations are successively removed. Each shortened-data fit holds the process standard deviations fixed and is compared with a full-data fit using the same conditional estimation method. This matching reference makes the retrospective differences interpretable within the fixed-SD configuration. The diagnostic therefore describes stability when the assumed magnitude of annual selectivity variation is held constant. Re-estimating that variation in each shortened dataset could change the retrospective pattern, and uncertainty in the process standard deviations remains outside this comparison. Improved fits to the Modern observations consequently provide one part of the model evaluation; broader retrospective reliability requires evidence across the relevant estimation choices. Model 26.0 remains the basis for the Tier 3 specifications, with Model 26.1.2 used to assess sensitivity to fishery structure and selectivity assumptions.

8.2 Population dynamics risk summary

Population dynamics risk is **Level 1 – Normal**. Cold spring conditions coincided with low egg and larval abundance; temperature-dependent timing complicates interpretation of these early-life indices. Southeastern Bering Sea (SEBS) age-0 abundance was also low. Above-average late-summer large copepod abundance, lipid-rich copepods and euphausiids, and above-average SEBS age-0 energy density indicate favorable feeding and potential survival through the first winter. Northern Bering Sea (NBS) age-0 energy density was below average, while adult-pollock cannibalism was low overall; age-1 specificity remains unverified. The regional contrasts and supporting evidence are described in Section 11.1 within Section 11.

8.3 Environmental and ecosystem risk summary

The environmental and ecosystem synthesis supports a **Level 1 – Normal** ecosystem score. Following the warm period of 2014–2021 and near-average thermal conditions in 2022–2025, the EBS shelf shifted to cool conditions in 2026. The expanded cold pool, abundant lipid-rich prey and improving adult condition provide favorable context for juvenile and adult pollock. Increased pollock biomass may increase competition; prey and condition indicators suggest adequate food, while the effects of increasing sablefish abundance remain a research question. Section 11 presents the detailed distribution, environmental, prey, competitor and predator evidence, including conditions for later juveniles and adults in Section 11.2. These category scores support the recommendation for no reduction from maximum permissible ABC.

8.4 Fishery-informed Stock Considerations

Fishing conditions differed between the two seasons. The assessment author's observations indicate that A-season fishing in 2026 appeared good. Later in the B-season, fishing occurred farther north than normal, and pollock of desirable sizes were limited. The late-season observations therefore qualify the favorable A-season picture and identify changes in where fishing occurred and the sizes available to the fishery.

We retain the **Level 1 – Normal** score for fishery-informed stock considerations. This score reflects the author's qualitative assessment of fishing conditions across the year. The observations provide context for interpreting fishery performance; their implications for stock-wide abundance remain uncertain. A standardized comparison of fishing locations, effort and size composition would help distinguish changes in fishery availability from changes in overall stock abundance.

8.5 Ecosystem and Socioeconomic Profile development

Development of the EBS pollock Ecosystem and Socioeconomic Profile (ESP) continues alongside the assessment (Shotwell et al. 2026). The archived September 2026 development report supplies exploratory background, and the AI-assisted draft appears as Appendix B. The 2026 Eastern Bering Sea Ecosystem Status Report describes an external Dynamic Structural Equation Modeling (DSEM) analysis that identified **sea ice extent**, **sea surface temperature (SST)** and **summer large copepods** as indicators with statistically supported relationships to age-1 recruitment (Siddon 2026). The exploratory Rceattle DSEM pilot in the appendix uses late-summer SST and cold-pool extent as two environmental proxies. Its scope supports a separate assessment of those proxies. Evaluating all three external relationships within Rceattle requires additional testing.

The first application of DSEM has been completed as a draft in the ESP draft. Development and implementation of an integrated version within Rceattle continue from the work presented in the September 2026 development document (Ianelli et al. 2026). We see the Ecosystem Status Report (ESR), the ESP, and integration of ecosystem information within the full assessment model as logical applications of ecosystem-based methods for providing informed catch advice to the Council.

Since all four categories are Level 1 – Normal, we recommend no reduction from maximum permissible ABC.

9 Status determination

Under the Magnuson–Stevens Fishery Conservation and Management Act, the Secretary of Commerce reports on whether each U.S. fishery is experiencing overfishing, is overfished, or is approaching an overfished condition.

As shown in the executive-summary status tables, both primary Model 26.0 and the Early-late fishery split Model 26.1.2 sensitivity support the following Tier 3 assessment conclusions:

1. **The stock was not subjected to overfishing in 2025.** The updated catch for the most recent complete year, **1,326,215.61 t** in 2025, is below the specified 2025 Bering Sea subarea OFL of **2,957,000 t** (2025 catch report, p. 1). Catch uses the updated annual estimates; the archived report provides the specified OFL, rather than the updated catch. This catch-to-OFL comparison is common to both models.
2. **The stock is not overfished in 2026.** Estimated female spawning biomass is **3.057 million t** for Model 26.0 and **3.065 million t** for the fixed-SD Model 26.1.2 comparison. These exceed their respective B35% values of **2.182 million t** and **2.204 million t**. Each model is therefore above its Tier 3 minimum stock size threshold.
3. **The stock is not approaching an overfished condition in 2026.** Scenario 7 female spawning biomass remains above each model's B35% through the two-year projection horizon ending in 2028, when it is **3.270 million t** for Model 26.0 and **3.330 million t** for the fixed-SD Model 26.1.2 comparison.

B35% is the Tier 3 BMSY proxy. Biomass above B35% satisfies the status criterion directly; the intermediate decision branch between half B35% and B35% is unnecessary for these results. The comparison below retains explicit biological years and model-specific reference points. Projections use fixed historical-mean recruitment. These are assessment conclusions for review, separate from final NMFS status determinations, scientific model adoption and adopted catch specifications.

Comparison of Model 26.0 and the sensitivity analysis			
Quantity	Year / basis	Model 26.0	26.1.2 Early-late fishery split
Estimated female spawning biomass	2026	3,057,000	3,065,000
BMSY proxy: B35%	Reference point	2,182,000	2,204,000
Lower status boundary: half B35%	Reference point	1,091,000	1,102,000
Scenario 6 female spawning biomass	2027	3,155,000	3,195,000
Scenario 7 female spawning biomass	2028	3,270,000	3,330,000
Maximum ABC: expected-catch scenario	2027	2,222,000	2,187,000
OFL: expected-catch scenario	2027	2,746,000	2,692,000
Maximum ABC: expected-catch scenario	2028	2,319,000	2,316,000
OFL: expected-catch scenario	2028	2,870,000	2,853,000
Biomasses are female spawning biomass at April 1. B35% is the Tier 3 BMSY proxy; B40% serves the ABC control rule. Half B35% is the lower status boundary. The biological-year comparisons remain separate from the national status-reporting year layout. Each model uses its own fitted population and reference points.			
Future ABC/OFL rows use each model's expected-catch scenario, with 1,375,000 t catch in each of 2027 and 2028. They are model-based maxima, separate from adopted specifications. Model 26.0 (Rceattle) is the adopted assessment model and main proposal for Tier 3 calculations and specifications; Model 26.1.2 is the sensitivity analysis. Values are rounded to the nearest 1,000 t; source exports retain their original precision. Projections condition on the revised fitted population, fixed biology and constant historical-mean recruitment.			

10 F limit

Following section 4.15 of the assessment guidelines, F limit is reverse-engineered as the annual fishing mortality that would have produced the previous year's OFL using the recommended model's estimated population for that year. For this 2026 assessment, the calculation therefore uses the adopted **2025 Bering Sea pollock OFL of 2.957 million t** (National Marine Fisheries Service (2025)) and the Model 26.0 beginning-of-year 2025 numbers at age, fishery weights, natural mortality and realized selectivity.

Solving the annual Baranov catch equation gives **F limit = 0.787 per year** on the maximum-F-across-ages scale used in this report, with the maximum at age 6. The equivalent scalar on Rceattle's native mean-normalized selectivity scale is **0.439 per year**. Both scalings give the same fishing mortality at age and reproduce the adopted 2025 OFL. This calculation holds the fitted population and biological inputs fixed while changing fishing mortality.

The calculation finds the value of F_{limit} satisfying

$$C_{2025}^{OFL} = \sum_{a=1}^{15+} N_{a,2025} w_{a,2025}^{fishery} \frac{F_{limit} s_{a,2025}^{*}}{M_{a,2025} + F_{limit} s_{a,2025}^{*}} \left[1 - \exp\left(-\left(M_{a,2025} + F_{limit} s_{a,2025}^{*}\right)\right)\right]. \quad (1)$$

Here N is beginning-of-year abundance, $w^{fishery}$ is fishery weight at age, M is annual natural mortality, and s^{*} is the fitted selectivity curve rescaled to a maximum of one. Thus F_{limit} is the maximum annual fishing mortality across ages. The sum includes ages 1–14 and the 15+ group once. Abundance in millions of fish and weight in kg give catch in thousand tons.

11 Ecosystem considerations

A draft Ecosystem and Socioeconomic Profile (ESP) is presented in Appendix B. It brings together indicators of pollock habitat, food-web conditions and fishery performance to provide context for the assessment. The ESP complements the population model by organizing evidence about processes that can affect year-class strength, fish growth and the distribution of fish available to surveys and the fishery. This broader perspective helps interpret changes in abundance and age composition alongside the environmental conditions experienced by successive cohorts.

The draft 2026 Eastern Bering Sea Ecosystem Status Report provides the environmental and food-web evidence summarized here (Siddon 2026). In 2026, cool spring conditions and an extensive cold pool coincided with low egg, larval and southeastern Bering Sea age-0 abundance. At the same time, favorable zooplankton abundance and lipid content, together with above-average southeastern Bering Sea age-0 energy density, indicated good feeding conditions for surviving juveniles. These contrasting signals emphasize the importance of timing, region and life stage: low early abundance and favorable later feeding conditions can occur within the same year. Their implications for recruitment depend on survival through the subsequent winter and later juvenile stages.

The following subsections connect these observations to pollock population processes. The first considers environmental conditions, prey, competitors and predators affecting eggs, larvae and early juveniles. The second addresses the distribution, feeding conditions and condition of later juveniles and adults, including how temperature and vertical habitat use can affect survey availability. Together, these indicators help distinguish changes in population abundance from changes in where fish occur and how effectively surveys sample them. The synthesis informs interpretation of the assessment and the risk-category evaluation summarized in Section 8; the individual indicators provide complementary evidence rather than a direct forecast of recruitment or an automatic adjustment to catch advice.

11.1 Ecosystem impacts of EBS pollock population dynamics

Environmental Processes

A very weak winter Aleutian Low in 2025/2026 brought cool Arctic air across the EBS (Lemagie and Wang 2026). **Sea ice extent** remained above average for six weeks during March–April 2026 (Thoman 2026). The resulting 2°C cold pool was approximately 150% larger than in 2025 and the fifth-largest during 1982–2026; the 0°C cold pool was the largest since 2013 (Rohan and Barnett 2026). **Sea surface temperatures** in the SEBS were colder than in 2025, especially during January–April, and remained below average for most of the year (Lemagie and Callahan 2026). In May 2026, spring winds favored Ekman transport off the shelf, influencing larval dispersal trajectories (Hennon 2026).

Prey

Large diatom anomalies were above average during 2023–2025 (Ostle and Batten 2026), and spring 2026 chlorophyll-a concentrations were average to high across the NBS and SEBS (Gann et al. 2026). These records suggest favorable prey-production conditions in their respective periods. Increased surface phosphate and a deeper mixed layer in 2024 imply wind mixing, larger phytoplankton cells, shorter food chains and more efficient energy transfer to juvenile fish (Gann 2026).

The spring Rapid Zooplankton Assessment indicated adequate small copepod abundance, with **above-average large copepod abundance by late summer** (Kimmel et al. 2026). Euphausiid abundance increased from 2025 (Levine et al. 2026; Spear et al. 2026). Copepod and euphausiid lipid content in the SEBS were both above average in 2026, indicating good prey quality for pelagic juveniles (Pinger et al. 2026).

Spring larval abundance and egg numbers were low, as was the larval condition ratio. Such observations are typical of cold spring conditions over the shelf, and temperature-dependent seasonal timing complicates their interpretation (Siddon 2026). SEBS age-0 abundance was low (Andrews, Domke, Murphy, Spear, et al. 2026), and age-0 energy density was above average (Page et al. 2026). The suggested increase in age-1 abundance in this juvenile-survey comparison remains unverified. Length–weight residuals for small pollock (100–250 mm) in the EBS survey improved to just below the long-term mean (Prohaska and Rohan 2026). In the NBS, age-0 abundance was average (Andrews, Domke, Murphy, Spear, et al. 2026) and energy density was below average (Page et al. 2026).

Competitors

Pelagic forager biomass increased from 2025 to 2026 to above its long-term mean, largely driven by a 58% increase in pollock biomass (Siddon 2026, 7–9). Above-average zooplankton quantity and quality moderate concerns about competition for prey among young pollock. SEBS jellyfish biomass remained average (Buser 2026) to below average (Domke et al. 2026). NBS jellyfish biomass continued its slow decline since 2023 and remained just above the time-series mean (Domke et al. 2026).

Pacific herring and capelin biomass were below average in both the NBS and SEBS (Andrews, Domke, Murphy, Dimond, et al. 2026). Bristol Bay sockeye salmon continued to decline from peak 2021/2022 levels (Cunningham et al. 2026). Strong 2020 and 2022 sablefish year classes contributed to a growing, aging population that could compete for pelagic juvenile prey; direct evidence for that competition remains unavailable (Siddon 2026).

Predators

Adult pollock diets consisted primarily of copepods and euphausiids, with a low overall proportion of cannibalism in 2026 (Aydin 2026). CNRG estimates show that age-1 predation mortality declined from 2025, while mortality due to cannibalism remained slightly above its long-term mean. The biomass of pollock consumed by conspecifics was approximately average and lower than in 2024 and 2025 (Holsman et al. 2026). Adult Bristol Bay sockeye salmon, which prey on juvenile pollock, decreased from 2025 (Cunningham et al. 2026).

Northern fur seal pup weights at St. Paul Island exceeded the average by more than one standard deviation, and maternal trip durations were the shortest in the time series. These indicators suggest high foraging success near the island. Age-0 pollock and herring had the highest frequency of occurrence in the September 2025 fur-seal diet samples; analysis of September 2026 samples was ongoing in the draft (McHuron et al. 2026). The apex predator guild increased from 2025, driven by a 17% increase in Pacific cod, and remained below its long-term mean (Siddon 2026, 7–9). This guild level suggests low predation pressure on juvenile pollock (Siddon 2026).

11.2 Ecosystem considerations for later juveniles and adults

Distribution

Pollock biomass estimated by the standard bottom-trawl survey increased approximately 50% from 2025 to 2026, alongside a pronounced spatial pattern associated with the extensive cold pool. Adult pollock occurred primarily outside bottom waters colder than 2°C, with concentrations along the northwestern outer shelf (Siddon 2026). The cold pool therefore delineated the near-bottom habitat occupied by much of the adult population during the survey. This pattern is relevant to interpreting the biomass increase because survey catches reflect both the number and size of fish present and their availability to bottom-trawl gear. The higher survey estimate and the concentration of adults along the cold-pool margin should be considered together when comparing years. The spatial association supports a role for temperature in shaping survey availability, while the contribution of redistribution to the biomass increase remains unresolved.

Environmental Processes

Bottom temperatures in 2026 were comparable to the historical cool years 1986, 1992, 2006 and 2010 (Siddon 2026). The large cold pool provided further evidence of cool bottom conditions (Rohan and Barnett 2026). **Below-average SSTs** characterized the SEBS, particularly during spring (Lemagie and Callahan 2026). Adult pollock condition was above its long-term mean (Prohaska and Rohan 2026). These temperature and condition patterns describe concurrent conditions; their causal relationship remains an interpretation.

Prey

Adult pollock diets continued to be dominated by **copepods** and euphausiids (Aydin 2026). Summer and late-summer euphausiid abundance increased from 2025 to 2026 (Levine et al. 2026; Spear et al. 2026), and above-average SEBS copepod and euphausiid lipid content provided lipid-rich prey (Pinger et al. 2026). Cannibalism formed a low proportion of adult pollock diets (Aydin 2026). CNRG estimates separately indicate slightly above-average mortality of age-1 pollock from cannibalism, with approximately average biomass consumed by conspecifics (Holsman et al. 2026).

Morphometric condition, measured as length–weight residuals, for large adult pollock (>250 mm) in the EBS survey increased from 2023 to 2026 to above the long-term mean. The NBS condition series extends through 2025, when large-pollock condition remained positive and continued the increasing trend observed since 2021 (Prohaska and Rohan 2026).

Competitors

The increase in pelagic forager biomass from 2025 to 2026 was primarily driven by pollock (Siddon 2026, 7–9), suggesting a potential increase in competition within the pollock population. Above-average prey quantity, prey quality and pollock length–weight residuals indicate adequate food resources. Increasing sablefish abundance represents potential competition for pelagic prey, with direct evidence still pending (Siddon 2026).

Predators

The apex predator guild, driven by Pacific cod, increased from 2025 and remained below its long-term mean (Siddon 2026, 7–9). This pattern suggests low predation pressure on pollock (Siddon 2026). These associations provide context for the stock; independent evidence is needed to establish the underlying mechanisms. The Plan Team requested further examination of ecosystem contributions to retrospective behavior alongside the direct data-influence tests.

12 Data gaps and research priorities

The 2026 assessment compares the separately fitted configurations and the Model 26.1.2 marginal-estimation sensitivity using their matching observations and conditional observation diagnostics. Model 26.0 posterior uncertainty, predictive observation checks and nine retrospective fits are complete. The Model 26.1.2 retrospective diagnostic uses fixed full-model process SDs and a conditional full 2026 reference. The two Model 26.0 simulation-retrospective experiments compare conditional biomass and recruitment revisions with the observed-data results (Section 6.3.1). Each experiment uses ten simulated datasets to examine biomass and recruitment revisions.

Additional age-1 BTS selectivity and BTS data-influence tests are outside the scope of this assessment (PT-02 and PT-05). The comparisons describe differences in retrospective patterns. The weight-at-age comparison in Section 5.2 retains the distinction between estimated schedules and uncertainty carried into population and projection outputs.

A high research priority is to refine the statistical treatment of length compositions and age observations from the fishery, bottom-trawl survey (BTS), and acoustic-trawl survey (ATS), together with paired Fourier-transform near-infrared spectroscopy (FT-NIR) and traditional morphological ageing (TMA) observations. A more statistically rigorous framework should account for sampling structure, ageing error, uncertainty in conversions between ageing methods, and dependence among observations derived from the same samples. This work should guide composition weighting and the integration of length and age information, with the resulting methods and uncertainty treatment carried into future model specifications.

The treatment of early fishery data should also be refined relative to more recent observations. Research should evaluate changes in sampling coverage, age and length information, and fishing practices across periods, and use those differences to inform selectivity assumptions and observation weights. Comparisons should assess how alternative treatments affect recruitment, biomass trends, and retrospective behavior.

Further development of random-effects models should address temporal variation in selectivity and other population or observation processes. Priorities include identifying which sources of variability the data can resolve, evaluating correlation across ages and years, and estimating process variation within a coherent framework. Simulation tests, retrospective analyses, and predictive checks should evaluate model performance and the propagation of uncertainty into stock status and catch projections.

The Ecosystem and Socioeconomic Profile (ESP) provides a basis for fuller implementation and use of dynamic structural equation models (DSEM). This work should develop and test hypothesized links among environmental conditions, prey, pollock condition, and recruitment, while accounting for observation uncertainty, missing years, and time lags. Evaluation should establish the stability and predictive value of these relationships and how ESP results can inform assessment interpretation, recruitment scenarios, and future model specifications.

Additional research questions include AVO uncertainty, the 2018 recruitment profile, and stock boundaries and connectivity.

13 Acknowledgements

We thank the scientifically trained observers and AFSC fisheries monitoring staff for their hard work and dedication. Survey staff contribute immensely through their diligence in collecting the samples that support this assessment, and their commitment under challenging circumstances is exceptional. We also thank the AFSC age-and-growth staff for their continued excellence in promptly processing these samples.

We thank the many colleagues who provided edits and suggestions to improve this document, particularly Dr. Cole Monnahan for his timely and thoughtful review.

14 Assessment source archive

The private source archive retains the guidelines and Plan Team excerpt, a downloaded copy of the public September report, the local September-development source and data products, and the previous assessment chapter sources. Main tables and figures follow Literature cited in separate sections; appendix-specific objects remain with their appendices. The final SAFE package will retain its table and figure inventory and electronic assessment outputs.

The report draws on saved assessment and survey results. The technical appendix documents the software, source identities and steps needed to reproduce those results (the technical supplement). Scientific review and accessibility checks retain their separate status.

Literature cited

AFSC-RACE MACE. 2026. *Preliminary Results, AVO 2026*. Unpublished report supplied privately.

Alaska Fisheries Science Center, National Marine Fisheries Service. 2025. *Assessment of Pollock in the Bering Sea and Aleutian Islands*. Bering Sea and Aleutian Islands SAFE catch report. North Pacific Fishery Management Council. https://files.npfmc.org/SAFE/2025/BSAIpollock.pdf.

Anderson, S. C., E. J. Ward, P. A. English, and L. A. K. Barnett. 2022. “sdmTMB: An r Package for Fast, Flexible, and User-Friendly Generalized Linear Mixed Effects Models with Spatial and Spatiotemporal Random Fields.” *BioRxiv*, 2022–03. https://doi.org/10.1101/2022.03.24.485545.

Andrews, A., L. Domke, J. Murphy, A. Dimond, C. Lawrence, and R. Suryan. 2026. “Pacific Herring and Capelin from Surface Trawl Surveys, 2003–2026.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Andrews, A., L. Domke, J. Murphy, A. Spear, A. Dimond, and C. Lawrence. 2026. “Age-0 Pollock and Pacific Cod from Surface Trawl Surveys, 2003–2026.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Aydin, K. 2026. “Summer Food Habits of Walleye Pollock and Pacific Cod.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Bacheler, N. M., L. Ciannelli, K. M. Bailey, and J. T. Duffy-Anderson. 2010. “Spatial and Temporal Patterns of Walleye Pollock (*Theragra Chalcogramma*) Spawning in the Eastern Bering Sea Inferred from Egg and Larval Distributions.” *Fish. Oceanogr.* 19 (2): 107–20.

Bailey, K. M. 2000. “Shifting Control of Recruitment of Walleye Pollock *Theragra Chalcogramma* After a Major Climatic and Ecosystem Change.” *Mar. Ecol. Prog. Ser.,* 198: 215–24. https://doi.org/10.3354/meps198215.

Bailey, K. M., T. J. Quinn, P. Bentzen, and W. S. Grant. 1999. “Population Structure and Dynamics of Walleye Pollock, *Theragra Chalcogramma*.” *Advances in Mar. Biol.* 37: 179–255.

Barbeaux, S. J., S. Gaichas, J. N. Ianelli, and M. W. Dorn. 2005. “Evaluation of Biological Sampling Protocols for at-Sea Groundfish Observers in Alaska.” *Alaska Fisheries Research Bulletin* 11 (2): 82–101.

Brodeur, R. D., M. T. Wilson, L. Ciannelli, M. J. Doyle, and J. M. Napp. 2002. “Interannual and Regional Variability in Distribution and Ecology of Juvenile Pollock and Their Prey in Frontal Structures of the Bering Sea.” *Deep-Sea Research II* 49: 6051–67.

Buckley, T. W., A. Greig, and J. L. Boldt. 2009. *Describing Summer Pelagic Habitat over the Continental Shelf in the Eastern Bering Sea, 1982–2006*. NMFS-AFSC-196. United States Department of Commerce, NOAA Technical Memorandum.

Buckley, T. W., I. Ortiz, S. Kotwicki, and K. Aydin. 2015. “Summer Diet Composition of Walleye Pollock and Predator-Prey Relationships with Copepods and Euphausiids in the Eastern Bering Sea, 1987-2011.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 302–11. http://doi.org/10.1016/j.dsr2.2015.10.009.

Buser, T. 2026. “Eastern and Northern Bering Sea – Jellyfishes.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Butterworth, D. S., J. N. Ianelli, and R. Hilborn. 2003. “A Statistical Model for Stock Assessment of Southern Bluefin Tuna with Temporal Changes in Selectivity.” *African Journal of Marine Science* 25 (1): 331–61. https://doi.org/10.2989/18142320309504021.

Canino, M. F., P. T. O’Reilly, L. Hauser, and P. Bentzen. 2005. “Genetic Differentiation in Walleye Pollock (*Theragra Chalcogramma*) in Response to Selection at the Pantophysin (Pan i) Locus.” *Can. J. Fish. Aquat. Sci.* 62: 2519–29.

Ciannelli, L., R. D. Brodeur, and J. M. Napp. 2004. “Foraging Impact on Zooplankton by Age-0 Walleye Pollock (*Theragra Chalcogramma*) Around a Front in the Southeast Bering Sea.” *Marine Biology* 144: 515–25.

Cunningham, C. J., S. Vega, and J. Terry-Shindelman. 2026. “Temporal Trend in the Annual Inshore Run Size of Bristol Bay Sockeye Salmon (Oncorhynchus nerka).” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Domke, L., A. Andrews, A. Dimond, C. Lawrence, and J. Murphy. 2026. “Jellyfish from Surface Trawl Surveys, 2004–2026.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Duffy-Anderson, J. T., S. J. Barbeaux, E. Farley, et al. 2016. “The Critical First Year of Life of Walleye Pollock (*Gadus Chalcogrammus*) in the Eastern Bering Sea: Implications for Recruitment and Future Research.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 283–301. http://doi.org/10.1016/j.dsr2.2015.02.001.

Fournier, D. A., J. R. Sibert, J. Majkowski, and J. Hampton. 1990. “MULTIFAN a Likelihood-Based Method for Estimating Growth Parameters and Age Composition from Multiple Length Frequency Samples with an Application to Southern Bluefin Tuna (Thunnus Maccoyii).” *Can. J. Fish. Aquat. Sci.* 47: 301–17.

Fournier, D., and C. Archibald. 1982. “A General Theory for Analyzing Catch at Age Data.” *Canadian Journal of Fisheries and …*. http://www.nrcresearchpress.com/doi/abs/10.1139/f82-157.

Francis, R. I. C. C. 1992. “Use of Risk Analysis to Assess Fishery Management Strategies: A Case Study Using Orange Roughy (Hoplostethus Atlanticus) on the Chatham Rise, New Zealand.” *Can. J. Fish. Aquat. Sci.* 49: 922–30.

Francis, R. I. C. C. 2011. “Data Weighting in Statistical Fisheries Stock Assessment Models.” *Canadian Journal of Fisheries and Aquatic Sciences* 68: 1124–38. https://doi.org/10.1139/F2011-025.

Gann, J. C. 2026. “Surface Phosphate Concentrations and Mixed Layer Depth Correlate with Larger Phytoplankton in the Eastern Bering Sea.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Gann, J. C., M. W. Callahan, J. M. Nielsen, C. W. Mordy, S. W. Bell, and P. Stabeno. 2026. “Spring Satellite Chlorophyll-a Concentrations.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Gann, J. C., L. B. Eisner, S. Porter, et al. 2015. “Possible Mechanism Linking Ocean Conditions to Low Body Weight and Poor Recruitment of Age-0 Walleye Pollock (*Gadus Chalcogrammus*) in the Southeast Bering Sea During 2007.” *Deep Sea Research Part II: Topical Studies in Oceanography* 134: 1–13. http://doi.org/10.1016/j.dsr2.2015.07.010.

Guthrie, C. M., III, Hv. T. Nguyen, A. E. Thomson, and J. R. Guyon. 2017. *Genetic Stock Composition Analysis of the Chinook Salmon Bycatch from the 2015 Bering Sea Walleye Pollock (Gadus chalcogrammus) Trawl Fishery*. NOAA Technical Memorandum NMFS-AFSC-342. U.S. Department of Commerce, National Oceanic; Atmospheric

Administration, National Marine Fisheries Service, Alaska Fisheries Science Center. https://doi.org/10.7289/V5/TM-AFSC-342.

Heintz, R. a., E. C. Siddon, E. V. Farley, and J. M. Napp. 2013. "Correlation Between Recruitment and Fall Condition of Age-0 Pollock (*Theragra Chalcogramma*) from the Eastern Bering Sea Under Varying Climate Conditions." *Deep Sea Research Part II: Topical Studies in Oceanography* 94: 150–56. https://doi.org/10.1016/j.dsr2.2013.04.006.

Hennon, T. 2026. "Winds at the Shelf Break." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Hilborn, R., and C. J. Walters. 1992. *Quantitative Fisheries Stock Assessment*. Springer US. https://doi.org/10.1007/978-1-4615-3598-0.

Holsman, K. K., J. Ianelli, K. Shotwell, et al. 2026. "Bering Sea Multispecies Model (CNRG) Estimates of Time-Varying Natural Mortality." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Honkalehto, T., and A. McCarthy. 2015. *Results of the Acoustic-Trawl Survey of Walleye Pollock (Gaddus Chalcogrammus) on the u.s. And Russian Bering Sea Shelf in June - August 2014*. AFSC Processed Rep. 2015-07, 62 p. Alaska Fish. Sci. Cent., NOAA, Natl. Mar. Fish. Serv., 7600 Sand Point Way NE, Seattle WA 98115. http://www.afsc.noaa.gov/Publications/ProcRpt/PR2015-07.pdf.

Honkalehto, T., P. H. Ressler, R. Towler, and C. D. Wilson. 2011. "Using Acoustic Data from Fishing Vessels to Estimate Walleye Pollock Abundance in the Eastern Bering Sea." *Can. J. Fish. Aquat. Sci.* 68: 1231–42.

Hunt Jr., G. L., K. O. Coyle, L. B. Eisner, et al. 2011. "Climate Impacts on Eastern Bering Sea Foodwebs: A Synthesis of New Data and an Assessment of the Oscillating Control Hypothesis." *ICES J. Mar. Sci.* 68 (6): 1230–43. http://dx.doi.org/10.1093/icesjms/fsr036.

Ianelli, J. 2026. *AKsf: Alaska Fisheries Spatial Features*. R package. Version 0.1.0. https://jimianelli.github.io/AKsf/.

Ianelli, J. N., and D. A. Fournier. 1998. *Alternative Age-Structured Analyses of the NRC Simulated Stock Assessment Data*. NOAA Tech. Memo. NMFS-F/SPO-30. NOAA.

Ianelli, J., T. Honkalehto, Allen-Akselrud C., S. Stienessen, and E. Siddon. 2021. *Chapter 1 : Assessment of the Walleye Pollock Stock in the Eastern Bering Sea*. (1007 West Third, Suite 400, Anchorage, Alaska 99501). https://apps-afsc.fisheries.noaa.gov/Plan_Team/2021/EBSPollock.pdf.

Ianelli, J., T. Honkalehto, Wasserman S., N. Lauffenburger, McGilliard C., and E. Siddon. 2023. *Chapter 1 : Assessment of the Walleye Pollock Stock in the Eastern Bering Sea*. (1007 West Third, Suite 400, Anchorage, Alaska 99501). https://apps-afsc.fisheries.noaa.gov/Plan_Team/2023/EBSPollock.pdf.

Ianelli, J., T. Honkalehto, S. Wassermann, et al. 2024. *Assessment of Walleye Pollock in the Eastern Bering Sea*. North Pacific Fishery Management Council. https://files.npfmc.org/SAFE/2024/EBSpollock.pdf.

Ianelli, J., S. Kotwicki, and T. Honkalehto. 2015. *Chapter 1 : Assessment of the Walleye Pollock Stock in the Eastern Bering Sea*. (Anchorage, Alaska), no. December.

Ianelli, J., S. Wassermann, C. McGilliard, et al. 2026. *September 2026 Assessment Developments: Eastern Bering Sea Walleye Pollock*. Assessment development working paper; Alaska Fisheries Science Center, National Marine Fisheries Service. https://noaa-afsc.github.io/EBS_pollock/doc/EBSpollock_Sept_2026.html.

Ianelli, S. B., James N., and N. Williamson. 2001. *Chapter 1 : Assessment of the Walleye Pollock Stock in the Eastern Bering Sea*. (Anchorage, Alaska), no. December.

Indivero, J., T. E. Essington, J. N. Ianelli, and J. T. Thorson. 2023. "Incorporating distribution shifts and spatio-temporal variation when estimating weight-at-age for stock assessments: a case study involving the Bering Sea pollock (*Gadus chalcogrammus*) ." *ICES Journal of Marine Science* 80 (2): 258–71. https://doi.org/10.1093/icesjms/fsac236.

Jost, L. 2006. "Entropy and Diversity." *Oikos* 113 (2): 363–75. https://doi.org/10.1111/j.2006.0030-1299.14714.x.

Kimmel, D. G., C. E. Harpold, J. Keister, et al. 2026. "Current and Historical Trends for Zooplankton in the Bering Sea." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Kimura, D. K. 1989. "Variability in Estimating Catch-in-Numbers-at-Age and Its Impact on Cohort Analysis." In *Effects on Ocean Variability on Recruitment and an Evaluation of Parameters Used in Stock Assessment Models*, edited by R. J. Beamish and G. A. McFarlane. Can. Spec. Publ. Fish. Aq. Sci.

Lang, G. M., R. D. Brodeur, J. M. Napp, and R. Schabetsberger. 2000. “Variation in Groundfish Predation on Juvenile Walleye Pollock Relative to Hydrographic Structure Near the Pribilof Islands, Alaska.” *ICES Journal of Marine Science* 57: 265–71.

Lang, G. M., P. A. Livingston, and K. A. Dodd. 2005. *Groundfish Food Habits and Predation on Commercially Important Prey Species in the Eastern Bering Sea from 1997 Through 2001*. No. 158. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC. http://www.afsc.noaa.gov/Publications/AFSC-TM/NOAA-TM-AFSC-158.pdf.

Lauffenburger, N., T. Honkalehto, S. C. Stienessen, and D. Stevenson. 2025. *Acoustic Vessel-of-Opportunity (AVO) Index for Midwater Bering Sea Walleye Pollock: A Reanalysis and New Results for 2022–2023*. AFSC Processed Report Nos. 2025-09. Alaska Fisheries Science Center, NOAA, National Marine Fisheries Service. https://repository.library.noaa.gov/view/noaa/71126/noaa_71126_DS1.pdf.

Lemagie, E., and M. Callahan. 2026. “Marine Heatwave Index.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Lemagie, E., and M. Wang. 2026. “State of the North Pacific Ocean.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Levine, M., A. De Robertis, P. Ressler, and B. Lucca. 2026. “Summer Eastern Bering Sea Euphausiids (‘Krill’).” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Livingston, P. A. 1991. *Walleye Pollock*. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-F/NWC-207.

McAllister, M. K., and J. N. Ianelli. 1997. “Bayesian Stock Assessment Using Catch-Age Data and the Sampling - Importance Resampling Algorithm.” *Canadian Journal of Fisheries and Aquatic Sciences* 54 (2): 284–300. https://doi.org/10.1139/f96-285.

McHuron et al. 2026. “Integrated Marine Mammal Information.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Monnahan, C., J. Sullivan, and S. Barbeaux. 2026. *afscOSA: Calculate and Visualize Fits to AFSC Compositional Stock Assessment Data*. https://github.com/noaa-afsc/afscOSA.

Monnahan, C., J. T. Thorson, S. Kotwicki, N. Lauffenburger, J. N. Ianelli, and A. E. Punt. 2021. “Incorporating Vertical Distribution in Index Standardization Accounts for Spatiotemporal Availability to Acoustic and Bottom Trawl Gear for Semi-Pelagic Species.” *ICES Journal of Marine Science*, ahead of print, May 19. https://doi.org/10.1093/icesjms/fsab085.

National Marine Fisheries Service. 2007. *Fisheries of the Exclusive Economic Zone Off Alaska; Prohibited Species Bycatch Management*. Federal Register 72:61070–61076, October 29. Final rule implementing Amendment 84; effective November 28, 2007. https://www.govinfo.gov/content/pkg/FR-2007-10-29/pdf/E7-21256.pdf.

National Marine Fisheries Service. 2010. *Fisheries of the Exclusive Economic Zone Off Alaska; Chinook Salmon Bycatch Management in the Bering Sea Pollock Fishery*. Federal Register 75:53026–53074, August 30. Final rule implementing Amendment 91; effective September 29, 2010. https://www.govinfo.gov/content/pkg/FR-2010-08-30/pdf/2010-20618.pdf.

National Marine Fisheries Service. 2016. *Fisheries of the Exclusive Economic Zone Off Alaska; Bycatch Management in the Bering Sea Pollock Fishery*. Federal Register 81:37534–37556, June 10. Final rule implementing Amendment 110; effective July 11, 2016. https://www.govinfo.gov/content/pkg/FR-2016-06-10/pdf/2016-13697.pdf.

National Marine Fisheries Service. 2023. *Notice of Intent to Prepare an Environmental Impact Statement for Minimizing Non-Chinook Salmon Bycatch in the Bering Sea Pollock Fishery in the Bering Sea/Aleutian Islands Fishery Management Plan Area*. Federal Register 88:44096–44100, July 11. Historical background in an environmental-impact-statement notice; not implementation of a new salmon-avoidance measure. https://www.govinfo.gov/content/pkg/FR-2023-07-11/pdf/2023-14581.pdf.

National Marine Fisheries Service. 2025. *Fisheries of the Exclusive Economic Zone Off Alaska; Bering Sea and Aleutian Islands; Final 2025 and 2026 Harvest Specifications for Groundfish*. Federal Register 90(51):12640, document 2025-04406. https://www.govinfo.gov/content/pkg/FR-2025-03-18/pdf/2025-04406.pdf.

National Marine Fisheries Service. 2026a. *Bering Sea and Aleutian Islands; 2026 and 2027 Harvest Specifications for Groundfish*. Federal Register 91:11750, March 10, document 2026-04684. https://www.govinfo.gov/content/pkg/FR-2026-03-10/pdf/2026-04684.pdf.

National Marine Fisheries Service. 2026b. *Pollock Fishing in the Winter Herring Savings Area of the Bering Sea and Aleutian Islands Management Area*. Federal Register 91:62343, October 1, document 2026-20135. https://www.govinfo.gov/content/pkg/FR-2026-10-01/pdf/2026-20135.pdf.

National Marine Fisheries Service. 2026c. *Reallocation of Pollock in the Bering Sea and Aleutian Islands*. Federal Register 91:51598–51599, August 11, document 2026-16336. https://www.govinfo.gov/content/pkg/FR-2026-08-11/pdf/2026-16336.pdf.

National Marine Fisheries Service Alaska Region. 2026a. *Bering Sea Aleutian Islands Catch Report, Includes CDQ: 2026*. https://www.fisheries.noaa.gov/sites/default/files/akro/car110_bsai_with_cdq2026.html.

National Marine Fisheries Service Alaska Region. 2026b. *Bering Sea and Aleutian Islands Inseason Management Report: December 2025*. https://www.fisheries.noaa.gov/s3/2026-02/BSAI-Inseason-Report-akro-2025.pdf.

National Marine Fisheries Service Alaska Regional Office. 2025. *Electronic Monitoring in Alaska*. https://www.fisheries.noaa.gov/alaska/resources-fishing/electronic-monitoring-alaska.

O'Leary, C. A., L. B. DeFilippo, J. T. Thorson, et al. 2022. "Understanding Transboundary Stocks' Availability by Combining Multiple Fisheries-Independent Surveys and Oceanographic Conditions in Spatiotemporal Models." *ICES Journal of Marine Science* 79 (4): 1063–74. https://doi.org/10.1093/icesjms/fsac046.

O'Leary, C. A., J. T. Thorson, J. N. Ianelli, and S. Kotwicki. 2020. "Adapting to Climate-driven Distribution Shifts Using Model-based Indices and Age Composition from Multiple Surveys in the Walleye Pollock (Gadus Chalcogrammus) Stock Assessment." *Fisheries Oceanography* 29 (6): 541–57. https://doi.org/10.1111/fog.12494.

Ostle, C., and S. Batten. 2026. "Continuous Plankton Recorder Data from the Eastern Bering Sea." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Page, J., J. Maselko, R. Suryan, E. Fergusson, and E. Siddon. 2026. "Fall Condition of Young-Of-The-Year Walleye Pollock in the Northern and Southeastern Bering Sea, 2002–2026." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Parma, A. M. 1993. "Retrospective Catch-at-Age Analysis of Pacific Halibut: Implications on Assessment of Harvesting Policies." *Proceedings of the International Symposium on Management Strategies of Exploited Fish Populations*.

Petitgas, P. 1993. "Geostatistics for Fish Stock Assessments: A Review and an Acoustic Application." *ICES J. Mar. Sci.* 50: 285–98.

Pinger, C., B. Cormack, C. Harpold, et al. 2026. "Lipid Content of Copepods and Euphausiids in the Bering Sea." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Prohaska, B., and S. Rohan. 2026. "Eastern and Northern Bering Sea Groundfish Condition." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Rohan, S. K., L. A. K. Barnett, and N. Charriere. 2022. *Evaluating Approaches to Estimating Mean Temperatures and Cold Pool Area from AFSC Bottom Trawl Surveys of the Eastern Bering Sea*. NOAA Technical Memorandum NMFS-AFSC-456. U.S. Department of Commerce, NOAA. https://doi.org/10.25923/1wwh-q418.

Rohan, S., and L. Barnett. 2026. "Cold Pool Extent - AFSC Bottom Trawl Survey." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Schnute, J. T., and L. J. Richards. 1995. "The Influence of Error on Population Estimates from Catch-Age Models." *Canadian Journal of Fisheries and Aquatic Sciences* 52 (10): 2063–77. https://doi.org/10.1139/f95-800.

Shotwell, S. K., J. Ianelli, and E. Siddon. 2026. *Ecosystem and Socioeconomic Profile of the Pollock Stock in the Eastern Bering Sea*. Draft Ecosystem and Socioeconomic Profile. North Pacific Fishery Management Council.

Siddon, E. 2026. *Ecosystem Status Report 2026: Eastern Bering Sea*. Stock Assessment and Fishery Evaluation Report. North Pacific Fishery Management Council.

Spear, A., D. Kimmel, J. Lamb, M. Levine, and P. Ressler. 2026. "Late Summer Eastern Bering Sea Euphausiids ('Krill')." In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Stewart, I. J., and C. C. Monnahan. 2025. "Diagnosing Common Sources of Lack of Fit to Composition Data in Fisheries Stock Assessment Models Using One-Step-Ahead (OSA) Residuals." *Canadian Journal of Fisheries and Aquatic Sciences* 82: 1–13. https://doi.org/10.1139/cjfas-2025-0158.

Thoman, R. 2026. “Bering Sea Daily Ice Extent.” In *Ecosystem Status Report 2026: Eastern Bering Sea*, edited by E. Siddon. North Pacific Fishery Management Council.

Urmy, S. S., P. Ressler, and A. De Robertis. 2025. “Estimating Uncertainty from Multiple Sources in Acoustic-Trawl Surveys.” *Unpublished Manuscript*, May.

Walline, P. D. 2007. “Geostatistical Simulations of Eastern Bering Sea Walleye Pollock Spatial Distributions, to Estimate Sampling Precision.” *ICES Journal of Marine Science* 64: 559–69.

Williamson, N., and J. Traynor. 1996. “Application of a One-Dimensional Geostatistical Procedure to Fisheries Acoustic Surveys of Alaskan Pollock.” *ICES Journal of Marine Science* 53: 423–28.

Tables

Table 1: Catch and adopted harvest limits for EBS pollock, 1964–2026 (round-weight metric tonnes). Catch from 1991 onward uses the catch estimates; the highlighted 2026 value is the selected annual model input, with the extract coverage cutoff unspecified.

EBS catch and adopted harvest limits				
Year	OFL (t)	ABC (t)	TAC (t)	Catch (t)
1964	—	—	—	174,792
1965	—	—	—	230,551
1966	—	—	—	261,678
1967	—	—	—	550,362
1968	—	—	—	702,181
1969	—	—	—	862,789
1970	—	—	—	1,256,565
1971	—	—	—	1,743,763
1972	—	—	—	1,874,534
1973	—	—	—	1,758,919
1974	—	—	—	1,588,390
1975	—	—	—	1,356,736
1976	—	—	—	1,177,822
1977	—	950,000	950,000	978,370
1978	—	950,000	950,000	979,431
1979	—	1,100,000	950,000	935,714
1980	—	1,300,000	1,000,000	958,280
1981	—	1,300,000	1,000,000	973,502
1982	—	1,300,000	1,000,000	955,964
1983	—	1,300,000	1,000,000	981,450
1984	—	1,300,000	1,200,000	1,092,055
1985	—	1,300,000	1,200,000	1,139,676
1986	—	1,300,000	1,200,000	1,141,993
1987	—	1,300,000	1,200,000	859,416
1988	—	1,500,000	1,300,000	1,228,721
1989	—	1,340,000	1,340,000	1,229,600
1990	—	1,450,000	1,280,000	1,455,193
1991	—	1,676,000	1,300,000	1,195,644
1992	1,770,000	1,490,000	1,300,000	1,390,299
1993	1,340,000	1,340,000	1,300,000	1,326,602
1994	1,590,000	1,330,000	1,330,000	1,329,352
1995	1,500,000	1,250,000	1,250,000	1,264,247
1996	1,460,000	1,190,000	1,190,000	1,192,781
1997	1,980,000	1,130,000	1,130,000	1,124,433
1998	2,060,000	1,110,000	1,110,000	1,102,159
1999	1,720,000	992,000	992,000	989,680
2000	1,680,000	1,139,000	1,139,000	1,132,710
2001	3,536,000	1,842,000	1,400,000	1,387,197
2002	3,530,000	2,110,000	1,485,000	1,480,774
2003	3,530,000	2,330,000	1,491,760	1,490,779
2004	2,740,000	2,560,000	1,492,000	1,480,552

EBS catch and adopted harvest limits				
Year	OFL (t)	ABC (t)	TAC (t)	Catch (t)
2005	2,100,000	1,960,000	1,478,500	1,483,022
2006	2,090,000	1,930,000	1,485,000	1,488,031
2007	1,640,000	1,394,000	1,394,000	1,354,502
2008	1,440,000	1,000,000	1,000,000	990,578
2009	977,000	815,000	815,000	810,784
2010	918,000	813,000	813,000	810,186
2011	2,450,000	1,270,000	1,252,000	1,199,031
2012	2,474,000	1,220,000	1,200,000	1,205,214
2013	2,550,000	1,375,000	1,247,000	1,270,760
2014	2,795,000	1,369,000	1,267,000	1,297,417
2015	3,330,000	1,637,000	1,310,000	1,321,574
2016	3,910,000	2,090,000	1,340,000	1,352,681
2017	3,640,000	2,800,000	1,345,000	1,359,181
2018	4,797,000	2,592,000	1,364,341	1,379,288
2019	3,914,000	2,163,000	1,397,000	1,409,346
2020	4,085,000	2,043,000	1,425,000	1,367,236
2021	2,594,000	1,626,000	1,375,000	1,376,258
2022	1,469,000	1,111,000	1,111,000	1,105,419
2023	3,381,000	1,910,000	1,300,000	1,310,592
2024	3,162,000	2,313,000	1,313,580	1,311,261
2025	2,957,000	2,417,000	1,389,000	1,326,216
2026	2,496,000	2,036,000	1,389,500	1,312,505
Catch for 1991–2026 uses the revised annual estimates. Model inputs use the same totals at full precision, converted to thousand tonnes. Pre-1991 catch and adopted harvest limits retain their previous sources. Dashes mean unavailable.				
All supplied Bering Sea pollock retained/discarded, gear and sector records are summed once. Explicit Bogoslof (518) and outside-stock area codes are excluded. The 23 records with blank reporting-area codes in 1998 retain their Bering Sea classification and contribute 83,077.58 t. The year-by-year reconciliation is in the catch-revision table.				
The 2026 CSV total is 1,312,504.580845496 t and is used as the annual input under the author's instruction. The CSV supplies no within-year catch cutoff; the author has confirmed this CSV as the correct annual input version. The previous 1,310,000-t forecast and October 17 catch-to-date are archived in the revision record. Annual source totals.				

Table 2: Selected management and monitoring milestones affecting the EBS pollock fishery, 1977–2017. Dates distinguish Council recommendations, enactment, voluntary programs and regulatory implementation. BSAI- or GOA-wide scope is stated where applicable; the table is selective rather than a complete regulatory history.

Historical pollock management and monitoring measures			
Year	Action stage	Measure and scope	Sources
1977	Implemented	Preliminary BSAI fishery management plan (FMP), including closure areas.	SAFE 2024; History 1997
1982	Implemented	BSAI groundfish FMP implemented; Chinook salmon bycatch limits established for foreign trawlers.	SAFE 2024; History 1997
1984	Implemented	BSAI groundfish optimum yield capped at 2 million t; Chinook salmon bycatch limits for foreign trawling reduced.	SAFE 2024; History 1997
1990	Implemented	Domestic observer program and associated data reporting began.	SAFE 2024; NOAA history
1992	Implemented	Pollock allocations under the Western Alaska Community Development Quota (CDQ) program began.	SAFE 2024

Historical pollock management and monitoring measures			
Year	Action stage	Measure and scope	Sources
1993–1994	Donation program	Prohibited Species Donation Program established in 1993; voluntary salmon donations in 1994 are reported in the accepted assessment history.	Council overview; SAFE 2024
1994	Council recommendation	Council recommended minimum codend mesh/design standards; an implemented NMFS mesh rule is not established by the reviewed sources.	History 1997; Gear review
1994	Reporting	NMFS began publishing individual-vessel bycatch rates online.	SAFE 2024; History 1997
1994–1995	Implemented	Emergency Chum Salmon Savings Area in 1994; permanent chum savings-area provisions under Amendment 35 in 1995, alongside the historical Chinook closure/trigger regime.	2023 notice history; SAFE 2024
1998	Implemented	Amendment 49 improved retention/utilization requirements for pollock and Pacific cod took effect January 3.	Amendment 49
1998	Enacted	American Fisheries Act (AFA) enacted, establishing the framework for sector allocations and fishery cooperatives.	SAFE 2024; AFA rule
1999	Implemented	AFA provisions implemented for catcher/processors.	SAFE 2024; AFA rule
1999	Emergency rule	Temporal/spatial pollock dispersal measures and trawl exclusion zones around selected Steller sea lion rookeries and haulouts in BSAI and GOA.	1999 emergency rule
2000	Implemented	AFA provisions implemented for the remaining catcher-vessel and mothership sectors.	SAFE 2024; AFA rule
2001	Voluntary	Pollock industry began the chum salmon rolling-hotspot program.	2023 notice history; SAFE 2024
2002	Voluntary	Pollock industry began the Chinook salmon rolling-hotspot program.	2023 notice history; SAFE 2024
2005	Council action	Council selected Amendment 84 in October to provide a salmon bycatch intercooperative agreement (ICA) alternative to fixed savings-area closures.	Amendment 84
2006	Exempted fishing	The B-season exempted fishing permit tested the ICA/rolling-hotspot approach before general regulatory implementation.	Amendment 84
2007	Implemented	Amendment 84 regulations, effective November 28, exempted pollock vessels participating in an approved salmon bycatch ICA from specified savings-area closures.	Amendment 84
2010–2011	Rule / program	Amendment 91 final rule effective September 29, 2010; Chinook PSC limits, incentive plan agreements (IPAs), performance standards and enhanced monitoring operated in the 2011 fishery.	Amendment 91
2015	Council action	Council took final action on Amendment 110 in April, recommending linked Chinook/chum avoidance measures and abundance-based Chinook limits.	Amendment 110

Historical pollock management and monitoring measures			
Year	Action stage	Measure and scope	Sources
2016	Implemented	Amendment 110 final rule published June 10, effective July 11. It strengthened IPAs and changed pollock seasonal allocations to 45% A season / 55% B season.	Amendment 110
2017	First index year	First fishing year informed by the Amendment 110 three-system abundance index supplied in 2016. Low-abundance limits apply when the index is at or below 250,000 Chinook salmon.	Amendment 110
Historical entries were checked against accepted 2024 SAFE Table 4 (p. 63) and the linked primary sources. The 1994 codend item is a Council recommendation; the reviewed sources do not establish NMFS implementation. The 2005 Amendment 84 action preceded exempted fishing and 2007 regulations. Council action and rule effective dates are distinct.			
The 45% / 55% allocation in the 2016 rule concerns pollock, distinct from Chinook PSC apportionments. Continuing and dated 2026 measures are described in Fishery and management history. Age-estimation and survey developments are documented in their respective report sections; they are not listed here as regulatory implementations.			
Supplied CSV; supplied Markdown; row-by-row verification and corrections. Original source records remain unchanged.			

Table 3: EBS pollock catch by reporting area and season, 1991–2026 (tons). Southeast (SE) comprises NMFS areas 520–539; Northwest (NW) comprises areas 500–517 and 519. Area 518 (Bogoslof) is excluded. A and B group records with week-ending dates before June 1 and from June 1 onward, respectively. The seasonal extract extends through October 17, 2026. Only the 1998 records included catch without an area; this catch is allocated between NW and SE in proportion to reported catch within each season. Totals include retained and discarded catch; values are rounded to whole tons.

Year	NW A	NW B	SE A	SE B	NW	SE	A	B	Total
1991	297,202	356,353	12,191	529,917	653,555	542,109	309,393	886,271	1,195,664
1992	454,947	375,612	101,597	458,143	830,559	559,741	556,544	833,755	1,390,299
1993	475,143	619,287	93,013	139,160	1,094,429	232,173	568,156	758,446	1,326,602
1994	586,105	566,470	11,026	165,751	1,152,575	176,777	597,131	732,221	1,329,352
1995	600,499	571,807	3,631	88,310	1,172,306	91,941	604,129	660,117	1,264,247
1996	549,297	537,545	1,804	104,135	1,086,843	105,939	551,101	641,680	1,192,781
1997	557,106	262,784	2,363	302,180	819,889	304,544	559,469	564,964	1,124,433
1998	515,420	443,514	1,632	141,593	958,934	143,226	517,052	585,107	1,102,159
1999	414,030	368,953	4,139	202,558	782,983	206,698	418,169	571,511	989,680
2000	439,023	400,154	19,363	274,170	839,177	293,532	458,386	674,324	1,132,710
2001	444,565	517,412	108,013	317,207	961,977	425,220	552,578	834,619	1,387,197
2002	561,061	599,272	36,693	283,749	1,160,334	320,442	597,754	883,022	1,480,776
2003	446,152	487,039	151,465	406,123	933,191	557,588	597,617	893,162	1,490,779
2004	556,469	533,538	50,661	339,883	1,090,008	390,544	607,130	873,422	1,480,552
2005	476,970	325,184	120,320	560,548	802,154	680,868	597,290	885,732	1,483,022
2006	585,507	241,700	18,789	642,034	827,207	660,824	604,297	883,734	1,488,031
2007	505,306	222,943	62,504	563,748	728,249	626,253	567,811	786,691	1,354,502
2008	358,936	123,762	44,801	463,079	482,698	507,880	403,738	586,840	990,578
2009	247,944	110,307	81,575	370,957	358,252	452,532	329,520	481,264	810,784
2010	124,670	130,443	199,011	356,061	255,114	555,072	323,681	486,505	810,186
2011	401,755	346,128	102,252	348,897	747,882	451,149	504,007	695,024	1,199,031
2012	380,373	238,497	104,926	481,418	618,870	586,344	485,299	719,915	1,205,214
2013	415,014	280,652	94,763	480,331	695,667	575,094	509,777	760,983	1,270,760
2014	469,973	388,267	50,465	388,712	858,240	439,177	520,438	776,979	1,297,417
2015	267,095	429,153	258,574	366,752	696,248	625,326	525,669	795,905	1,321,574
2016	458,044	709,028	81,531	104,078	1,167,072	185,609	539,576	813,106	1,352,681
2017	544,028	633,993	37,730	143,430	1,178,021	181,159	581,758	777,423	1,359,181
2018	598,533	450,160	3,842	326,753	1,048,692	330,595	602,375	776,913	1,379,288
2019	614,949	487,217	2,649	304,532	1,102,166	307,181	617,598	791,749	1,409,346
2020	560,438	299,978	85,717	421,102	860,416	506,820	646,156	721,080	1,367,236
2021	554,774	469,234	56,779	295,470	1,024,009	352,249	611,554	764,704	1,376,258
2022	487,216	411,573	9,018	197,610	898,790	206,629	496,235	609,184	1,105,419
2023	568,455	319,251	11,335	411,552	887,705	422,887	579,789	730,803	1,310,592
2024	530,902	347,432	55,087	377,840	878,334	432,927	585,989	725,272	1,311,261
2025	445,959	270,015	157,475	452,767	715,974	610,242	603,434	722,782	1,326,216
2026	528,440	205,467	75,601	458,034	733,908	533,635	604,041	663,501	1,267,543

Table 4: Retained and discarded EBS pollock catch by reporting area, 1991–2026 (tons), from the annual catch-accounting extract. Southeast (SE) comprises NMFS areas 520–539; Northwest (NW) comprises areas 500–517 and 519. Area 518 (Bogoslof) is excluded. Only the 1998 records included catch without an area; this catch is allocated between NW and SE in proportion to reported catch within each disposition. Discarded percentage equals discarded catch divided by total catch. Values are rounded to whole tons; 2026 estimates are preliminary.

Year	NW Discarded	SE Discarded	NW Retained	SE Retained	Total	Discarded %
1991	66,792	48,257	586,763	493,852	1,195,664	9.6
1992	71,194	57,578	759,365	502,163	1,390,299	9.3
1993	83,986	26,100	1,010,443	206,073	1,326,602	8.3
1994	88,098	16,084	1,064,476	160,693	1,329,352	7.8
1995	87,492	9,715	1,084,813	82,226	1,264,247	7.7
1996	71,368	4,838	1,015,474	101,100	1,192,781	6.4
1997	71,032	22,557	748,858	281,986	1,124,433	8.3
1998	14,291	1,581	944,511	141,777	1,102,159	1.4
1999	26,912	1,912	756,071	204,786	989,680	2.9
2000	19,678	1,942	819,499	291,591	1,132,710	1.9
2001	14,874	2,450	947,103	422,770	1,387,197	1.2
2002	19,430	1,441	1,140,904	319,002	1,480,776	1.4
2003	13,795	2,959	919,397	554,629	1,490,779	1.1
2004	20,380	2,781	1,069,628	387,763	1,480,552	1.6
2005	14,838	2,586	787,316	678,282	1,483,022	1.2
2006	11,877	3,677	815,330	657,147	1,488,031	1.0
2007	12,334	3,769	715,915	622,484	1,354,502	1.2
2008	5,968	1,643	476,730	506,237	990,578	0.8
2009	4,014	1,936	354,238	450,596	810,784	0.7
2010	2,490	1,270	252,623	553,802	810,186	0.5
2011	3,444	1,376	744,438	449,773	1,199,031	0.4
2012	4,080	1,190	614,791	585,154	1,205,214	0.4
2013	4,084	1,225	691,582	573,869	1,270,760	0.4
2014	12,556	1,786	845,684	437,391	1,297,417	1.1
2015	7,055	2,418	689,193	622,907	1,321,574	0.7
2016	8,124	1,036	1,158,948	184,574	1,352,681	0.7
2017	6,848	1,356	1,171,173	179,803	1,359,181	0.6
2018	9,170	2,005	1,039,523	328,590	1,379,288	0.8
2019	7,126	1,979	1,095,040	305,202	1,409,346	0.6
2020	9,363	2,450	851,053	504,370	1,367,236	0.9
2021	12,379	1,534	1,011,629	350,715	1,376,258	1.0
2022	9,898	3,538	888,891	203,091	1,105,419	1.2
2023	10,219	4,685	877,486	418,201	1,310,592	1.1
2024	12,935	4,131	865,399	428,796	1,311,261	1.3
2025	6,355	899	709,618	609,343	1,326,216	0.5
2026	5,815	3,182	769,440	534,068	1,312,505	0.7

Table 5: EBS pollock fishery catch-at-age estimates, 1964–2025 (million fish). Ages 15 and older are pooled as 15+. Catch numbers are calculated from assessment proportions, annual fishery weights at age and catch biomass where complete annual numbers are unavailable. Age-specific values are rounded to 0.1 million fish; All ages totals are rounded to the nearest million fish.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+	All ages
1964	8.3	34.8	54.6	63.8	31.5	88.5	39.8	8.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	330
1965	7.2	7.8	103.3	278.4	68.2	17.0	16.4	7.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	506
1966	16.9	100.2	34.2	250.2	157.8	14.4	15.7	6.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	596
1967	14.1	164.0	859.9	130.6	238.2	56.6	17.1	8.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1,489
1968	57.3	310.9	244.1	815.5	186.9	53.2	63.7	9.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1,741
1969	33.6	182.6	1051.1	722.5	248.2	5.3	27.8	17.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2,289
1970	730.1	1109.2	1743.2	823.0	159.5	51.9	25.9	14.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4,658
1971	766.6	229.0	1278.9	858.5	1272.7	123.6	81.2	22.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4,633
1972	155.0	747.6	1724.5	1381.2	578.7	227.0	86.2	31.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4,931
1973	36.3	327.2	396.1	861.1	659.1	450.3	250.5	181.5	81.4	14.7	1.8	0.5	0.1	0.5	0.0	3,261
1974	117.7	2558.9	945.7	269.0	361.3	195.1	124.9	105.3	103.3	43.5	17.7	4.6	4.6	0.8	0.7	4,853
1975	34.3	602.7	2774.6	213.3	56.7	106.6	102.1	55.6	32.7	23.0	5.7	2.3	0.4	0.5	0.0	4,010
1976	0.6	357.3	1351.8	909.8	89.7	41.4	57.3	66.5	51.7	25.5	11.8	8.0	2.1	0.0	0.0	2,973
1977	104.3	685.8	916.8	664.3	238.5	53.0	19.5	28.1	22.5	18.0	9.3	4.6	3.4	0.7	0.2	2,769
1978	29.4	455.2	752.9	638.6	368.2	99.7	28.8	17.3	19.7	15.8	11.1	4.3	1.8	0.5	0.2	2,444
1979	89.6	479.6	635.7	371.0	346.6	190.3	49.7	22.7	31.7	24.3	15.6	7.0	2.6	0.5	0.5	2,267
1980	9.0	424.0	754.9	406.7	231.2	193.5	76.8	34.5	19.9	21.9	23.3	14.6	7.1	2.3	1.1	2,221
1981	0.6	75.8	1062.5	669.3	238.2	107.9	54.2	31.1	16.9	9.8	7.9	4.8	1.6	0.7	0.3	2,282
1982	4.5	24.0	152.6	1108.4	399.3	98.1	34.0	34.1	20.4	8.6	5.1	3.0	1.8	0.7	0.3	1,895
1983	5.2	121.3	161.4	320.2	835.8	223.3	42.4	25.3	20.3	11.4	7.8	5.0	3.6	1.8	0.4	1,785
1984	1.9	41.8	80.8	392.7	448.3	596.3	122.0	32.4	22.9	14.3	6.5	2.3	2.7	1.5	1.8	1,768
1985	2.4	50.9	351.3	112.2	337.0	296.3	408.4	103.6	33.7	23.8	22.9	9.8	8.7	3.7	4.7	1,770
1986	2.8	78.5	84.2	683.0	195.4	345.0	202.5	195.5	54.5	13.9	3.0	2.4	0.2	1.1	0.0	1,862
1987	0.0	16.3	91.9	63.9	340.5	114.4	100.9	74.7	203.6	44.5	31.9	17.7	23.8	5.0	6.6	1,136
1988	0.0	9.7	410.9	381.6	228.2	492.7	203.4	95.0	35.5	87.6	16.5	9.2	3.4	5.0	5.6	1,984
1989	0.0	4.3	49.7	134.4	406.8	150.4	516.1	86.9	93.6	29.2	116.4	9.8	3.6	2.3	5.3	1,609
1990	1.4	36.1	62.2	239.8	219.3	522.1	141.2	402.6	71.8	111.4	9.9	65.6	9.3	5.1	10.1	1,908
1991	0.7	112.9	43.8	90.3	153.9	184.4	497.5	78.1	288.0	28.9	144.1	18.7	91.2	22.8	53.3	1,808
1992	1.1	84.7	675.8	130.1	79.6	108.7	133.8	253.6	102.3	147.0	58.0	46.3	13.4	43.8	34.5	1,913
1993	0.1	7.4	260.5	1146.7	103.0	66.2	66.4	56.5	86.2	21.2	32.7	12.4	13.5	6.8	16.1	1,895
1994	0.7	30.2	55.1	361.2	1059.7	175.7	53.5	19.1	13.1	20.2	9.7	9.4	7.6	4.0	8.3	1,827
1995	0.0	0.5	72.9	146.8	395.5	761.1	136.3	34.6	12.3	7.5	17.5	5.0	5.8	1.7	8.9	1,606
1996	0.0	21.7	48.1	71.8	161.0	361.9	481.7	184.6	33.6	13.4	7.9	8.8	4.3	5.8	5.3	1,410
1997	1.0	77.7	40.3	119.0	455.1	289.0	256.4	198.6	64.0	13.3	6.0	4.6	2.9	4.8	9.1	1,542
1998	0.3	42.0	84.5	70.5	153.4	702.8	199.6	131.7	110.7	27.8	6.1	5.6	2.6	2.5	4.5	1,545
1999	0.2	10.3	298.7	225.0	103.0	157.0	469.7	131.0	56.5	33.2	4.0	2.2	0.9	0.5	2.1	1,494
2000	0.0	16.1	82.4	428.5	346.5	106.7	168.4	357.7	84.9	29.7	22.0	5.2	1.4	0.6	1.0	1,651
2001	0.0	3.2	42.8	154.5	581.0	415.0	137.2	129.0	157.2	57.9	33.6	16.2	5.5	3.1	1.9	1,738

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+	All ages
2002	0.8	47.0	108.1	217.8	287.6	606.3	268.0	98.5	85.9	93.9	34.6	14.4	11.0	3.0	1.8	1,879
2003	0.0	14.5	411.8	324.1	360.4	301.5	337.6	158.6	49.4	39.3	35.7	22.9	6.6	3.6	3.2	2,069
2004	0.0	0.5	89.6	831.1	480.6	236.8	169.3	156.3	65.0	16.1	17.1	25.3	9.4	5.9	6.9	2,110
2005	0.0	4.8	52.2	392.9	863.8	484.6	159.5	68.1	66.6	30.1	10.0	9.1	3.2	2.5	3.4	2,151
2006	0.0	9.9	84.2	295.8	619.6	597.6	278.6	107.3	48.0	38.4	17.7	8.2	8.3	5.3	7.2	2,126
2007	1.7	15.7	59.2	139.2	389.4	512.0	300.8	137.1	47.6	27.5	21.9	8.9	6.5	3.4	10.8	1,682
2008	0.0	25.2	58.8	79.2	147.0	309.7	242.3	148.7	84.3	22.2	17.6	14.4	8.6	2.8	12.7	1,173
2009	0.0	1.3	175.5	200.6	82.6	114.4	124.3	104.3	66.6	40.2	23.5	7.6	7.5	2.7	8.7	960
2010	1.1	26.5	31.8	559.4	220.5	54.7	43.0	57.7	51.8	31.8	16.0	8.6	6.0	4.4	5.1	1,118
2011	0.4	10.4	193.3	115.4	809.2	284.7	63.6	37.5	38.5	41.4	25.8	12.5	1.8	4.0	4.3	1,643
2012	0.0	22.3	116.7	946.7	172.8	432.6	141.6	36.7	17.4	14.6	15.9	13.5	7.4	6.1	3.3	1,948
2013	1.8	1.0	64.0	342.4	955.9	194.4	156.5	70.0	20.7	12.8	12.8	10.9	7.8	4.8	5.7	1,861
2014	0.0	39.2	31.0	167.8	399.3	751.8	210.2	86.2	29.8	9.0	4.5	4.5	4.6	2.8	6.0	1,747
2015	0.0	15.5	632.4	196.4	228.6	384.3	510.1	88.8	42.1	17.7	2.9	2.1	3.1	2.7	1.2	2,128
2016	0.0	0.5	90.6	1390.3	159.9	174.5	174.8	224.7	34.1	13.8	8.0	0.5	1.2	0.7	1.0	2,274
2017	0.0	2.2	28.2	549.1	899.0	215.5	147.6	122.1	97.3	21.8	7.2	5.6	0.5	0.2	0.1	2,096
2018	0.0	1.3	13.8	115.0	1216.1	506.6	104.8	82.0	60.6	25.9	4.3	1.1	0.4	0.6	0.5	2,133
2019	0.7	10.9	12.3	18.4	157.6	916.8	422.5	93.2	52.1	53.0	10.1	2.9	0.8	0.0	0.0	1,751
2020	3.5	247.7	85.5	101.1	135.2	551.1	597.8	124.0	52.8	38.6	27.3	6.8	1.8	1.1	0.0	1,974
2021	0.0	111.5	1297.0	144.2	110.1	107.2	309.6	296.0	72.1	26.5	16.2	8.5	2.1	0.0	0.4	2,501
2022	0.0	62.3	174.5	1035.4	183.7	70.4	79.5	141.9	81.1	16.3	11.3	4.1	6.0	1.0	0.0	1,867
2023	0.0	25.2	86.6	263.0	1473.1	227.2	29.4	45.4	63.6	42.6	18.8	5.4	8.6	2.6	0.4	2,292
2024	0.7	2.9	104.8	131.5	362.9	1277.6	113.7	22.3	32.9	22.8	25.3	10.2	3.4	1.9	0.1	2,113
2025	0.0	28.8	50.7	100.3	129.5	526.2	852.1	92.5	29.3	26.0	8.0	19.7	0.0	0.0	6.2	1,869

Table 6: Numbers of EBS pollock measured for fishery length frequencies, by recorded sex and processing stratum, 1977–2025.

Year	A		B southeast		B northwest		Total
	Male	Female	Male	Female	Male	Female	
1977	26,411	25,923	4,301	4,511	29,075	31,219	121,440
1978	25,110	31,653	9,829	9,524	46,349	46,072	168,537
1979	59,782	62,512	3,461	3,113	62,298	61,402	252,568
1980	42,726	42,577	3,380	3,464	47,030	49,037	188,214
1981	64,718	57,936	2,401	2,147	53,161	53,570	233,933
1982	74,172	70,073	16,265	14,885	181,606	163,272	520,273
1983	94,118	90,778	16,604	16,826	193,031	174,589	585,946
1984	158,329	161,876	106,654	105,234	243,877	217,362	993,332
1985	119,384	109,230	96,684	97,841	284,850	256,091	964,080
1986	186,505	189,497	135,444	123,413	164,546	131,322	930,727
1987	373,163	399,072	14,170	21,162	24,038	22,117	853,722
1988	NA	NA	NA	NA	NA	NA	NA
1989	NA	NA	NA	NA	NA	NA	NA
1990	NA	NA	NA	NA	NA	NA	NA
1991	155,876	143,625	148,385	132,539	123,516	122,241	826,182
1992	152,566	148,024	150,829	152,003	93,073	94,701	791,196
1993	136,211	126,635	145,280	137,384	24,797	26,057	596,364
1994	138,179	146,067	154,311	148,497	26,431	26,380	639,865
1995	128,719	125,847	175,115	150,323	16,142	16,327	612,473
1996	147,992	139,905	193,493	149,814	18,101	18,288	667,593
1997	123,454	102,619	114,846	106,001	58,492	51,498	556,910
1998	135,136	109,119	205,282	174,676	31,968	39,475	695,656
1999	36,035	32,407	38,229	35,084	16,258	18,321	176,334
2000	64,430	58,030	63,746	41,027	40,839	39,105	307,177
2001	79,190	75,491	54,037	51,179	44,232	45,766	349,895
2002	71,502	69,467	65,299	64,243	37,661	39,285	347,457
2003	74,902	77,533	49,307	52,899	51,764	53,435	359,840
2004	75,208	75,811	63,146	61,957	47,261	44,220	367,603
2005	75,784	68,665	43,271	33,917	68,831	63,022	353,490
2006	72,543	63,349	35,378	27,939	77,620	67,219	344,048
2007	66,533	63,969	38,104	29,558	76,755	70,504	345,423
2008	51,303	46,296	23,467	20,462	64,126	60,678	266,332
2009	43,476	41,540	17,343	16,148	45,418	47,926	211,851
2010	41,019	39,495	20,577	19,194	40,914	40,449	201,648
2011	62,295	58,481	65,057	60,208	48,055	50,927	345,023
2012	57,946	53,557	46,942	45,024	53,243	49,968	306,680
2013	62,148	51,984	44,582	37,307	49,649	49,161	294,831
2014	58,066	55,954	51,743	46,568	46,067	46,642	305,040
2015	56,419	55,646	43,601	46,853	41,183	45,117	288,819
2016	58,915	57,478	69,654	72,973	9,015	10,264	278,299
2017	64,693	55,965	65,982	70,285	14,125	15,871	286,921
2018	64,628	57,156	49,653	56,243	32,796	35,811	296,287
2019	64,665	49,191	54,927	59,416	27,753	34,955	290,907
2020	65,609	60,018	47,791	53,161	48,459	53,985	329,023
2021	75,561	59,580	53,205	61,208	27,641	30,833	308,028
2022	114,918	93,686	86,598	94,900	33,622	36,012	459,736
2023	73,284	54,056	38,537	42,259	38,222	37,101	283,459
2024	72,356	60,933	47,300	53,442	33,511	33,693	301,397
2025	78,854	65,020	39,796	42,845	43,374	48,084	319,026

Table 7: Numbers of EBS pollock with paired fish-weight and length measurements, by recorded sex and processing stratum, 1977–2025.

Year	A		B southeast		B northwest		Total
	Male	Female	Male	Female	Male	Female	
1977	1,222	1,338	137	166	1,461	1,664	5,988
1978	1,991	2,686	409	516	2,200	2,623	10,425
1979	2,709	3,151	152	209	1,469	1,566	9,256
1980	1,849	2,156	99	144	612	681	5,541
1981	1,821	2,045	51	52	1,623	1,810	7,402
1982	2,030	2,208	181	176	2,852	3,043	10,490
1983	1,199	1,200	144	122	3,268	3,447	9,380
1984	980	1,046	117	136	1,273	1,378	4,930
1985	520	499	46	55	426	488	2,034
1986	689	794	518	501	286	286	3,074
1987	1,351	1,466	25	33	72	63	3,010
1988	NA	NA	NA	NA	NA	NA	NA
1989	NA	NA	NA	NA	NA	NA	NA
1990	NA	NA	NA	NA	NA	NA	NA
1991	2,893	2,791	1,209	1,116	2,536	2,408	12,953
1992	1,605	1,537	556	600	2,003	1,940	8,241
1993	1,278	1,205	451	437	1,412	1,459	6,242
1994	1,638	1,553	174	166	1,591	1,584	6,706
1995	1,258	1,220	232	223	1,352	1,331	5,616
1996	2,165	2,117	NA	NA	1,393	1,421	7,096
1997	629	630	552	536	674	620	3,641
1998	1,958	1,865	357	335	936	982	6,433
1999	4,813	5,337	3,767	3,546	7,182	7,954	32,599
2000	11,346	12,457	7,736	7,991	7,800	12,463	59,793
2001	14,411	14,965	9,064	8,803	10,460	10,871	68,574
2002	13,564	14,098	7,648	7,213	13,004	12,988	68,515
2003	15,535	14,857	10,272	10,031	10,111	9,437	70,243
2004	7,924	7,742	4,318	4,617	6,868	6,850	38,319
2005	7,039	7,428	6,426	6,947	4,114	5,139	37,093
2006	6,566	7,381	6,442	7,406	3,045	4,006	34,846
2007	6,640	6,695	7,081	7,798	3,202	4,305	35,721
2008	4,501	4,865	5,855	6,264	2,236	2,624	26,345
2009	4,033	4,382	4,655	4,511	1,723	1,934	21,238
2010	4,258	4,536	3,883	4,125	2,012	2,261	21,075
2011	5,845	6,388	4,954	4,647	5,929	6,456	34,219
2012	5,494	5,979	4,923	5,346	4,507	4,774	31,023
2013	5,689	6,525	4,844	4,920	3,599	4,313	29,890
2014	5,675	5,871	4,785	4,652	4,753	5,180	30,916
2015	5,310	5,323	4,648	4,194	4,365	4,064	27,904
2016	5,312	5,725	1,077	909	6,872	6,635	26,530
2017	5,238	6,047	1,586	1,343	6,575	6,254	27,043
2018	5,583	6,174	3,430	3,172	5,506	4,850	28,715
2019	4,513	6,086	3,594	2,953	5,809	5,499	28,454
2020	6,116	6,846	5,325	4,815	5,376	4,900	33,378
2021	5,852	7,368	6,247	5,468	2,886	2,698	30,519
2022	4,890	5,847	1,705	1,566	4,899	4,364	23,271
2023	10,676	14,487	6,171	6,173	5,911	5,436	48,854
2024	7,253	6,276	5,027	5,679	3,298	3,421	30,958
2025	7,391	6,333	4,257	4,597	4,379	4,922	31,927

Table 8: Numbers of EBS pollock contributing age estimates, by recorded sex and processing stratum, 1977–2025. The 2024–2025 entries are released FT-NIRS age predictions.

Year	A		B southeast		B northwest		Total
	Male	Female	Male	Female	Male	Female	
1977	1,229	1,344	137	166	1,415	1,613	5,904
1978	1,992	2,686	407	514	2,188	2,611	10,398
1979	2,647	3,088	152	209	1,464	1,561	9,121
1980	1,854	2,158	93	138	606	675	5,524
1981	1,819	2,042	51	52	1,620	1,807	7,391
1982	2,030	2,210	181	176	2,865	3,062	10,524
1983	1,200	1,200	144	122	3,249	3,420	9,335
1984	980	1,046	117	136	1,272	1,379	4,930
1985	520	499	46	55	426	488	2,034
1986	689	794	518	501	286	286	3,074
1987	1,351	1,466	25	33	72	63	3,010
1988	NA	NA	NA	NA	NA	NA	NA
1989	NA	NA	NA	NA	NA	NA	NA
1990	NA	NA	NA	NA	NA	NA	NA
1991	439	431	367	349	263	289	2,138
1992	399	396	178	180	391	375	1,919
1993	476	445	122	124	496	507	2,170
1994	201	200	142	132	574	571	1,820
1995	313	299	131	123	420	439	1,725
1996	465	479	NA	NA	436	443	1,823
1997	437	434	343	341	313	286	2,154
1998	663	595	237	222	311	316	2,344
1999	506	541	298	308	748	750	3,151
2000	629	667	293	254	596	847	3,286
2001	563	603	205	178	697	736	2,982
2002	672	663	247	202	890	839	3,513
2003	653	588	274	262	701	671	3,149
2004	547	561	221	245	698	600	2,872
2005	599	617	420	422	490	614	3,162
2006	528	609	507	568	367	459	3,038
2007	627	642	552	568	485	594	3,468
2008	513	497	538	650	342	368	2,908
2009	404	484	440	432	240	299	2,299
2010	545	624	413	466	418	505	2,971
2011	581	808	404	396	582	660	3,431
2012	517	571	485	579	480	533	3,165
2013	666	703	525	568	401	518	3,381
2014	609	629	413	407	475	553	3,086
2015	653	642	511	493	508	513	3,320
2016	488	599	157	125	929	969	3,267
2017	604	778	179	163	777	753	3,254
2018	569	662	366	358	621	591	3,167
2019	552	778	387	332	558	531	3,138
2020	757	899	405	420	450	408	3,339
2021	760	910	588	542	270	256	3,326
2022	609	777	209	212	616	563	2,986
2023	452	633	350	313	399	342	2,489
2024	700	616	493	534	291	350	2,985
2025	697	562	401	433	439	456	2,992

Table 9: Pollock sampling effort from the NMFS bottom-trawl survey, 1982–2026 (counts). Historical values are as reported; recent haul counts use the survey records and recent ages are released EBS assignments, including FT-NIRS ages.

Year	Hauls	Lengths	Aged / age assigned
1982	329	40,001	1,611
1983	354	78,033	1,931
1984	355	40,530	1,806
1985	434	48,642	1,913
1986	354	41,101	1,344
1987	356	40,144	1,607
1988	373	40,408	1,173
1989	373	38,926	1,227
1990	371	34,814	1,257
1991	371	43,406	1,083
1992	356	34,024	1,263
1993	375	43,278	1,385
1994	375	38,901	1,141
1995	376	25,673	1,156
1996	375	40,789	1,387
1997	376	35,536	1,193
1998	375	37,673	1,261
1999	373	32,532	1,385
2000	372	41,762	1,545
2001	375	47,335	1,641
2002	375	43,361	1,695
2003	376	46,480	1,638
2004	375	44,102	1,660
2005	373	35,976	1,676
2006	376	39,211	1,573
2007	376	29,679	1,484
2008	375	24,635	1,251
2009	376	24,819	1,342
2010	517	23,142	1,385
2011	376	36,227	1,734
2012	376	35,782	1,785
2013	376	35,908	1,847
2014	376	43,042	2,099
2015	376	54,241	2,320
2016	376	50,857	1,766
2017	519	47,873	1,623
2018	376	48,673	1,486
2019	520	42,382	1,519
2020	NA	NA	NA
2021	520	53,545	2,050
2022	520	36,687	1,980
2023	492	38,618	1,677
2024	350	36,137	1,526
2025	487	30,862	1,619
2026	350	31,296	1,582
NA denotes an unavailable count; the survey was absent in 2020. The 2025 haul count includes 350 EBS and 137 NBS records; the 2026 count includes 350 EBS records. Historical coverage varies among years.			
The 2025 and 2026 length counts use the EBS production summaries. Recent length and age counts cover EBS; haul counts include NBS where stated.			
Recent age assignments are distinct from traditional tester-pair counts; historical age counts retain their published definitions.			

Table 10: Bottom-trawl survey (BTS) pollock biomass for the combined eastern and northern Bering Sea, 1982–2026, with assessment standard errors (SE) and coefficients of variation (CV).

Year	Biomass (million t)	SE (thousand t)	CV (%)
1982	4.064	732	18.0%
1983	10.457	1,174	11.2%
1984	6.777	977	14.4%
1985	7.794	1,008	12.9%
1986	7.459	1,078	14.5%
1987	8.131	1,399	17.2%
1988	12.000	1,896	15.8%
1989	10.554	1,435	13.6%
1990	11.449	2,352	20.5%
1991	7.472	1,427	19.1%
1992	6.961	1,121	16.1%
1993	8.275	1,060	12.8%
1994	7.385	1,159	15.7%
1995	7.631	2,335	30.6%
1996	4.179	551	13.2%
1997	4.842	890	18.4%
1998	3.665	565	15.4%
1999	5.732	1,020	17.8%
2000	7.778	1,350	17.4%
2001	6.271	705	11.2%
2002	7.117	971	13.6%
2003	12.148	4,671	38.5%
2004	6.063	739	12.2%
2005	6.867	952	13.9%
2006	4.181	513	12.3%
2007	6.511	992	15.2%
2008	4.271	633	14.8%
2009	2.860	482	16.8%
2010	5.174	839	16.2%
2011	4.524	520	11.5%
2012	5.063	681	13.5%
2013	6.739	790	11.7%
2014	11.838	1,266	10.7%
2015	11.145	982	8.8%
2016	8.825	1,006	11.4%
2017	8.781	709	8.1%
2018	5.119	813	15.9%
2019	9.026	967	10.7%
2021	4.706	338	7.2%
2022	6.348	649	10.2%
2023	4.754	429	9.0%
2024	8.354	1,196	14.3%
2025	5.801	792	13.7%
2026	9.034	956	10.6%

CV is the annual standard error divided by biomass. These are the uncertainties used in the assessment, which also accounts for correlations among years. The survey was absent in 2020.

Table 11: Combined eastern and northern Bering Sea model-based pollock abundance at age used for the assessment, 1982–2026 (million fish). Ages 15 and older are pooled as 15+.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+	Total
1982	1220.6	2780.1	3221.8	4394.2	1481.2	196.2	126.3	60.7	37.7	23.7	14.8	8.8	3.3	1.4	1.4	13572.2
1983	4548.2	1080.4	2880.1	4538.0	9238.2	2681.1	458.1	210.7	97.6	87.7	62.5	22.9	8.3	7.7	3.6	25925.1
1984	494.0	404.5	803.7	2218.2	2460.1	5226.9	929.5	201.8	92.6	31.1	21.3	8.0	4.9	6.1	3.1	12905.9
1985	5028.8	787.5	3999.5	1409.8	4629.4	2808.1	1856.2	353.5	86.3	73.3	25.6	8.0	8.6	0.7	0.7	21075.9
1986	3061.3	918.2	793.9	3276.1	1686.3	2542.1	2103.4	1790.2	534.1	83.7	37.4	16.5	1.5	4.6	1.6	16850.8
1987	495.7	706.6	1227.4	956.9	5641.8	1477.3	1356.5	522.5	1628.6	256.8	79.2	30.0	5.9	2.9	2.6	14390.8
1988	1588.1	769.9	2287.4	4488.9	1959.7	6093.4	1659.9	1198.1	666.8	1611.9	155.7	91.5	20.6	24.6	14.8	22631.4
1989	944.5	576.1	979.3	2967.0	6523.0	1245.0	4451.9	628.4	703.5	263.0	825.0	141.4	126.8	59.2	95.4	20529.5
1990	2285.0	272.5	107.0	1089.6	2485.0	8094.0	1492.5	3554.8	337.0	587.7	89.3	832.7	71.8	52.6	73.2	21424.7
1991	3250.4	947.9	368.5	134.1	952.9	836.1	2526.4	881.9	1708.2	430.0	576.7	117.6	346.1	52.7	60.4	13189.9
1992	1617.8	418.5	2737.7	428.0	520.3	836.6	731.6	1010.7	438.8	810.1	281.6	355.1	153.0	121.0	97.3	10558.3
1993	2776.9	444.9	1162.3	5158.9	1074.2	819.5	409.8	538.4	681.7	416.6	359.3	261.2	209.2	115.1	140.3	14568.1
1994	1722.8	695.5	525.9	2019.7	5779.3	950.8	229.2	178.9	196.0	361.9	221.7	308.3	118.8	117.1	198.7	13624.6
1995	1631.4	204.8	456.2	1978.6	2724.3	4580.8	1714.5	388.3	229.3	147.8	272.0	112.0	206.2	84.3	134.9	14865.5
1996	2000.9	437.8	198.4	408.0	1127.1	1494.9	1288.5	423.6	102.1	111.4	74.7	142.5	48.6	87.4	128.8	8074.7
1997	2338.7	398.5	216.5	265.8	3005.0	1397.2	817.3	1038.9	178.6	90.3	66.3	71.4	124.0	40.3	152.6	10201.3
1998	902.8	708.6	371.2	262.0	548.3	3044.7	782.6	476.6	364.3	94.0	41.5	14.1	32.9	35.1	86.7	7765.4
1999	1000.6	1041.4	1090.9	1116.1	634.3	1180.5	2883.0	751.1	362.3	337.0	126.5	54.0	22.4	33.6	123.5	10757.3
2000	1053.0	376.4	703.0	2187.0	2206.2	1225.8	970.7	2977.1	1110.5	573.6	246.2	158.1	50.3	23.0	103.9	13964.7
2001	1985.9	1224.0	712.7	713.1	1687.0	1656.8	688.4	335.0	987.1	704.8	269.8	212.6	82.1	30.2	87.2	11376.7
2002	1319.8	393.9	794.7	1211.9	1379.5	1789.1	892.4	430.4	575.4	1076.3	538.2	249.8	149.1	48.0	57.8	10906.2
2003	565.2	170.1	1149.5	1885.8	2127.9	1800.1	2248.6	1153.8	477.1	626.8	1226.1	562.9	236.1	98.8	109.5	14438.2
2004	401.3	266.2	191.1	1614.3	1597.9	1188.0	665.2	702.3	346.8	219.4	220.1	406.2	181.4	48.1	40.8	8089.1
2005	397.3	153.0	251.1	1214.6	3364.3	2117.6	1068.1	485.3	369.1	269.1	75.0	156.8	261.1	108.4	123.4	10414.1
2006	822.5	76.0	113.3	525.4	1405.2	1644.7	972.2	442.0	245.2	204.2	98.3	61.0	87.9	119.3	123.5	6940.6
2007	2730.3	65.0	224.4	719.4	2148.5	2411.1	1628.2	1147.3	449.4	195.6	180.2	154.0	71.7	88.1	178.4	12391.4
2008	501.3	122.5	141.0	269.9	777.1	1573.6	1152.4	734.2	437.9	158.4	129.9	97.2	44.2	24.6	159.1	6323.3
2009	863.4	210.1	435.1	497.8	325.1	488.6	690.0	521.6	371.6	171.0	107.8	35.1	35.2	18.3	80.1	4850.6
2010	480.0	136.6	315.5	4277.0	1722.7	479.2	379.3	389.4	391.1	268.4	227.6	82.8	49.9	29.5	68.1	9296.8
2011	1331.7	136.3	359.0	431.6	2165.5	1048.1	294.4	160.1	244.1	240.6	205.5	158.0	69.4	33.1	98.2	6975.5
2012	1341.7	211.1	485.1	4164.0	951.6	1431.6	456.1	171.2	115.0	158.5	127.7	110.9	96.5	34.9	67.6	9923.3
2013	1257.5	134.9	273.7	1127.2	5422.6	1228.5	750.4	262.1	88.5	79.4	105.0	79.8	77.2	44.9	63.6	10995.3
2014	2531.9	477.4	279.9	497.3	2312.4	7741.2	3820.1	797.9	435.3	157.6	62.1	89.1	93.0	46.9	141.4	19483.5
2015	1460.1	947.2	2835.6	767.6	1528.6	2738.5	5150.1	1522.0	385.3	192.1	27.0	25.2	49.9	30.5	70.4	17730.0
2016	782.9	507.3	744.5	4155.6	1720.4	1196.9	1767.8	2703.3	547.2	224.0	79.7	17.8	18.7	6.8	14.1	14486.9
2017	789.3	312.7	526.1	2619.5	3497.6	1602.5	1236.2	1161.9	1488.5	425.9	163.7	55.1	8.4	4.8	16.2	13908.4
2018	1179.6	474.1	251.5	376.6	2531.9	1556.8	541.1	393.9	405.0	333.0	93.2	14.6	1.9	0.3	5.8	8159.1
2019	2685.3	712.4	584.7	619.6	1763.6	6308.4	2520.5	597.1	431.0	254.9	133.9	60.8	24.1	6.9	2.7	16705.9
2020	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
2021	1027.6	827.5	1953.4	765.3	703.6	508.1	1291.4	1479.8	391.6	152.5	127.5	82.6	18.2	12.9	4.4	9346.2

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+	Total
2022	821.4	348.4	595.5	4025.4	1888.9	733.5	815.5	1143.0	855.9	271.4	116.2	96.7	37.6	21.3	2.2	11773.0
2023	1152.8	388.7	341.0	630.2	2710.5	804.7	436.7	424.2	702.6	503.9	126.4	68.4	38.1	11.7	10.6	8350.8
2024	1683.9	957.6	893.4	610.1	1944.1	6404.9	1426.1	447.7	424.6	389.1	275.1	107.7	33.5	14.6	5.4	15617.7
2025	1534.2	344.2	318.5	504.0	491.9	1366.7	3303.4	827.7	327.6	208.0	299.8	162.0	81.4	50.4	26.5	9846.2
2026	3466.8	608.7	1637.4	948.4	1034.1	1390.7	2587.6	2390.8	738.5	253.8	442.7	369.9	114.1	11.4	69.8	16064.7
Mean	1615.3	549.5	921.3	1692.5	2315.4	2225.9	1444.8	863.9	495.2	321.1	205.4	142.5	80.1	41.2	69.3	12983.4

The full current revised series is shown; historical values differ from the earlier published VAST series. Northern survey coverage varies among years. These survey-expansion estimates are distinct from model population numbers at age.

The 2025 and 2026 FT-NIRS products use their year-specific traditional-age keys; the 2026 key retains 307 pairs after excluding 22 flagged pairs. Exact traditional ages 1–18 are converted before 15+ pooling, conserving total abundance. Empirical-key uncertainty is excluded. NA in 2020 denotes no survey observation; the mean uses the 44 observed-year estimates. Values are rounded to 0.1 million fish.

Table 12: Annual BTS-derived population weight-at-age inputs used in Model 26.0, 1982–2026 (kg per fish). Ages 15 and older are pooled as 15+.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
1982	0.027	0.076	0.180	0.369	0.550	0.862	1.151	1.326	1.463	1.648	1.790	1.778	2.142	1.770	2.633
1983	0.017	0.091	0.179	0.341	0.491	0.691	0.988	1.304	1.494	1.539	1.634	1.600	1.807	1.555	1.961
1984	0.017	0.062	0.187	0.293	0.465	0.605	0.955	1.375	1.520	1.710	1.957	1.829	1.880	1.675	2.091
1985	0.017	0.070	0.183	0.333	0.485	0.753	0.941	1.238	1.667	1.648	1.779	2.062	2.135	1.817	2.505
1986	0.014	0.066	0.137	0.356	0.433	0.657	0.829	1.025	1.270	1.635	1.714	1.952	2.051	2.212	* 2.191
1987	0.028	0.105	0.239	0.291	0.424	0.555	0.705	0.861	0.993	1.224	1.506	1.539	1.931	2.215	2.339
1988	0.019	0.079	0.246	0.331	0.444	0.527	0.661	0.883	0.910	1.101	1.366	1.390	1.872	1.169	1.715
1989	0.022	0.085	0.166	0.293	0.384	0.509	0.710	0.802	1.088	1.072	1.201	1.268	1.303	1.290	1.400
1990	0.019	0.101	0.145	0.337	0.441	0.536	0.633	0.741	0.916	1.134	1.256	1.313	1.393	1.498	1.692
1991	0.025	0.109	0.202	0.339	0.549	0.629	0.714	0.809	0.986	1.108	1.364	1.400	1.552	1.824	1.930
1992	0.018	0.096	0.270	0.381	0.510	0.575	0.774	0.916	0.969	1.083	1.262	1.455	1.601	1.662	1.630
1993	0.015	0.087	0.271	0.470	0.524	0.646	0.757	0.895	1.101	1.112	1.318	1.406	1.561	1.673	1.786
1994	0.019	0.083	0.209	0.388	0.559	0.675	0.719	1.053	1.239	1.199	1.322	1.413	1.525	1.680	1.758
1995	0.016	0.057	0.139	0.339	0.485	0.657	0.693	0.912	1.079	1.328	1.314	1.373	1.501	1.481	1.747
1996	0.021	0.077	0.137	0.297	0.500	0.632	0.797	0.903	1.043	1.176	1.386	1.511	1.680	1.682	1.789
1997	0.018	0.078	0.193	0.299	0.380	0.623	0.754	0.899	1.122	1.185	1.319	1.529	1.583	1.645	1.764
1998	0.018	0.070	0.185	0.323	0.460	0.524	0.801	0.895	1.012	1.082	1.286	1.349	1.485	1.886	1.886
1999	0.016	0.078	0.219	0.347	0.407	0.561	0.700	0.897	1.087	1.152	1.292	1.323	1.754	1.756	1.982
2000	0.014	0.079	0.236	0.400	0.477	0.562	0.704	0.786	0.946	1.164	1.364	1.476	1.720	1.752	2.001
2001	0.018	0.078	0.172	0.359	0.530	0.669	0.797	0.881	0.996	1.118	1.337	1.499	1.589	1.794	1.880
2002	0.017	0.094	0.244	0.389	0.521	0.674	0.760	0.935	1.033	1.121	1.127	1.401	1.563	1.741	1.934
2003	0.027	0.113	0.279	0.461	0.621	0.735	0.815	0.968	1.045	1.201	1.250	1.371	1.548	1.550	1.744
2004	0.023	0.107	0.275	0.421	0.567	0.689	0.819	0.928	1.041	1.152	1.210	1.304	1.456	1.748	1.746
2005	0.022	0.088	0.217	0.395	0.510	0.630	0.809	0.942	1.020	1.108	1.361	1.314	1.299	1.480	1.599
2006	0.011	0.087	0.170	0.421	0.598	0.696	0.815	0.894	1.042	1.164	1.275	1.390	1.467	1.434	1.489
2007	0.014	0.102	0.263	0.497	0.637	0.802	0.907	1.039	1.146	1.406	1.372	1.411	1.723	1.636	1.689
2008	0.014	0.056	0.221	0.455	0.585	0.717	0.846	0.997	1.058	1.237	1.537	1.509	1.726	1.850	1.894
2009	0.012	0.074	0.224	0.499	0.675	0.791	0.999	1.135	1.174	1.481	1.568	1.759	2.049	2.277	2.360
2010	0.020	0.077	0.249	0.510	0.675	0.875	1.094	1.188	1.245	1.352	1.453	1.651	1.876	2.213	2.443

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
2011	0.017	0.085	0.209	0.499	0.663	0.773	0.938	1.118	1.247	1.296	1.455	1.530	1.638	1.727	2.015
2012	0.015	0.075	0.195	0.420	0.606	0.784	0.933	1.137	1.333	1.401	1.432	1.529	1.558	1.845	2.015
2013	0.018	0.079	0.213	0.501	0.582	0.744	0.993	1.166	1.354	1.409	1.542	1.582	1.710	1.860	1.923
2014	0.019	0.109	0.257	0.438	0.571	0.716	0.791	1.106	1.231	1.423	1.676	1.606	1.672	1.847	1.903
2015	0.020	0.090	0.327	0.445	0.588	0.686	0.784	0.905	1.166	1.229	1.496	1.550	1.432	1.492	1.750
2016	0.023	0.080	0.279	0.531	0.595	0.697	0.779	0.828	0.904	1.004	1.137	1.554	1.351	1.536	1.421
2017	0.021	0.092	0.231	0.472	0.625	0.673	0.755	0.814	0.892	0.919	1.057	0.993	1.289	1.123	1.485
2018	0.023	0.092	0.215	0.421	0.591	0.684	0.777	0.816	0.896	0.933	1.036	0.944	1.410	* 1.720	1.540
2019	0.022	0.105	0.286	0.487	0.634	0.731	0.822	0.925	0.937	1.035	0.965	1.066	1.038	1.063	* 1.910
2020	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
2021	0.022	0.101	0.245	0.442	0.585	0.713	0.803	0.857	0.977	1.118	1.040	1.117	1.298	0.999	1.361
2022	0.018	0.112	0.270	0.447	0.590	0.697	0.771	0.873	0.969	1.034	1.254	1.251	1.324	1.205	1.812
2023	0.015	0.106	0.283	0.416	0.536	0.657	0.812	0.955	0.968	1.013	1.179	1.200	1.414	1.759	1.487
2024	0.017	0.090	0.309	0.454	0.560	0.607	0.693	0.923	0.997	1.020	1.126	1.137	1.259	1.406	1.754
2025	0.020	0.079	0.226	0.526	0.587	0.676	0.734	0.812	0.903	1.087	1.066	1.150	1.258	1.399	1.547
2026	0.020	0.066	0.397	0.544	0.640	0.725	0.809	0.874	0.949	1.027	1.048	1.192	1.416	1.449	1.360
Mean	0.019	0.086	0.227	0.409	0.542	0.673	0.815	0.967	1.101	1.218	1.351	1.431	1.587	1.645	1.838

* The selected value is retained. Its current treatment has not been reconciled with an older preparation record identifying these historical cells for filling; the technical supplement preserves that provenance qualification.

The table reproduces the selected schedules, including historical revisions, rather than labeling every cell as an unmodified survey measurement. NA in 2020 marks absence of a survey observation; the model retains its separate annual schedule. The mean uses the 44 displayed annual schedules. Values are rounded to 0.001 kg.

Table 13: Sampling effort and biological collections from eastern Bering Sea acoustic-trawl surveys, 1979–2026 (counts). E and W denote collections east and west of 170°W within the US EEZ; US is the reported US subtotal and RU is the Russian-region count. The 2024 US aged-fish count is 2,260 (McCarthy et al. 2026, p. 15).

Year	Hauls				Length records				Otoliths				Number aged			
	E	W	US	RU	E	W	US	RU	E	W	US	RU	E	W	US	RU
1979	NA	NA	25	NA	NA	NA	7722	NA	NA	NA	0	NA	NA	NA	2610	NA
1982	13	31	48	NA	1725	6689	8687	NA	840	2324	3164	NA	783	1958	2741	NA
1985	NA	NA	73	NA	NA	NA	19872	NA	NA	NA	2739	NA	NA	NA	2739	NA
1988	NA	NA	25	NA	NA	NA	6619	NA	NA	NA	1471	NA	NA	NA	1471	NA
1991	NA	NA	62	NA	NA	NA	16343	NA	NA	NA	2062	NA	NA	NA	1663	NA
1994	25	51	76	19	4553	21011	25564	8930	1560	3694	4966	1270	612	932	1770	455
1996	15	42	57	NA	3551	13273	16824	NA	669	1280	1949	NA	815	1111	1926	NA
1997	25	61	86	NA	6493	23043	29536	NA	966	2669	3635	NA	936	1349	2285	NA
1999	41	77	118	NA	13841	28521	42362	NA	1945	3001	4946	NA	946	1500	2446	NA
2000	29	95	124	NA	7721	36008	43729	NA	850	2609	3459	NA	850	1403	2253	NA
2002	47	79	126	NA	14601	25633	40234	NA	1424	1883	3307	NA	1000	1200	2200	NA
2004	33	57	90	15	8896	18262	27158	5893	1167	2002	3169	461	798	1192	2351	461
2006	27	56	83	NA	4939	19326	24265	NA	822	1871	2693	NA	822	1870	2692	NA
2007	23	46	69	4	5492	14863	20355	1407	871	1961	2832	319	823	1737	2560	315
2008	9	53	62	6	2394	15354	17748	1754	341	1698	2039	177	338	1381	1719	176
2009	13	33	46	3	1576	9257	10833	282	308	1210	1518	54	306	1205	1511	54
2010	11	48	59	9	2432	20263	22695	3502	653	1868	2521	381	652	1598	2250	379
2012	17	60	77	14	4422	23929	28351	5620	650	2045	2695	418	646	1483	2129	416
2014	52	87	139	3	28857	8645	37502	747	1739	849	2588	72	845	1735	2580	72
2016	37	71	108	NA	10912	24134	35046	NA	880	1514	2394	NA	876	1513	2388	NA
2018	36	64	100	NA	11031	21173	32204	NA	1105	1750	2855	NA	1100	1743	2843	NA
2020	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
2022	26	36	62	NA	4954	11994	16948	NA	671	1691	2362	NA	668	1688	2356	NA
2024	14	41	55	NA	4535	14472	19007	NA	725	1561	2286	NA	NA	NA	2260	NA
2026	36	62	99	NA	11916	23031	35128	NA	1051	1571	2642	NA	NA	NA	NA	NA

Table 14: Midwater pollock biomass by area from summer ATS surveys of the U.S. EEZ Bering Sea shelf, 1994–2026 (million tons; near surface to 0.5 m above the seafloor). The core shelf excludes the Northern Extension. Assessment CVs average 20%, with 45% assigned to 2000; SE equals biomass × CV. The 2020 survey used uncrewed sailing vessels; 2026 estimates are preliminary.

Year	Date	Area (nmi^2)	Biomass (million t)				Rel. error	Assessment CV	Assessment SE (kt)
			SCA	E170–SCA	W170	0.5 m total			
1994	9 Jul - 7 Sept	78,251	0.382	0.652	2.594	3.628	0.040	17.0%	618.4
1996	20 Jul - 30 Aug	93,810	0.275	0.492	2.185	2.952	0.032	13.6%	402.5
1997	17 Jul - 5 Sept	102,770	0.272	0.703	2.470	3.446	0.029	12.4%	425.8
1999	12 Jun - 29 Jul	103,670	0.383	0.757	3.060	4.200	0.044	18.8%	787.5
2000	12 Jun - 1 Aug	106,140	0.458	0.715	2.441	3.615	0.028	45.0%	1,626.6
2002	9 Jun - 21 Jul	99,526	0.762	0.938	2.601	4.301	0.027	11.5%	494.9
2004	4 Jun -29 Jul	99,659	0.550	0.918	2.528	3.996	0.031	13.2%	527.9
2006	3 Jun -25 Jul	89,550	0.147	0.340	1.387	1.874	0.033	14.1%	263.6
2007	2 Jun -30 Jul	92,944	0.136	0.245	1.902	2.283	0.038	16.2%	369.7
2008	2 Jun -31 Jul	95,374	0.122	0.087	1.196	1.404	0.056	23.9%	335.2
2009	9 Jun -7 Aug	91,414	0.156	0.058	1.117	1.331	0.069	29.4%	391.5
2010	5 Jun -7 Aug	92,849	0.098	0.193	2.345	2.636	0.054	23.0%	606.6
2012	7 Jun -10 Aug	96,852	0.196	0.319	1.764	2.279	0.034	14.5%	330.3
2014	12 Jun -13 Aug	94,361	0.571	1.458	2.714	4.743	0.034	14.5%	687.2
2016	12 Jun -17 Aug	100,674	0.542	1.268	3.028	4.838	0.019	8.1%	391.7
2018	12 Jun -22 Aug	92,283	0.234	0.510	1.753	2.497	0.039	16.6%	415.1
2020	4 Jul - 20 Aug	102,320	0.398	0.531	2.688	3.617	0.096	40.9%	1,479.8
2022	1 Jun -5 Aug	103,942	0.538	0.823	2.473	3.834	0.068	29.0%	1,111.1
2024	5 Jun - 22 Jul	100,365	0.497	0.593	1.785	2.875	0.056	23.9%	686.1
2026	28 May - 12 Aug	95,091	0.239	0.706	2.738	3.682	0.034	14.5%	533.6

Table 15: Biomass and abundance summed across the 2026 survey intervals, including the below-3 m analysis and northern extension. The three core-shelf regions sum to the historical EBS estimate; the Northern Extension is additional.

Region	EDSU records	Biomass (t)	Abundance (million fish)
SCA	1,079	238,999.6	720.50
East of 170	3,232	705,748.5	1,396.55
West of 170	5,137	2,737,718.7	10,545.58
EBS Northern Extension	1,476	78,311.1	104.29

Table 16: ATS estimates of EBS pollock abundance at age, 1979–2026 (million fish). Ages 10 and older are pooled as 10+. The 2026 estimates use the BTS age–length key applied to ATS length distributions and remain provisional; ages from the 2026 ATS samples are expected in 2027. Mean and median summarize surveys with age compositions during 1994–2026; 1979–1991 provides historical context.

Year	Age										Age 2+	Total at age
	1	2	3	4	5	6	7	8	9	10+		
1979	69,110	41,132	3,884	413	534	128	30	4	28	161	46,314	115,424
1982	108	3,401	4,108	7,637	1,790	283	141	178	90	177	17,805	17,913
1985	2,076	929	8,149	898	2,186	1,510	1,127	130	21	15	14,965	17,041
1988	11	1,112	3,586	3,864	739	1,882	403	151	130	414	12,280	12,292
1991	639	5,942	967	215	224	133	120	39	37	53	7,730	8,369
1994	1,140	4,969	1,424	1,819	2,252	389	109	96	56	221	11,335	12,475
1996	1,800	567	552	2,741	915	634	585	142	39	165	6,338	8,139
1997	13,227	2,881	440	536	2,330	546	313	290	75	220	7,633	20,860
1999	607	1,780	3,717	1,810	652	398	1,548	526	180	249	10,859	11,466
2000	460	1,322	1,230	2,588	1,012	327	308	950	278	252	8,266	8,726
2002	796	4,944	3,385	1,295	661	935	538	140	162	493	12,554	13,351
2004	83	313	1,217	3,123	1,634	567	288	283	121	265	7,811	7,894
2006	525	217	291	654	783	659	390	145	75	171	3,386	3,910
2007	5,775	1,041	345	478	794	729	407	241	98	135	4,267	10,042
2008	71	2,915	1,047	166	161	288	235	136	102	120	5,169	5,240
2009	5,197	816	1,734	281	77	94	129	111	77	114	3,433	8,630
2010	2,568	6,404	984	2,295	446	73	33	37	38	91	10,400	12,968
2012	177	1,989	1,693	2,710	280	367	113	36	25	103	7,315	7,492
2014	4,751	8,655	969	1,161	1,119	1,770	740	170	79	99	14,762	19,513
2016	174	1,038	4,496	4,476	715	348	392	420	96	64	12,046	12,220
2018	450	517	249	621	2,268	944	198	112	107	104	5,120	5,570
2022	142	332	975	6,578	819	211	133	239	166	79	9,533	9,674
2024	5,075	1,146	681	379	820	2,485	450	110	86	127	6,284	11,359
2026	2,026	3,729	3,958	375	319	403	714	656	198	285	10,637	12,663
Mean 1994–2026	2,371	2,399	1,547	1,794	950	640	401	255	108	177	8,271	10,642
Median 1994–2026	796	1,322	1,047	1,295	794	403	313	145	96	135	7,811	10,042

Table 17: Recent core-area ATS pollock biomass and abundance estimates. The 2026 estimates are preliminary.

Recent core-area ATS estimates		
Survey year	Biomass (million t)	Abundance (billion fish)
2022	3.8	9.67
2024	2.87	11.4
2026 (preliminary)	3.68	12.7

Table 18: Provisional core-area 2026 ATS abundance by age, in million fish, restricted to supported lengths. FT-NIRS source age 15 retains its released label; the corresponding TMA row pools target ages 15–18. The upper-length tail remains unallocated.

Age (FT-NIRS / TMA)	FT-NIRS abundance	TMA abundance
1	2,194.690	2,025.867
2	5,340.635	3,729.246
3	1,978.834	3,958.120
4	597.222	374.975
5	338.742	318.742
6	299.224	403.011
7	619.494	713.961
8	786.844	656.327
9	343.438	197.732
10	99.819	63.788
11	45.953	106.936
12	12.907	80.216
13	2.903	19.546
14	0.836	1.844
15 / 15+	0.980	12.210

Table 19: Provisional 2026 ATS biomass allocation using the same BTS FT-NIRS length and conditional TMA keys as abundance. Biomass columns are tonnes. Unsupported lengths remain unallocated; core EBS excludes the Northern Extension.

Region	Input (t)	FT-NIRS (t)	TMA (t)	Unallocated (t)	Unallocated (%)
All supplied regions	3,760,777.888	3,759,915.746	3,759,915.746	862.142	0.022925
Core EBS	3,682,466.780	3,682,141.718	3,682,141.718	325.062	0.008827
EBS Northern Extension	78,311.108	77,774.028	77,774.028	537.080	0.685829
East of 170	705,748.481	705,694.837	705,694.837	53.644	0.007601
SCA	238,999.623	238,999.623	238,999.623	0.000	0.000000
West of 170	2,737,718.675	2,737,447.258	2,737,447.258	271.417	0.009914

Table 20: Provisional core-area 2026 ATS TMA biomass, abundance and mean weight at age over supported lengths. The 15+ row sums exact ages 15–18 before dividing biomass by abundance. Unsupported upper-length biomass and fish are excluded from these age-specific estimates and retained separately.

TMA age	Biomass (t)	Abundance (million fish)	Mean weight (kg/fish)
1	89,550.280	2,025.867	0.0442
2	525,587.105	3,729.246	0.1409
3	949,648.466	3,958.120	0.2399
4	182,528.304	374.975	0.4868
5	185,878.529	318.742	0.5832
6	263,257.538	403.011	0.6532
7	525,152.651	713.961	0.7355
8	518,643.709	656.327	0.7902
9	168,006.366	197.732	0.8497
10	57,043.013	63.788	0.8943
11	96,454.798	106.936	0.9020
12	81,491.146	80.216	1.0159
13	22,896.782	19.546	1.1714
14	2,181.579	1.844	1.1832
15+	13,821.450	12.210	1.1320

Table 21: Assumed age-composition likelihood sample sizes used in the completed full-data Model 23.0 fit, 1964–2026. Fishery inputs for 2023–2025 are the applied rounded values 443, 578 and 530; other values are rounded for display.

Year	Fishery	BTS	ATS
1964–1977	10		
1978	39		
1979	39		
1980	39		
1981	39		
1982	39	153	
1983	39	134	
1984	39	61	
1985	39	76	
1986	39	153	
1987	39	60	
1988	39	81	
1989	39	55	
1990	39	48	
1991	260	79	

Year	Fishery	BTS	ATS
1992	228	86	
1993	344	92	
1994	286	91	43
1995	257	103	
1996	189	227	32
1997	319	77	49
1998	355	64	
1999	476	97	67
2000	482	88	70
2001	328	178	
2002	487	187	72
2003	441	130	
2004	391	195	51
2005	494	188	
2006	506	321	47
2007	500	184	39
2008	524	238	35
2009	420	148	26
2010	549	107	34
2011	728	172	
2012	607	108	44
2013	753	64	
2014	606	117	79
2015	821	104	
2016	704	124	61
2017	607	241	
2018	668	156	50
2019	701	120	
2020	565		1
2021	818	150	
2022	641	181	52
2023	443	135	
2024	578	63	79
2025	530	111	
2026		131	94

Year	Fishery	BTS	ATS
Mean	503	129	51

Table 22: Revised mean-one AVO display index for 17 observations during 2009–2026, excluding 2020, including newly processed 2011 and 2013 values. Sampling relative errors, standard errors and scaled source 95% confidence intervals are retained. The blank 2020 row preserves calendar-year coverage.

Year	Index (million m^2)	Relative error (%)	SE (million m^2)	Relative index	SE (relative)	95% CI (relative)
2009	0.6951	12.22%	0.0849	0.2990	0.0365	0.2274–0.3707
2010	1.9218	6.56%	0.1261	0.8268	0.0542	0.7205–0.9331
2011	1.5679	4.51%	0.0707	0.6745	0.0304	0.6149–0.7342
2012	1.5206	5.32%	0.0810	0.6542	0.0348	0.5859–0.7225
2013	2.2347	4.64%	0.1037	0.9614	0.0446	0.8740–1.0488
2014	3.0769	4.11%	0.1263	1.3238	0.0543	1.2172–1.4303
2015	3.5927	2.46%	0.0884	1.5457	0.0380	1.4711–1.6202
2016	2.8319	2.91%	0.0825	1.2184	0.0355	1.1488–1.2879
2017	2.2630	3.05%	0.0691	0.9736	0.0297	0.9153–1.0319
2018	2.0841	2.66%	0.0555	0.8966	0.0239	0.8498–0.9434
2019	2.8292	4.89%	0.1383	1.2172	0.0595	1.1006–1.3338
2020						
2021	2.4101	4.79%	0.1154	1.0369	0.0496	0.9396–1.1342
2022	2.9031	3.41%	0.0990	1.2490	0.0426	1.1655–1.3325
2023	2.4810	2.94%	0.0730	1.0674	0.0314	1.0058–1.1289
2024	2.0070	2.72%	0.0546	0.8635	0.0235	0.8174–0.9095
2025	2.4954	5.26%	0.1313	1.0736	0.0565	0.9628–1.1843
2026	2.5997	4.55%	0.1182	1.1185	0.0508	1.0188–1.2181

Source: NOAA AFSC MACE AVO workbook, 2009–2026. The 2020 observation is unavailable, so its data cells are blank.

SE = index × CV. Relative index and SE share the mean-1 scaling; relative error is unchanged. Source 95% intervals are scaled by the same constant.

Table 23: Counts of AVO sampling intervals during 2009–2026, including positive intervals and recorded zeros; these counts describe survey coverage. There are no AVO observations in 2020.

Year	Survey intervals	Positive intervals	Recorded zeros
2009	5,557	2,430	56.3%
2010	5,699	2,584	54.7%
2011	7,829	5,171	34.0%
2012	6,740	4,750	29.5%
2013	8,021	4,725	41.1%
2014	10,032	8,315	17.1%
2015	10,524	9,700	7.8%
2016	8,603	7,912	8.0%
2017	8,798	6,833	22.3%
2018	8,880	8,044	9.4%
2019	8,836	7,301	17.4%
2020	—	—	—
2021	6,939	6,038	13.0%
2022	9,350	8,502	9.1%
2023	8,740	7,435	14.9%
2024	12,229	10,406	14.9%
2025	7,320	6,952	5.0%
2026	8,699	7,462	14.2%

Table 24: ATS biomass (million tons) and AVO backscatter (million m²), 2006–2026. AVO indices and sampling relative errors for 2009–2026 match Table 22. Assessment CVs from the saved Model 26.0 inputs average 23% across 20 observations; SE equals index × CV. The 2020 USV survey provides ATS biomass; AVO has no observation that year.

Year	ATS biomass (million t)	AVO index (million m^2)	Sampling RE	Assessment CV	Assessment SE (million m^2)
2006	1.8741	1.7415	0.0510	27.64%	0.4814
2007	2.2827	2.0019	0.0865	53.90%	1.0790
2008	1.4045	0.9915	0.0643	19.84%	0.1968
2009	1.3314	0.6951	0.1222	56.89%	0.3955
2010	2.6357	1.9218	0.0656	30.55%	0.5872
2011	—	1.5679	0.0451	21.01%	0.3294
2012	2.2794	1.5206	0.0532	24.80%	0.3771
2013	—	2.2347	0.0464	21.60%	0.4827
2014	4.7428	3.0769	0.0411	19.12%	0.5882
2015	—	3.5927	0.0246	11.46%	0.4118
2016	4.8377	2.8319	0.0291	13.57%	0.3843
2017	—	2.2630	0.0305	14.22%	0.3218
2018	2.4973	2.0841	0.0266	12.40%	0.2585
2019	—	2.8292	0.0489	22.77%	0.6441
2020	3.6170	—	—	—	—
2021	—	2.4101	0.0479	22.29%	0.5372
2022	3.8340	2.9031	0.0341	15.88%	0.4610
2023	—	2.4810	0.0294	13.70%	0.3400
2024	2.8750	2.0070	0.0272	12.67%	0.2543
2025	—	2.4954	0.0526	24.51%	0.6117
2026	3.6825	2.5997	0.0455	21.17%	0.5504
Mean	2.9149	2.2125	0.0486	23.00%	0.4646

Table 25: Annual fishery mean weight at age (kg) used in the full-data Model 23.0 main input through 2026, with the completed run's annual weight-model forecasts for 2027–2028. Observational inputs extend through 2025; the 2026 row is predicted. Age 15 is a 15+ group. The fixed early schedule and mean/CV definitions are described below.

Year / basis	Age (years)														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
1964–1990	0.007	0.170	0.303	0.447	0.589	0.722	0.840	0.942	1.029	1.102	1.163	1.212	1.253	1.286	1.312
1991	0.007	0.150	0.285	0.483	0.604	0.730	0.844	0.882	1.020	1.123	1.135	1.237	1.236	1.296	1.247
1992	0.007	0.179	0.397	0.465	0.651	0.714	0.819	0.986	1.030	1.200	1.237	1.269	1.193	1.357	1.431
1993	0.007	0.331	0.495	0.612	0.652	0.775	0.934	1.062	1.198	1.240	1.423	1.540	1.576	1.609	1.508
1994	0.007	0.233	0.400	0.652	0.732	0.746	0.727	1.070	1.380	1.325	1.335	1.409	1.397	1.278	1.370
1995	0.007	0.153	0.386	0.505	0.729	0.843	0.847	0.970	1.232	1.296	1.401	1.402	1.392	1.095	1.306
1996	0.007	0.293	0.336	0.445	0.684	0.797	0.948	0.956	1.025	1.100	1.418	1.489	1.521	1.702	1.602
1997	0.007	0.187	0.327	0.477	0.559	0.748	0.889	1.074	1.095	1.236	1.287	1.400	1.561	1.363	1.338
1998	0.007	0.191	0.369	0.589	0.618	0.622	0.780	1.040	1.169	1.276	1.316	1.428	1.448	1.437	1.528
1999	0.007	0.188	0.404	0.507	0.643	0.702	0.729	0.894	1.038	1.253	1.224	1.422	0.995	0.616	1.239
2000	0.007	0.218	0.353	0.526	0.630	0.732	0.780	0.807	0.968	1.015	1.253	1.286	1.108	1.084	1.359
2001	0.006	0.227	0.329	0.505	0.668	0.786	0.964	0.986	1.061	1.133	1.320	1.411	1.568	1.472	1.495
2002	0.007	0.231	0.385	0.510	0.667	0.799	0.911	1.026	1.113	1.102	1.284	1.442	1.579	1.290	1.568
2003	0.006	0.276	0.489	0.549	0.652	0.769	0.863	0.953	1.086	1.202	1.212	1.194	1.374	1.355	1.709
2004	0.007	0.135	0.408	0.584	0.641	0.760	0.888	0.924	1.036	1.176	1.127	1.167	1.309	1.254	1.185
2005	0.007	0.283	0.351	0.508	0.641	0.742	0.880	0.960	1.062	1.074	1.216	1.268	1.217	1.076	1.342
2006	0.007	0.174	0.306	0.448	0.606	0.755	0.858	0.959	1.060	1.117	1.190	1.218	1.280	1.384	1.417
2007	0.007	0.155	0.349	0.507	0.642	0.783	0.961	1.100	1.192	1.266	1.327	1.488	1.444	1.729	1.512
2008	0.007	0.208	0.328	0.519	0.653	0.774	0.900	1.054	1.117	1.289	1.452	1.528	1.560	1.874	1.645
2009	0.007	0.136	0.340	0.525	0.705	0.879	0.999	1.130	1.398	1.479	1.558	1.576	1.807	2.026	2.222
2010	0.050	0.175	0.381	0.490	0.668	0.909	1.114	1.277	1.374	1.586	1.679	1.923	1.948	2.077	2.271
2011	0.031	0.205	0.290	0.508	0.666	0.809	0.971	1.224	1.342	1.513	1.582	1.623	2.080	1.707	2.242
2012	0.029	0.142	0.271	0.409	0.643	0.824	0.974	1.170	1.303	1.509	1.599	1.637	1.680	2.031	2.062
2013	0.095	0.144	0.290	0.442	0.564	0.781	1.130	1.281	1.440	1.685	1.827	1.786	1.934	2.159	2.182
2014	0.014	0.193	0.319	0.454	0.617	0.751	0.894	1.156	1.307	1.386	1.669	1.773	1.704	1.623	2.215
2015	0.025	0.181	0.404	0.462	0.571	0.690	0.786	0.887	1.141	1.195	1.315	1.671	1.389	1.559	2.600
2016	0.025	0.181	0.409	0.531	0.557	0.646	0.732	0.800	0.941	1.043	1.178	0.788	0.911	1.684	1.429
2017	0.025	0.191	0.408	0.499	0.650	0.694	0.752	0.827	0.894	0.911	1.028	0.961	0.312	0.701	0.688
2018	0.025	0.186	0.377	0.467	0.573	0.734	0.809	0.853	0.906	1.039	0.936	1.110	0.568	0.435	1.130
2019	0.025	0.186	0.422	0.565	0.643	0.759	0.878	0.962	1.007	1.065	1.035	1.182	0.754	0.698	0.769
2020	0.025	0.186	0.387	0.522	0.632	0.716	0.799	0.955	1.006	1.040	1.189	1.072	1.208	0.961	0.769
2021	0.025	0.186	0.393	0.480	0.574	0.690	0.757	0.841	1.011	1.130	1.160	1.269	1.215	0.936	0.408
2022	0.025	0.186	0.440	0.506	0.574	0.724	0.837	0.883	0.984	0.972	1.271	1.044	1.235	0.910	0.770
2023	0.025	0.186	0.494	0.483	0.524	0.602	0.812	0.923	0.957	1.041	1.095	1.294	1.331	1.218	1.131

Year / basis	Age (years)														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
2024	0.025	0.186	0.435	0.481	0.560	0.617	0.696	0.825	1.016	1.017	1.078	1.251	1.308	1.328	1.559
2025	0.025	0.186	0.314	0.556	0.621	0.693	0.730	0.855	0.966	1.140	1.194	1.082	1.314	1.343	1.268
2026 predicted	0.025	0.186	0.643	0.500	0.696	0.754	0.831	0.881	0.968	1.082	1.190	1.238	1.309	1.418	1.440
2027 forecast			0.370	0.757	0.619	0.818	0.876	0.952	0.998	1.081	1.190	1.292	1.334	1.399	1.501
2028 forecast			0.370	0.484	0.877	0.742	0.940	0.996	1.069	1.111	1.188	1.291	1.387	1.423	1.482
Mean	0.018	0.196	0.373	0.508	0.629	0.745	0.863	0.987	1.111	1.205	1.301	1.361	1.356	1.362	1.472
CV (%)	97.9%	22.7%	15.6%	10.0%	7.8%	9.1%	12.1%	13.2%	13.6%	15.1%	15.5%	18.1%	26.8%	31.0%	33.7%

1964–1990 uses the identical fixed schedule in every year. Annual rows through 2025 retain the actual main-data inputs: observed means at ages 3–15+ where available, with model-predicted fills for 2025 ages 13 and 14. The 2026 ages 3–15+ are the saved annual WtAgeRe predictions copied into that main input.

Ages 1–2 in 2025 and 2026 retain the 2024 input values. Forecast rows 2027–2028 have blank ages 1–2 because that annual model estimates ages 3–15+ only. Blank cells indicate unavailable estimates. Future rows use the completed Model 23.0 annual fishery weight forecasts, which differ from the A-season spawning-weight schedule. Detailed source records are in the technical appendix.

Mean and CV use the 35 main-input schedules from 1991–2025, including the identified fills and younger-age values retained from the baseline. The fixed early row and all 2026–2028 predictions/forecasts are excluded. CV = sample SD / mean, expressed as percent; it describes variation in the weight schedules among years.

Table 26: Effects of cumulative input updates within Model 23.0, expressed as 2026 estimates. Biomass uses the same reference population weight schedule in every fit, and fishing mortality is the maximum realized rate across ages. Each row includes preceding updates. ATS and AVO contrasts combine new observations, historical revisions and the stated uncertainty calibrations; the fits estimate their combined effect.

Input updates within Model 23.0				
Cumulative input update	Biomass (million t)	Spawning biomass (million t)	Age-1 fish (billion)	Fishing mortality (per year)
Catch + endpoint	9.030	2.623	18.525	0.348
+ Fishery weights 2024	8.856	2.542	18.541	0.348
+ Fishery weights 2026	8.857	2.533	18.541	0.348
+ Fishery age data	9.323	2.746	18.523	0.341
+ BTS	9.966	2.834	24.345	0.335
+ ATS	11.043	2.761	31.731	0.345
+ AVO	11.227	2.808	31.948	0.341

Table 27: Deterministic 2026 population estimates with the selected catch inputs. Model 26.1.2 is the Early-late fishery split comparison; its marginal sampling estimator is excluded.

2026 estimates after the catch revision				
Model	Biomass (million t)	Spawning biomass (million t)	Age-1 fish (billion)	Maximum F (per year)
Model 23.2	12.267	3.031	37.532	0.304
Model 23.0	11.227	2.808	31.948	0.341
Model 26.0	12.162	3.057	36.347	0.299
Model 26.1.0	12.153	3.053	36.334	0.300
Model 26.1.1	12.198	3.064	36.826	0.316
Model 26.1 decline penalty 0	12.351	3.110	37.608	0.338
Model 26.1.2 Early-late fishery split	12.235	3.065	36.518	0.315
ADMB biomass uses the same population weights as Model 26.0. Model 26.1.2 is a deterministic native joint mode conditional on fixed Foreign and Modern SDs; its sampling-based marginal estimator was excluded by author instruction. This table presents point estimates; scientific model adoption remains separate.				

Table 28: Model 26.0 estimated population quantities in 2026. Ranges are approximate 95% confidence intervals under the selected model and fixed biological schedules. Fishing mortality is the largest realized rate across ages in 2026.

Estimated 2026 population					
Quantity	Estimate	SE	95% lower	95% upper	Units
Total biomass	12.162	1.511	9.534	15.515	Million tons
Spawning biomass	3.057	0.320	2.489	3.754	Million tons
Age-1 recruitment	36.347	19.395	12.772	103.440	Billion fish
Fishing mortality: most exposed age	0.299	0.067	0.194	0.463	Per year

Table 29: Model 26.0 annual estimates and approximate 95% confidence intervals, 1964–2026. Each cell gives the estimate followed by its 95% interval. Biomass and spawning biomass are million tons, recruitment is billion age-1 fish and fishing mortality is the maximum realized rate across ages per year. Recruitment is labeled by the calendar year fish enter age 1.

Annual population estimates and 95% intervals				
Year	Biomass (million t)	Spawning biomass (million t)	Age-1 fish (billion)	Fishing mortality (per year)
1964	4.827 (3.351–6.953)	1.348 (0.909–1.998)	5.727 (2.316–14.165)	0.077 (0.043–0.135)
1965	5.201 (3.701–7.308)	1.376 (0.939–2.016)	23.494 (13.757–40.121)	0.085 (0.049–0.146)
1966	5.420 (3.893–7.545)	1.398 (0.946–2.067)	15.880 (8.083–31.199)	0.090 (0.053–0.153)
1967	6.159 (4.520–8.392)	1.508 (1.027–2.215)	27.213 (16.078–46.059)	0.161 (0.091–0.286)
1968	6.803 (5.019–9.223)	1.674 (1.131–2.476)	22.441 (12.605–39.953)	0.178 (0.107–0.297)
1969	7.387 (5.544–9.843)	1.895 (1.292–2.781)	26.281 (15.584–44.321)	0.179 (0.104–0.308)
1970	8.087 (6.171–10.599)	2.006 (1.388–2.899)	23.326 (13.486–40.346)	0.265 (0.126–0.556)

Annual population estimates and 95% intervals				
Year	Biomass (million t)	Spawning biomass (million t)	Age-1 fish (billion)	Fishing mortality (per year)
1971	8.012 (6.127–10.478)	2.117 (1.476–3.036)	13.629 (6.794–27.342)	0.386 (0.178–0.839)
1972	7.273 (5.512–9.596)	1.978 (1.384–2.825)	10.896 (5.305–22.380)	0.426 (0.196–0.929)
1973	6.521 (4.990–8.522)	1.697 (1.159–2.484)	26.743 (18.276–39.133)	0.471 (0.221–1.005)
1974	5.485 (4.193–7.175)	1.293 (0.834–2.007)	19.281 (13.001–28.596)	0.499 (0.303–0.821)
1975	4.711 (3.663–6.059)	1.200 (0.784–1.835)	16.451 (11.522–23.489)	0.467 (0.327–0.668)
1976	4.304 (3.468–5.342)	1.121 (0.794–1.581)	12.944 (9.258–18.097)	0.423 (0.278–0.643)
1977	4.036 (3.351–4.860)	1.305 (0.984–1.732)	14.023 (10.506–18.716)	0.377 (0.171–0.830)
1978	4.311 (3.700–5.023)	1.158 (0.892–1.504)	26.513 (21.903–32.094)	0.436 (0.197–0.968)
1979	5.530 (4.931–6.200)	1.068 (0.839–1.359)	62.439 (54.916–70.993)	0.462 (0.220–0.969)
1980	6.294 (5.684–6.969)	1.268 (1.044–1.541)	29.662 (24.950–35.264)	0.424 (0.263–0.684)
1981	7.831 (7.139–8.589)	1.934 (1.693–2.210)	34.243 (29.110–40.280)	0.308 (0.201–0.472)
1982	9.443 (8.615–10.352)	2.680 (2.387–3.008)	19.180 (15.520–23.704)	0.201 (0.139–0.291)
1983	10.311 (9.464–11.233)	3.594 (3.219–4.013)	52.001 (46.089–58.672)	0.174 (0.113–0.267)
1984	10.272 (9.441–11.176)	3.732 (3.350–4.158)	14.229 (11.444–17.692)	0.172 (0.130–0.227)
1985	12.273 (11.351–13.271)	4.262 (3.876–4.686)	32.543 (28.408–37.280)	0.181 (0.132–0.249)
1986	11.664 (10.861–12.526)	4.145 (3.805–4.516)	12.743 (10.478–15.497)	0.168 (0.115–0.246)
1987	10.839 (10.212–11.504)	4.157 (3.857–4.480)	7.323 (5.855–9.158)	0.129 (0.086–0.194)
1988	9.780 (9.224–10.370)	3.909 (3.657–4.179)	5.712 (4.552–7.168)	0.181 (0.122–0.268)
1989	8.195 (7.706–8.714)	3.382 (3.165–3.613)	11.908 (10.240–13.848)	0.209 (0.159–0.275)
1990	7.731 (7.305–8.183)	2.870 (2.674–3.080)	50.462 (46.960–54.226)	0.292 (0.214–0.396)
1991	8.525 (8.086–8.988)	2.320 (2.144–2.511)	27.363 (24.774–30.224)	0.406 (0.289–0.569)
1992	9.146 (8.688–9.628)	2.202 (2.049–2.366)	23.766 (21.304–26.512)	0.484 (0.330–0.711)
1993	10.389 (9.875–10.931)	2.878 (2.704–3.063)	46.285 (42.750–50.113)	0.304 (0.211–0.438)
1994	10.238 (9.733–10.769)	3.341 (3.143–3.551)	15.931 (14.075–18.031)	0.233 (0.157–0.346)
1995	8.867 (8.408–9.350)	3.572 (3.367–3.790)	11.171 (9.736–12.818)	0.296 (0.252–0.348)
1996	9.339 (8.859–9.845)	3.509 (3.307–3.723)	22.654 (20.657–24.844)	0.376 (0.308–0.458)
1997	8.875 (8.422–9.352)	3.412 (3.210–3.626)	30.307 (28.043–32.753)	0.339 (0.229–0.504)
1998	8.748 (8.302–9.217)	2.901 (2.726–3.087)	14.951 (13.463–16.604)	0.275 (0.199–0.380)
1999	8.937 (8.494–9.403)	3.020 (2.847–3.204)	16.029 (14.509–17.708)	0.213 (0.180–0.251)
2000	9.070 (8.635–9.528)	2.903 (2.741–3.075)	24.585 (22.686–26.642)	0.272 (0.224–0.330)
2001	9.067 (8.643–9.512)	2.939 (2.776–3.113)	34.415 (32.234–36.744)	0.291 (0.239–0.354)
2002	9.528 (9.109–9.966)	2.728 (2.577–2.889)	22.938 (21.267–24.742)	0.374 (0.318–0.440)
2003	10.763 (10.308–11.238)	2.748 (2.605–2.899)	14.341 (13.115–15.681)	0.415 (0.338–0.510)
2004	9.630 (9.214–10.065)	3.032 (2.882–3.189)	6.525 (5.751–7.404)	0.361 (0.279–0.467)
2005	7.764 (7.409–8.136)	2.663 (2.529–2.804)	4.624 (3.988–5.362)	0.335 (0.282–0.399)
2006	6.805 (6.471–7.156)	2.431 (2.302–2.567)	11.539 (10.424–12.774)	0.406 (0.348–0.474)
2007	6.328 (6.007–6.666)	1.994 (1.876–2.119)	24.986 (23.081–27.048)	0.432 (0.370–0.505)
2008	4.826 (4.568–5.099)	1.528 (1.427–1.636)	13.553 (12.109–15.170)	0.443 (0.372–0.528)
2009	5.899 (5.593–6.221)	1.617 (1.512–1.730)	49.521 (46.316–52.948)	0.381 (0.256–0.569)
2010	7.701 (7.321–8.101)	1.654 (1.549–1.767)	23.135 (21.102–25.363)	0.317 (0.237–0.425)
2011	8.485 (8.068–8.924)	2.029 (1.907–2.158)	14.211 (12.693–15.910)	0.433 (0.356–0.526)
2012	8.811 (8.359–9.287)	2.475 (2.331–2.628)	12.310 (10.960–13.827)	0.429 (0.266–0.691)
2013	9.756 (9.251–10.287)	2.824 (2.655–3.003)	43.285 (40.546–46.208)	0.428 (0.253–0.723)
2014	10.381 (9.869–10.919)	2.782 (2.606–2.969)	49.480 (46.207–52.984)	0.403 (0.226–0.719)
2015	11.568 (11.013–12.151)	2.597 (2.436–2.768)	21.258 (19.132–23.620)	0.449 (0.245–0.823)
2016	12.325 (11.722–12.960)	2.855 (2.690–3.030)	8.625 (7.265–10.239)	0.312 (0.244–0.400)

Annual population estimates and 95% intervals				
Year	Biomass (million t)	Spawning biomass (million t)	Age-1 fish (billion)	Fishing mortality (per year)
2017	11.425 (10.826–12.056)	3.300 (3.110–3.502)	7.890 (6.545–9.511)	0.295 (0.211–0.414)
2018	9.899 (9.330–10.502)	3.152 (2.957–3.361)	16.823 (14.472–19.557)	0.276 (0.232–0.329)
2019	10.741 (10.071–11.457)	3.067 (2.851–3.299)	87.831 (78.998–97.651)	0.438 (0.360–0.532)
2020	11.795 (10.940–12.717)	2.364 (2.171–2.575)	27.572 (23.257–32.688)	0.419 (0.353–0.497)
2021	11.818 (10.823–12.905)	2.619 (2.394–2.866)	12.958 (9.881–16.992)	0.416 (0.332–0.523)
2022	13.137 (11.866–14.545)	3.742 (3.383–4.139)	12.683 (9.030–17.815)	0.291 (0.220–0.385)
2023	11.901 (10.613–13.345)	3.779 (3.369–4.238)	11.616 (7.811–17.274)	0.232 (0.155–0.348)
2024	10.767 (9.397–12.337)	3.526 (3.076–4.041)	39.254 (25.030–61.560)	0.243 (0.201–0.294)
2025	10.134 (8.563–11.993)	3.228 (2.732–3.814)	27.976 (16.327–47.937)	0.293 (0.225–0.380)
2026	12.162 (9.534–15.515)	3.057 (2.489–3.754)	36.347 (12.772–103.440)	0.299 (0.194–0.463)

Table 30: Final 2026 population summaries from the regularized Model 26.0 posterior. All 5,000 saved draws contribute. Intervals are central 95% credible intervals. Priors, parameter support and supplied biology remain fixed. Production-fit estimates and projections retain their separate roles.

Posterior population uncertainty				
Quantity and units	Mean	Median	2.5%	97.5%
SSB 2026 (million t)	2.9645	2.9465	2.4497	3.5844
Biomass 2026 (million t)	11.5860	11.4616	9.2151	14.7192
Recruitment 2026 (billion fish)	36.4680	31.1518	12.2880	91.0616

Table 31: Model 26.0 start-of-year numbers at age, 1964–2026, in billion fish. Age 15+ includes all older fish. Values are rounded to 0.001 billion fish; full precision is retained in the supporting tables.

	Numbers at age (billion fish)														
Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
1964	5.727	4.350	2.577	0.877	0.514	0.633	0.390	0.247	0.198	0.173	0.149	0.127	0.106	0.088	0.159
1965	23.494	2.328	2.763	1.872	0.612	0.356	0.434	0.269	0.171	0.137	0.120	0.104	0.089	0.074	0.172
1966	15.880	9.549	1.476	1.987	1.274	0.417	0.243	0.297	0.184	0.117	0.094	0.083	0.072	0.061	0.170
1967	27.213	6.454	6.040	1.053	1.345	0.864	0.283	0.165	0.202	0.125	0.080	0.064	0.057	0.049	0.158
1968	22.441	11.052	4.038	4.077	0.664	0.850	0.548	0.180	0.105	0.129	0.080	0.051	0.041	0.036	0.132
1969	26.281	9.109	6.877	2.693	2.528	0.414	0.533	0.345	0.113	0.066	0.081	0.050	0.032	0.026	0.106
1970	23.326	10.660	5.638	4.445	1.667	1.574	0.259	0.334	0.216	0.071	0.041	0.051	0.031	0.020	0.082
1971	13.629	9.439	6.459	3.435	2.654	0.984	0.930	0.152	0.193	0.124	0.041	0.024	0.029	0.018	0.058
1972	10.896	5.500	5.584	3.709	1.888	1.418	0.529	0.500	0.079	0.100	0.064	0.021	0.012	0.015	0.038
1973	26.743	4.398	3.158	2.995	1.895	0.965	0.726	0.271	0.249	0.039	0.049	0.031	0.010	0.006	0.026
1974	19.281	10.796	2.452	1.583	1.418	0.896	0.457	0.345	0.126	0.116	0.018	0.023	0.014	0.005	0.014
1975	16.451	7.788	5.813	1.103	0.714	0.641	0.407	0.208	0.155	0.057	0.052	0.008	0.010	0.007	0.009
1976	12.944	6.658	4.425	2.700	0.526	0.347	0.314	0.200	0.102	0.077	0.028	0.026	0.004	0.005	0.007
1977	14.023	5.244	3.863	2.220	1.311	0.261	0.174	0.159	0.102	0.052	0.039	0.014	0.013	0.002	0.006
1978	26.513	5.685	3.073	2.134	1.152	0.671	0.135	0.090	0.083	0.053	0.027	0.020	0.007	0.007	0.004
1979	62.439	10.753	3.356	1.697	1.095	0.568	0.333	0.067	0.045	0.041	0.025	0.013	0.009	0.003	0.005
1980	29.662	25.334	6.512	1.953	0.894	0.526	0.271	0.160	0.032	0.021	0.019	0.012	0.006	0.004	0.004
1981	34.243	12.042	15.722	4.120	1.061	0.440	0.255	0.132	0.078	0.016	0.010	0.009	0.006	0.003	0.004
1982	19.180	13.910	7.571	10.735	2.474	0.577	0.240	0.140	0.073	0.043	0.009	0.006	0.005	0.003	0.004
1983	52.001	7.794	8.800	5.406	7.005	1.499	0.350	0.146	0.085	0.044	0.026	0.005	0.003	0.003	0.004
1984	14.229	21.135	4.935	6.345	3.678	4.454	0.933	0.219	0.091	0.053	0.028	0.017	0.003	0.002	0.005
1985	32.543	5.783	13.399	3.558	4.364	2.346	2.778	0.583	0.137	0.057	0.033	0.017	0.010	0.002	0.004
1986	12.743	13.228	3.666	9.626	2.464	2.876	1.450	1.721	0.363	0.086	0.036	0.021	0.011	0.007	0.004
1987	7.323	5.180	8.390	2.635	6.620	1.641	1.821	0.908	1.088	0.229	0.054	0.023	0.013	0.007	0.007
1988	5.712	2.977	3.291	6.063	1.859	4.529	1.089	1.207	0.591	0.709	0.150	0.036	0.015	0.009	0.009
1989	11.908	2.322	1.889	2.308	4.180	1.215	2.892	0.674	0.753	0.369	0.446	0.095	0.023	0.009	0.011
1990	50.462	4.841	1.474	1.348	1.573	2.721	0.768	1.739	0.412	0.467	0.229	0.278	0.059	0.014	0.013
1991	27.363	20.513	3.069	1.051	0.883	0.939	1.582	0.426	0.962	0.232	0.268	0.128	0.158	0.034	0.015
1992	23.766	11.123	13.004	2.212	0.709	0.545	0.550	0.835	0.237	0.517	0.130	0.133	0.065	0.080	0.025
1993	46.285	9.660	7.040	9.058	1.484	0.440	0.299	0.266	0.386	0.109	0.241	0.059	0.061	0.030	0.048
1994	15.931	18.816	6.141	5.016	5.788	0.975	0.266	0.164	0.145	0.212	0.061	0.136	0.033	0.034	0.043
1995	11.171	6.476	11.970	4.488	3.422	3.424	0.588	0.156	0.097	0.087	0.127	0.037	0.082	0.020	0.046
1996	22.654	4.542	4.119	8.785	3.202	2.200	1.886	0.337	0.090	0.057	0.052	0.077	0.022	0.050	0.040
1997	30.307	9.210	2.882	3.010	6.386	2.202	1.297	0.960	0.172	0.049	0.032	0.030	0.045	0.013	0.053

Numbers at age (billion fish)															
Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+
1998	14.951	12.321	5.830	2.098	2.142	4.358	1.377	0.729	0.524	0.091	0.027	0.018	0.017	0.026	0.037
1999	16.029	6.078	7.822	4.233	1.487	1.457	2.659	0.843	0.421	0.295	0.052	0.016	0.011	0.010	0.037
2000	24.585	6.517	3.865	5.567	2.949	1.011	0.946	1.592	0.514	0.259	0.182	0.032	0.010	0.007	0.029
2001	34.415	9.995	4.144	2.792	3.765	1.891	0.648	0.560	0.899	0.307	0.159	0.112	0.020	0.006	0.022
2002	22.938	13.991	6.359	3.017	1.923	2.277	1.047	0.364	0.318	0.524	0.184	0.095	0.068	0.012	0.017
2003	14.341	9.325	8.893	4.605	2.049	1.174	1.160	0.540	0.192	0.175	0.301	0.108	0.057	0.040	0.017
2004	6.525	5.830	5.931	6.270	3.117	1.213	0.615	0.567	0.271	0.102	0.098	0.176	0.063	0.033	0.034
2005	4.624	2.653	3.710	4.296	3.941	1.884	0.684	0.318	0.295	0.147	0.058	0.058	0.105	0.038	0.040
2006	11.539	1.880	1.688	2.690	2.846	2.166	0.998	0.370	0.176	0.165	0.085	0.034	0.035	0.064	0.047
2007	24.986	4.691	1.195	1.193	1.747	1.593	1.069	0.507	0.191	0.093	0.089	0.047	0.020	0.020	0.063
2008	13.553	10.158	2.982	0.845	0.768	0.978	0.766	0.526	0.262	0.101	0.050	0.048	0.026	0.011	0.045
2009	49.521	5.510	6.459	2.150	0.555	0.441	0.465	0.369	0.260	0.132	0.053	0.026	0.025	0.013	0.029
2010	23.135	20.133	3.507	4.643	1.417	0.335	0.226	0.238	0.190	0.135	0.068	0.027	0.013	0.013	0.022
2011	14.211	9.405	12.817	2.564	2.970	0.863	0.197	0.127	0.128	0.103	0.074	0.037	0.014	0.007	0.019
2012	12.310	5.777	5.985	9.326	1.780	1.497	0.415	0.096	0.062	0.063	0.050	0.036	0.019	0.007	0.013
2013	43.285	5.005	3.675	4.338	6.114	1.155	0.723	0.201	0.047	0.031	0.031	0.024	0.018	0.009	0.010
2014	49.480	17.597	3.185	2.667	2.916	3.703	0.691	0.385	0.098	0.023	0.015	0.015	0.012	0.008	0.009
2015	21.258	20.116	11.197	2.315	1.823	1.817	2.084	0.356	0.201	0.051	0.011	0.007	0.008	0.006	0.009
2016	8.625	8.642	12.808	7.777	1.536	1.146	1.015	1.080	0.185	0.105	0.025	0.005	0.004	0.004	0.007
2017	7.890	3.507	5.505	9.386	4.577	0.972	0.694	0.596	0.586	0.103	0.059	0.014	0.003	0.002	0.006
2018	16.823	3.208	2.233	4.045	6.465	2.573	0.548	0.391	0.334	0.323	0.058	0.033	0.008	0.002	0.004
2019	87.831	6.840	2.042	1.641	2.896	3.781	1.446	0.314	0.224	0.194	0.191	0.034	0.020	0.005	0.004
2020	27.572	35.707	4.349	1.497	1.185	1.992	2.008	0.691	0.164	0.123	0.109	0.113	0.020	0.012	0.005
2021	12.958	11.209	22.578	3.140	1.035	0.768	1.044	0.979	0.355	0.082	0.062	0.056	0.060	0.011	0.009
2022	12.683	5.268	7.061	15.644	2.187	0.668	0.471	0.534	0.478	0.181	0.042	0.033	0.030	0.032	0.011
2023	11.616	5.156	3.322	5.072	10.705	1.455	0.427	0.283	0.296	0.276	0.107	0.024	0.019	0.018	0.025
2024	39.254	4.722	3.268	2.384	3.526	6.678	0.891	0.267	0.171	0.174	0.164	0.063	0.014	0.011	0.025
2025	27.976	15.959	3.002	2.342	1.659	2.308	3.879	0.546	0.166	0.104	0.106	0.100	0.038	0.009	0.022
2026	36.347	11.374	10.149	2.172	1.644	1.112	1.276	2.157	0.320	0.098	0.062	0.064	0.060	0.023	0.019

Table 32: Model 26.0 start-of-2026 numbers at age and uncertainty, in billion fish. Ranges are approximate 95% confidence intervals, accounting for covariance in the fitted model. Age 15+ includes all older fish.

2026 numbers at age and uncertainty					
Age	Estimate	SE	CV	95% lower	95% upper
1	36.347	19.395	53.4%	12.772	103.440
2	11.374	3.125	27.5%	6.638	19.489
3	10.149	2.335	23.0%	6.466	15.932
4	2.172	0.449	20.7%	1.448	3.259
5	1.644	0.309	18.8%	1.138	2.376
6	1.112	0.184	16.5%	0.804	1.537
7	1.276	0.176	13.8%	0.973	1.672
8	2.157	0.259	12.0%	1.704	2.730
9	0.320	0.045	13.9%	0.243	0.420
10	0.098	0.016	16.6%	0.071	0.136
11	0.062	0.011	18.1%	0.043	0.088
12	0.064	0.012	18.0%	0.045	0.091
13	0.060	0.012	19.7%	0.041	0.089
14	0.023	0.006	25.1%	0.014	0.038
15+	0.019	0.006	31.3%	0.010	0.034

Table 33: Model 26.0 biological inputs at ages 1–15+, with weights for 2024–2026. Mortality is an annual instantaneous rate; maturity and female fraction are proportions; weights are kg per fish. Population weights are the BTS schedule. Native 2027–2028 projections carry the 2026 schedules. These inputs are treated as fixed within the fit.

Biological schedules and recent weights												
Age	Natural mortality	Mature	Female	Population weights (kg)			Fishery weights (kg)			Spawning weights (kg)		
				2024	2025	2026	2024	2025	2026	2024	2025	2026
1	0.900	0.000	0.500	0.017	0.020	0.020	0.025	0.025	0.025	0.085	0.085	0.085
2	0.450	0.008	0.500	0.090	0.079	0.066	0.186	0.186	0.186	0.196	0.196	0.196
3	0.300	0.289	0.500	0.309	0.226	0.397	0.435	0.314	0.644	0.387	0.415	0.317
4	0.300	0.641	0.500	0.454	0.526	0.544	0.481	0.556	0.501	0.452	0.523	0.541
5	0.300	0.842	0.500	0.560	0.587	0.640	0.560	0.621	0.697	0.521	0.596	0.657
6	0.300	0.901	0.500	0.607	0.676	0.725	0.617	0.693	0.756	0.633	0.668	0.732
7	0.300	0.947	0.500	0.693	0.734	0.809	0.696	0.730	0.834	0.750	0.779	0.804
8	0.300	0.963	0.500	0.923	0.812	0.874	0.825	0.855	0.883	0.879	0.893	0.911
9	0.300	0.970	0.500	0.997	0.903	0.949	1.016	0.966	0.970	0.987	1.016	1.019
10	0.300	1.000	0.500	1.020	1.087	1.027	1.017	1.140	1.087	1.055	1.116	1.135
11	0.300	1.000	0.500	1.126	1.066	1.048	1.078	1.194	1.192	1.127	1.175	1.227
12	0.300	1.000	0.500	1.137	1.150	1.192	1.251	1.082	1.240	1.245	1.239	1.279
13	0.300	1.000	0.500	1.259	1.258	1.416	1.308	1.314	1.312	1.294	1.347	1.333
14	0.300	1.000	0.500	1.406	1.399	1.449	1.328	1.343	1.420	1.336	1.388	1.434
15+	0.300	1.000	0.500	1.754	1.547	1.360	1.559	1.268	1.442	1.391	1.421	1.466

Table 34: Model 26.0 selectivity at age in 2026. Cells give the estimate and approximate 95% confidence interval. ATS and AVO share the same estimated curve. Relative selectivity can exceed one because the existing curve convention uses a different normalization than division by its maximum.

2026 selectivity and uncertainty			
Age	Fishery	BTS	ATS and AVO
1	0.000 (0.000–0.003)	0.054 (0.019–0.155)	0.000 (0.000–0.000)
2	0.016 (0.006–0.045)	0.026 (0.012–0.055)	0.776 (0.516–1.169)
3	0.145 (0.055–0.382)	0.066 (0.039–0.112)	0.789 (0.535–1.163)
4	0.337 (0.133–0.852)	0.156 (0.099–0.244)	0.709 (0.483–1.039)
5	0.646 (0.261–1.594)	0.326 (0.207–0.513)	0.779 (0.541–1.122)
6	1.589 (1.018–2.481)	0.559 (0.374–0.836)	0.974 (0.690–1.376)
7	1.556 (1.076–2.249)	0.769 (0.583–1.013)	1.141 (0.831–1.567)
8	1.536 (1.098–2.148)	0.897 (0.770–1.046)	1.125 (0.992–1.275)
9	1.429 (1.042–1.960)	0.958 (0.889–1.033)	1.125 (0.992–1.275)
10	1.354 (0.999–1.837)	0.984 (0.950–1.018)	1.125 (0.992–1.275)
11	1.295 (0.963–1.743)	0.994 (0.979–1.009)	1.125 (0.992–1.275)
12	1.274 (0.937–1.733)	0.998 (0.991–1.004)	1.125 (0.992–1.275)
13	1.274 (0.937–1.733)	0.999 (0.996–1.002)	1.125 (0.992–1.275)
14	1.274 (0.937–1.733)	1.000 (0.998–1.001)	1.125 (0.992–1.275)
15+	1.274 (0.937–1.733)	1.000 (0.999–1.000)	1.125 (0.992–1.275)

Table 35: Estimated parameter groups and their uncertainty in Model 26.0. Standard errors use each group's estimation scale; differences in scales make direct comparisons among groups inappropriate.

Estimated parameter groups					
Parameter group	Count	Minimum estimate	Maximum estimate	Median SE	Maximum SE
BTS age-1 deviation	44	−0.401	0.359	0.138	0.211
BTS age-1 log selectivity	1	−3.048	−3.048	0.307	0.307
BTS midpoint deviation	44	−0.351	0.192	0.099	0.123
BTS slope deviation	44	0.246	1.589	0.234	0.398
Initial-age deviation	14	−1.480	−0.395	0.651	0.803
Log catchability	3	−8.225	−0.248	0.071	0.185
Log fishing scale	63	−2.939	−0.821	0.134	0.233
Log logistic slope	1	−0.838	−0.838	0.196	0.196
Logistic midpoint	1	6.981	6.981	0.641	0.641
Recruitment deviation	63	−1.234	1.710	0.142	0.538
Recruitment level	1	9.673	9.673	0.121	0.121
Selectivity increment	987	−1.459	1.287	0.280	0.597

Table 36: Contributions to the Model 26.0 fit, grouped by data source or stabilizing assumption. Rounded rows may differ slightly from their sum. Negative values arise from likelihood constants. Contributions across different data and model structures use different scales and assumptions.

Contributions to the fit	
Contribution	Value
Catch data	−127.771
Composition data	421.207
Index data	160.956
Initial abundance deviates	19.860
Non-parametric selectivity	18.325
Recruitment deviates	71.193
Selectivity deviates	188.221

Table 37: Complete nonzero Model 26.0 likelihood and penalty contributions by data source or population-level process. Zero terms are retained in the complete numerical file. The saved exact total is retained in the supporting CSV. Negative contributions reflect likelihood normalization constants.

Likelihood and penalty contributions			
Component	Data stream or process	Weighted contribution	Unweighted contribution
Catch data	Fishery	−127.771	−127.771
Composition data	Fishery	177.319	177.319
Non-parametric selectivity	Fishery	16.538	16.538
Selectivity deviates	Fishery	129.726	129.726
Initial abundance deviates	Population	19.860	19.860
Recruitment deviates	Population	71.193	71.193
Index data	AVO	10.616	10.616
Non-parametric selectivity	AVO	1.788	1.788
Selectivity deviates	AVO	9.313	9.313
Index data	BTS	30.725	30.725
Composition data	BTS	205.885	205.885
Selectivity deviates	BTS	49.182	49.182
Index data	ATS	−5.488	−5.488
Composition data	ATS	38.003	38.003
Index data	ATS_1	30.596	30.596
Index data	CPUE	94.507	94.507

Table 38: Observed-index residual summaries for Model 26.0. Residuals measure observed-minus-predicted differences relative to the selected observation errors. The retained 2020 ATS placeholders are excluded.

Model 26.0 index residuals			
Series	Observations	Mean residual	Residual SD
ATS	20	−0.10	1.04
ATS age 1	19	0.06	1.13
AVO	20	0.12	0.97
BTS	44	0.09	1.19
Historical CPUE	12	0.05	0.53
BTS residuals account for the covariance among survey years. A spread near 1 agrees with the selected observation-error scale. These are fitted-model diagnostics.			

Table 39: Model 26.0 profile likelihood points and conditional 2027 female spawning biomass (thousand tons, kt), with approximate 95% confidence intervals. ATS catchability is fixed separately at each point; all other active parameters are re-estimated. The projection assumes 1,375 kt catch in 2027 and the mean 2023–2025 fishery selectivity curve.

2027 female spawning biomass at fixed ATS catchability				
ATS q	ΔNLL	2027 SSB (kt)	Lower 95% (kt)	Upper 95% (kt)
0.45000	2.477	3,567.3	2,737.3	4,649.0
0.45837	1.866	3,533.9	2,709.7	4,608.9
0.46674	1.349	3,501.6	2,682.8	4,570.3
0.47510	0.922	3,470.2	2,656.8	4,532.6
0.48347	0.581	3,439.7	2,631.5	4,496.0
0.49184	0.322	3,410.1	2,607.0	4,460.5
0.50021	0.141	3,381.2	2,583.6	4,425.0
0.50858	0.035	3,353.0	2,560.3	4,391.0
0.51694	0.000	3,325.5	2,538.0	4,357.3
0.53358	0.130	3,272.7	2,494.8	4,293.2
0.55021	0.505	3,222.5	2,453.8	4,232.0
0.56684	1.102	3,174.7	2,414.2	4,174.7
0.58347	1.901	3,129.3	2,376.6	4,120.4
0.60010	2.886	3,086.1	2,340.4	4,069.5
0.61674	4.039	3,045.0	2,306.3	4,020.4
0.63337	5.347	3,005.8	2,273.8	3,973.5
0.65000	6.797	2,968.3	2,242.7	3,928.7
Intervals use the fixed-q penalized Hessian and the delta method on log SSB. They propagate fitted-parameter uncertainty, including fitted abundance and selectivity, conditional on the stated catch, biological schedules and ATS q. They exclude uncertainty in ATS q, future recruitment process variation and weight-at-age schedules. Spawning occurs on April 1; age 15+ is retained.				

Table 40: Model 26.0 Mohn’s rho against the full-2026 fit. Rho is the mean proportional terminal revision.

Model 26.0 retrospective Mohn's rho			
Quantity	2024–2025 (2 peels)	2023–2025 (3 peels)	2017–2025 (9 peels)
Maximum fishing mortality	0.04411	0.15391	−0.06364
Age-1 recruitment	−0.26348	0.04013	−0.09848
Total biomass	−0.10792	−0.10774	0.00006
Female spawning biomass	−0.07636	−0.09393	0.12096

Table 41: Model 26.0 retrospective terminal estimates and revisions: All rows use the full-2026 reference. Each row omits all observations after its terminal year, using the same data-lag rules. Relative revisions are 100 × (peel estimate / full-2026-fit estimate for the same calendar year − 1). SSB is female spawning biomass; F is the maximum age-specific fishing mortality. These are point-estimate comparisons; the peel comparisons cover population estimates only.

Retrospective revisions in Model 26.0						
Terminal year	F revision (%)	Recruitment revision (%)	Biomass revision (%)	SSB revision (%)	Peel SSB (million t)	Full-fit SSB (million t)
2025	−9.7	−30.8	−11.0	−4.5	3.08	3.23
2024	18.5	−21.9	−10.5	−10.8	3.15	3.53
2023	37.4	64.7	−10.7	−12.9	3.29	3.78
2022	7.6	−35.0	1.6	2.3	3.83	3.74
2021	34.7	44.6	−9.8	−13.8	2.26	2.62
2020	−35.6	−59.0	−11.0	40.7	3.33	2.36
2019	−53.4	−79.4	0.8	28.2	3.93	3.07
2018	−17.2	−21.3	14.3	26.9	4.00	3.15
2017	−39.5	49.4	36.5	52.8	5.04	3.30

Table 42: Native fitting objectives with the selected catch inputs, including the Early-late fishery split comparison. Model-specific penalties and normalizations preclude direct likelihood ranking.

model	objective	method
26_0	751.989578	Established deterministic penalized fit
26_1_0	746.780971	Established deterministic penalized fit
26_1_1	726.234542	Established deterministic penalized fit
26_1_decline_zero	706.054843	Established deterministic penalized fit
26_1_2_conditional	731.019469	Early-late fishery split; Foreign 0.1235; Modern fixed at prior marginal estimate
Objectives have model-specific penalties and normalizations. The Early-late fishery split comparison supplies its native joint objective. These values support run verification; direct AIC or likelihood-ratio ranking is excluded.		

Table 43: Maximum ABC and OFL for the four separately fitted Rceattle configurations and the Early-late fishery split Model 26.1.2 comparison under matched Scenario 2 assumptions, in million tons. Model 26.0 (Rceattle) is the adopted assessment model and main proposal for Tier 3 calculations and specifications; Model 26.1.2 is the sensitivity analysis. Adopted catch specifications remain separate.

Maximum ABC and OFL under matched Scenario 2 catches				
Model	2027		2028	
	Maximum ABC (difference)	OFL	Maximum ABC (difference)	OFL
Model 26.0	2.222 (+0.00%)	2.746	2.319 (+0.00%)	2.870
Model 26.1.0	2.220 (-0.09%)	2.744	2.317 (-0.08%)	2.868
Model 26.1.1	2.191 (-1.39%)	2.699	2.308 (-0.47%)	2.846
26.1 decline penalty = 0	2.199 (-1.04%)	2.700	2.330 (+0.50%)	2.864
26.1.2 Early-late fishery split	2.187 (-1.59%)	2.692	2.316 (-0.10%)	2.853
Million tonnes; differences are relative to Model 26.0 in the same year. Recruitment is fixed at each model's historical arithmetic mean. The 26.1.2 row is the Early-late fishery split comparator. Simulation, retrospective and MCMC analyses are excluded. Adopted specifications remain unchanged.				

Table 44: Final Model 26.0 posterior stock and selectivity summaries from 5,000 saved draws. SD is posterior standard deviation; intervals are central 95% credible intervals. R-hat and chain ESS describe the saved chains.

Posterior stock and selectivity summaries					
Quantity and units	Mean	Posterior SD	Median	2.5%	97.5%
SSB 2026 (million t)	2.9645	0.2928	2.9465	2.4497	3.5844
Biomass 2026 (million t)	11.5860	1.4111	11.4616	9.2151	14.7192
Recruitment 2026 (billion fish)	36.4680	21.2479	31.1518	12.2880	91.0616
Ages1-3 2026 (billion fish)	56.0356	22.4635	50.9012	28.5376	113.5991
BTS selectivity age 1	0.0702	0.0354	0.0635	0.0207	0.1578
BTS selectivity age 2	0.0330	0.0120	0.0314	0.0152	0.0605
BTS selectivity age 5	0.3631	0.0795	0.3541	0.2332	0.5394
BTS selectivity age 10	0.9772	0.0215	0.9839	0.9172	0.9985
BTS selectivity age 15	0.9995	0.0009	0.9999	0.9969	1.0000

Table 45: Reference points for revised Model 26.0 and the Early-late fishery split Model 26.1.2 comparison. Biomass is female spawning biomass in million tons; F is the annual rate on the most selected age. Each column uses that model’s primary projection selectivity and fixed biological schedules.

Quantity	Model 26.0	26.1.2 Early-late fishery split	Units
B100%	6.234	6.296	Million tonnes
B40%	2.493	2.518	Million tonnes
B35%	2.182	2.204	Million tonnes
F40%	0.534	0.544	Per year
F35%	0.688	0.696	Per year
Revised fits; model-specific 1978–2023 arithmetic mean recruitment. These SPR reference points condition on fixed biology and the stated selectivity window.			

Table 46: Projections from the revised fits. Harvest scenarios under the Model 26.0 base fit. Each scenario retains its own rule; specified catches apply only to Scenario 2.

Scenario	Rule
1	Fishing at maximum permissible ABC each year.
2	Expected catch of 1,375 kt in 2027 and 2028, followed by maximum permissible ABC; author F fraction 1.
3	Mean annual maximum fishery F over 2021–2025, held constant.
4	One-half the maximum permissible ABC fishing mortality, without fixed-catch conditioning.
5	No fishing.
6	Fishing at the OFL threshold each year.
7	Maximum ABC fishing in 2027 and 2028, followed by OFL fishing from 2029.

Table 47: Projections from the revised fits. Short-term Tier 3 calculations under Scenario 2. Catch, spawning biomass, maximum ABC and OFL are thousand tons; F is the annual rate on the most selected age. The 2028 catch is a planning assumption.

Year	Assumed catch	Spawning biomass	Maximum ABC	OFL	F
2027	1,375	3,326	2,222	2,746	0.310
2028	1,375	3,670	2,319	2,870	0.296

Table 48: Projections from the revised fits. Thirteen-year catch table for the proposed harvest scenarios, in thousand tons.

Year	Catch (kt) by harvest scenario						
	1: Maximum ABC	2: Expected catch	3: Recent F	4: Half FABC	5: No fishing	6: OFL	7: ABC, then OFL
2026	1,313	1,313	1,313	1,313	1,313	1,313	1,313
2027	2,222	1,375	1,317	1,200	0	2,746	2,222
2028	2,103	1,375	1,379	1,274	0	2,446	2,103
2029	2,295	2,709	1,621	1,512	0	2,564	2,832
2030	2,136	2,370	1,600	1,505	0	2,320	2,458
2031	2,059	2,185	1,605	1,519	0	2,161	2,265
2032	1,881	1,943	1,505	1,431	0	1,889	1,916
2033	1,805	1,836	1,453	1,385	0	1,832	1,837
2034	1,771	1,786	1,426	1,360	0	1,823	1,823
2035	1,753	1,760	1,407	1,343	0	1,822	1,821
2036	1,744	1,748	1,396	1,332	0	1,822	1,822
2037	1,740	1,742	1,390	1,326	0	1,823	1,823
2038	1,737	1,738	1,385	1,321	0	1,823	1,823

Table 49: Projections from the revised fits. Thirteen-year female spawning-biomass table for the proposed harvest scenarios, in thousand tons.

Year	SSB (kt) by harvest scenario						
	1: Maximum ABC	2: Expected catch	3: Recent F	4: Half FABC	5: No fishing	6: OFL	7: ABC, then OFL
2026	3,057	3,057	3,057	3,057	3,057	3,057	3,057
2027	3,223	3,326	3,333	3,346	3,475	3,155	3,223
2028	3,270	3,670	3,691	3,746	4,341	3,037	3,270
2029	3,252	3,742	3,894	3,985	5,038	2,928	3,182
2030	3,045	3,320	3,831	3,948	5,439	2,680	2,809
2031	2,811	2,958	3,673	3,809	5,670	2,444	2,500
2032	2,656	2,732	3,542	3,688	5,845	2,323	2,338
2033	2,580	2,618	3,462	3,612	5,990	2,296	2,299
2034	2,538	2,558	3,405	3,556	6,081	2,290	2,290
2035	2,516	2,526	3,368	3,518	6,143	2,289	2,289
2036	2,505	2,510	3,342	3,491	6,177	2,289	2,289
2037	2,499	2,501	3,326	3,474	6,202	2,289	2,289
2038	2,496	2,497	3,317	3,463	6,217	2,289	2,289

Table 50: Thirteen-year fishing-mortality table for the proposed harvest scenarios. F is the annual rate on the most selected age.

Year	F (yr^{-1}) by harvest scenario						
	1: Maximum ABC	2: Expected catch	3: Recent F	4: Half FABC	5: No fishing	6: OFL	7: ABC, then OFL
2026	0.299	0.299	0.299	0.299	0.299	0.299	0.299
2027	0.534	0.310	0.295	0.267	0.00	0.688	0.534
2028	0.534	0.296	0.295	0.267	0.00	0.688	0.534
2029	0.534	0.534	0.295	0.267	0.00	0.688	0.688
2030	0.534	0.534	0.295	0.267	0.00	0.688	0.688
2031	0.534	0.534	0.295	0.267	0.00	0.674	0.688
2032	0.534	0.534	0.295	0.267	0.00	0.639	0.643
2033	0.534	0.534	0.295	0.267	0.00	0.631	0.632
2034	0.534	0.534	0.295	0.267	0.00	0.629	0.629
2035	0.534	0.534	0.295	0.267	0.00	0.629	0.629
2036	0.534	0.534	0.295	0.267	0.00	0.629	0.629
2037	0.534	0.534	0.295	0.267	0.00	0.629	0.629
2038	0.534	0.534	0.295	0.267	0.00	0.629	0.629

Table 51: Definitions of the nine Model 26.0 catch-decision indicators. All benchmarks are calculated from the Model 26.0 fit.

Probability event	Definition	Purpose	Model
P(F2027 > F35%)	2027 fishing mortality exceeds the rate giving 35% of unfished spawning potential per recruit.	Compare the catch scenario with the Tier 3 reference rate.	Both
P(B2028 < mean B)	2028 female spawning biomass at spawning is below mean estimated SSB over 1978–2025.	Compare the following year's stock with its historical mean.	Both
P(B2031 < mean B)	2031 female spawning biomass at spawning is below mean estimated SSB over 1978–2025.	Compare medium-term stock condition with its historical mean.	Both
P(B2031 < B2027)	2031 female spawning biomass at spawning is below the first projected-year value under the same scenario.	Describe medium-term change relative to 2027.	Both
P(B2029 < B20%)	2029 female spawning biomass at spawning is below 20% of the model-specific unfished reference biomass.	Describe the historical low-biomass benchmark used in this decision table; Model 26.0 uses mean-recruitment B100 and ADMB uses stock–recruit Bzero.	Both
P(Y2031 > Yhist)	The 2031 abundance ratio of ages 1–5 to ages 6–15+ exceeds the corresponding ratio pooled over 1964–2026.	Describe relative abundance of young fish. The historical threshold pools abundance across years, rather than averaging annual proportions.	Both
P(D2028 < D1994)	The effective number of mature-biomass age classes in 2028 is below its estimated 1994 value.	Describe near-term age diversity relative to a historical low-diversity year.	Both
P(D2031 < D1994)	The effective number of mature-biomass age classes in 2031 is below its estimated 1994 value.	Describe medium-term age diversity relative to 1994.	Both
P(E2027 > E2026)	Mean age-specific fishing mortality in 2027 exceeds mean age-specific fishing mortality in 2026.	Provide an F-based relative-effort proxy. Observed vessel effort, catchability changes and costs are outside this indicator.	Both
B denotes female spawning biomass at spawning. Y denotes the young-to-old abundance ratio. D is exp(−sum(p × log(p))), with p the beginning-year mature-biomass age shares. E is the mean-age F proxy. Model-specific biological schedules and reference points are retained.			

Table 52: Model 26.0 probabilities under eight alternative 2027 catches (kt), expressed as chances out of 100. Fishing mortality is set to achieve each 2027 catch and then held constant through 2031. Probabilities use local estimation uncertainty from the fit with updated AVO data, fixed biological schedules and mean future recruitment.

Probability event	2027 catch (kt)							
	10.0	317.5	635.0	952.5	1270.0	1587.5	1905.0	2540.0
P(F2027 > F35%)	0	0	0	0	0	0	0	27
P(B2028 < mean B)	1	2	3	5	9	13	18	32
P(B2031 < mean B)	0	0	1	3	8	17	32	71
P(B2031 < B2027)	0	0	2	7	20	43	69	98
P(B2029 < B20%)	0	0	0	0	0	0	0	1
P(Y2031 > Yhist)	0	0	0	1	6	17	32	59
P(D2028 < D1994)	1	1	1	2	3	4	6	12
P(D2031 < D1994)	0	0	0	0	0	1	5	29
P(E2027 > E2026)	0	0	0	0	45	93	99	100
Nine supported events. Assessment covariance is propagated through the catch solution, projections and estimated benchmarks. Future recruitment equals the estimated 1978–2023 mean; independent future recruitment innovations and weight-estimation uncertainty are excluded.								
Entries are approximate chances out of 100, rounded to whole numbers. A displayed 0 or 100 reflects rounding. The F solving each 2027 catch is held constant through 2031; subsequent catches vary with abundance.								

Table 54: Bycatch estimates of prohibited species in the BSAI directed pollock fishery, 1991-2026.

Year	Bairdi	Chinook	Halib	Halib.M	Herring	NonChin	Snow Crab	Red King	Blue King	Golden King
1991	1,398,107	39,054	2,156	NA	3,159	28,709	4,380,023	17,777	NA	NA
1992	1,500,765	33,672	2,220	NA	647	40,187	4,569,662	43,874	NA	NA
1993	1,649,103	36,619	1,326	NA	527	241,980	738,259	58,140	NA	NA
1994	371,214	31,890	963	689	1,627	92,011	811,734	42,361	NA	NA
1995	153,993	13,403	492	397	905	17,755	206,651	4,644	NA	NA
1996	89,416	55,472	382	321	1,242	77,174	63,398	5,934	NA	NA
1997	17,046	44,320	257	200	1,135	65,415	216,152	137	NA	NA
1998	57,071	51,245	353	278	801	60,677	123,758	14,287	NA	NA
1999	7,549	11,978	154	125	800	45,169	20,386	92	NA	NA
2000	1,517	4,961	110	91	483	58,568	6,580	NA	NA	NA
2001	5,146	33,444	243	200	225	57,004	5,709	106	NA	NA
2002	2,172	34,495	165	138	109	80,641	2,908	20	NA	NA
2003	771	45,661	89	74	967	188,339	750	52	9	0
2004	1,139	51,751	96	92	1,096	442,148	712	27	4	2
2005	631	68,126	120	112	594	703,080	2,260	0	0	1
2006	1,345	82,653	132	122	434	305,793	2,796	289	0	3
2007	1,497	121,956	312	290	352	86,380	3,165	7	0	3
2008	9,179	21,242	373	336	128	15,133	9,327	676	8	33
2009	6,204	12,731	541	457	65	45,960	7,392	1,137	20	0
2010	12,846	9,835	335	267	349	13,734	10,291	1,124	35	0
2011	10,986	25,499	460	378	377	193,754	6,584	577	26	0
2012	5,621	11,343	463	388	2,353	22,297	6,250	344	0	0
2013	12,427	13,091	334	272	959	125,525	8,704	508	34	107
2014	12,522	15,135	239	200	160	219,837	19,455	368	0	148
2015	8,872	18,329	152	130	1,487	237,803	8,339	0	0	0
2016	2,295	22,204	121	103	1,431	343,208	1,166	440	0	106
2017	7,270	30,079	97	89	964	467,750	3,407	203	0	67
2018	2,249	13,726	75	62	474	295,818	5,143	565	0	53
2019	3,146	25,038	134	113	1,102	348,631	6,228	453	99	445
2020	10,749	32,204	128	102	3,861	343,625	40,005	479	1	522
2021	8,417	13,852	145	131	1,709	546,472	4,668	52	0	115
2022	4,758	6,415	170	156	1,708	242,375	1,952	311	59	88
2023	11,997	11,874	88	70	3,087	112,512	4,100	54	0	132
2024	10,120	8,054	81	53	1,281	35,152	10,553	230	1	4
2025	4,822	19,825	79	66	3,030	151,457	14,424	1,018	1	2
2026	3,336	15,153	64	56	3,354	149,468	4,353	148	0	5

Source: author-provided AKFIN / NMFS Alaska Regional Office export, received from AKR October 27, 2026 and loaded November 7, 2026 (earlier source vintages remain in some historical records). Both pollock target categories and all reported gears are included. Herring, halibut catch (Halib), and halibut mortality (Halib.M) are metric tons; other columns are estimated numbers of individuals. Values are summed before rounding. NA denotes unavailable estimates; zero denotes a reported zero. Sums use available values when individual records are missing; halibut mortality is unavailable for 1991–1993 and has some missing records in 1998–2003. The 2026 values are preliminary, through the week ending October 24. Other King Crab is outside the species columns of the example table.

Pacific ocean perch (3,995 t) and Pacific cod (3,315 t) were the largest named bycatch categories in the pollock fishery in 2025, followed by flathead sole (1,813 t). Preliminary 2026 estimates through October 24 were 2,412 t of Pacific ocean perch, 6,549 t of Pacific cod, and 3,367 t of flathead sole.

Table 55: Bycatch estimates (t) of other species caught in the Bering Sea directed pollock fishery, 1991-2026.

Year	P.cod	Rock.Sole	Flathead	Arrowtooth	POP	Yellowfin	Sablefish	Shark	Atka.Mkrl	Skates	Other
1991	24,310	5,121	NA	5,719	418	418	10	NA	24	NA	10,973
1992	24,005	7,234	3	4,311	173	892	8	NA	261	NA	15,552
1993	20,931	8,714	NA	1,223	283	1,102	2	NA	40	NA	8,127
1994	14,409	3,010	NA	2,011	171	1,208	2	NA	44	NA	4,435
1995	19,777	2,180	2,175	1,177	142	675	12	NA	211	NA	1,897
1996	15,174	2,042	3,208	1,844	303	1,798	7	NA	384	NA	3,718
1997	8,263	1,523	2,350	985	428	606	2	NA	83	NA	3,187
1998	6,255	770	2,048	1,713	617	1,745	2	NA	10	NA	2,449
1999	3,220	1,059	1,885	273	121	350	7	NA	158	NA	1,740
2000	3,433	2,688	2,510	979	21	1,466	12	NA	2	NA	1,859
2001	3,880	1,673	2,199	530	574	594	21	NA	41	NA	3,226
2002	5,887	1,886	1,845	607	543	768	34	NA	221	NA	2,775
2003	5,968	1,419	1,501	618	935	210	48	NA	762	NA	1,992
2004	6,437	2,554	2,104	557	394	841	17	NA	1,053	NA	2,649
2005	7,413	1,125	2,352	651	652	63	11	NA	678	NA	2,759
2006	7,291	1,361	2,862	1,089	736	256	9	NA	789	NA	4,094
2007	5,630	510	4,226	2,795	625	86	12	NA	315	NA	3,731
2008	6,971	2,150	4,315	1,715	336	552	5	NA	15	NA	5,217
2009	7,875	7,591	4,666	2,202	114	271	3	NA	25	NA	4,969
2010	6,908	2,240	4,357	1,454	231	1,057	2	NA	55	NA	2,877
2011	10,043	8,481	4,886	1,602	658	1,083	1	66	900	2,352	1,156
2012	10,066	6,702	3,968	749	706	1,517	1	55	263	2,018	1,483
2013	8,955	6,379	3,146	966	611	2,097	0	43	70	1,750	746
2014	5,242	4,360	2,554	758	1,301	1,954	1	75	117	813	2,526
2015	8,303	1,709	2,260	403	2,525	863	0	52	202	824	2,875
2016	4,980	1,095	1,554	291	3,191	870	18	60	68	423	1,716
2017	6,151	1,708	908	204	4,438	578	96	91	61	448	2,120
2018	4,271	1,077	974	271	3,804	743	396	62	543	508	2,048
2019	6,213	1,117	1,087	422	7,972	444	1,237	101	367	509	790
2020	9,184	866	1,971	688	5,970	1,211	3,453	132	569	827	1,465
2021	9,104	830	1,531	413	2,468	754	1,105	173	545	912	566
2022	3,785	677	948	242	1,445	887	321	55	157	561	357
2023	3,821	549	844	245	1,345	750	487	271	40	400	458
2024	2,923	888	961	332	1,811	523	103	130	45	371	518
2025	3,315	557	1,813	293	3,995	231	51	155	12	438	873
2026	6,549	1,586	3,367	489	2,412	566	125	194	13	823	636

Source: supplied AKFIN / NMFS Alaska Regional Office FMP catch export. Estimates sum retained and discarded catch across both pollock target categories and all reported gears in the BS subarea. Pollock, salmon, halibut, Non TAC Species, and records without a species group are excluded. Other combines remaining named source groups, including other flatfish and rockfish, Greenland turbot, Alaska plaice, squid, octopus, sculpins, and historical Other / Other Species groups; definitions follow the source export rather than a fixed modern FMP list. NA means no records for that category and year, rather than a measured zero. Source group names and the complete column map are available with the annual CSV. Values are summed before rounding; units are metric tons. The 2026 estimates are preliminary through the week ending October 24.

Pollock catch in identified non-pollock fisheries declined from 46,996 t in 2024 to 29,263 t in 2025. Rock sole, yellowfin sole, and Pacific cod fisheries accounted for 10,776 t, 8,710 t, and 6,427 t, respectively, in 2025. The preliminary 2026 total through October 31 was 34,729 t; the different cutoffs and partial-year coverage qualify comparisons with the other bycatch tables.

Table 56: Pollock bycatch (t) in identified non-pollock directed Bering Sea fisheries, 1991-2026.

Year	Rock.sole	Other.flats	Other.spp	P.cod	Yellowfin.sole	Total
1991	9,711	7,992	756	10,695	NA	29,155
1992	9,825	1,372	568	20,778	13,101	45,644
1993	18,583	2,581	524	31,299	15,253	68,240
1994	15,784	6,770	89	26,594	33,200	82,438
1995	7,766	5,211	58	25,691	27,041	65,768
1996	7,699	5,457	205	22,382	22,255	57,997
1997	9,123	3,480	13	33,658	24,101	70,375
1998	3,960	3,011	61	10,469	15,339	32,840
1999	5,207	4,772	258	21,131	8,701	40,070
2000	5,481	7,068	43	14,509	13,425	40,526
2001	4,578	4,740	488	11,571	16,503	37,879
2002	9,942	2,220	52	15,256	14,490	41,960
2003	4,925	3,672	261	15,926	11,579	36,362
2004	8,976	6,396	194	18,650	10,384	44,600
2005	7,235	5,058	202	14,110	10,313	36,917
2006	6,986	3,826	143	15,168	5,967	32,091
2007	3,245	4,353	276	20,320	4,021	32,215
2008	4,931	4,822	17	9,534	9,828	29,132
2009	6,172	3,505	14	7,876	7,037	24,603
2010	6,074	3,317	86	6,576	5,179	21,232
2011	6,931	2,302	301	8,981	8,674	27,189
2012	6,704	1,751	413	8,377	11,198	28,443
2013	7,327	4,048	238	9,801	20,172	41,586
2014	11,259	6,404	203	11,502	24,713	54,080
2015	9,387	4,994	430	9,062	21,282	45,154
2016	11,850	3,687	450	9,071	22,306	47,365
2017	5,616	3,612	508	8,319	23,414	41,471
2018	5,182	3,525	957	8,008	28,235	45,908
2019	3,176	7,972	1,367	7,594	23,153	43,262
2020	6,401	2,373	728	5,512	31,651	46,666
2021	2,399	5,098	1,415	4,316	24,845	38,073
2022	2,977	5,505	2,627	6,260	26,515	43,884
2023	11,048	4,586	3,076	7,182	22,348	48,239
2024	15,126	4,356	1,929	6,690	18,895	46,996
2025	10,776	2,070	1,279	6,427	8,710	29,263
2026	13,847	7,040	1,345	6,599	5,898	34,729

Source: supplied AKFIN / NMFS Alaska Regional Office FMP catch export, BS subarea. Estimates include retained and discarded pollock catch and all reported gears. Pollock-directed trips, blank trip targets, and No retained catch targets are excluded. Other.flats combines flathead sole, arrowtooth flounder, other flatfish, Alaska plaice, Kamchatka flounder, and Greenland turbot targets. Other.spp combines the remaining identified targets: rockfish, sablefish, Atka mackerel, Other Species, and halibut. Rock.sole is the source Rock Sole - BSAI target; it is not restricted to a separately identified northern rock sole target. NA means no records for that target group and year. Total sums the included unrounded values; displayed columns may differ from Total by rounding. Units are metric tons. The 2026 estimates are preliminary through the week ending October 31. The complete target map and exclusion totals accompany the annual CSV.

Jellyfish dominated the selected nontarget-species catch by weight. Estimated jellyfish catch declined from 4,402 t in 2024 to 1,769 t in 2025. Miscellaneous fish contributed 99 t in 2025, with smaller catches in each of the other displayed groups. Preliminary jellyfish catch through October 24, 2026 was 4,418 t. These catch estimates describe fishery encounters and should be interpreted with fishing effort, distribution, and reporting coverage.

The historical source discrepancy extends beyond jellyfish: five other same-named columns also approximately doubled in 2019-2022, while the remaining four columns changed non-uniformly. Current estimates are rebuilt from named species groups rather than scaled from the 2024 SAFE table.

Table 57: Estimated catch (t) of selected nontarget-species groups in the BSAI directed pollock fishery, 2003-2026.

Year	Jellies	MiscFish	SeaStar	Osmerid	Grenadier	Eelpouts	Sea.pen	Anemone	Snails	Other
2003	5,644	101	89	10	21	7	1	0	1	2
2004	6,590	90	7	22	14	1	1	0	1	1
2005	5,198	158	10	13	14	1	2	0	7	3
2006	2,718	154	11	102	16	21	2	1	0	14
2007	2,403	203	5	140	28	119	4	0	0	13
2008	4,185	121	20	4	28	9	1	1	2	9
2009	8,118	135	10	6	5	4	3	1	1	3
2010	2,518	150	12	1	2	1	3	2	1	10
2011	8,233	277	27	2	2	1	3	2	1	8
2012	3,519	143	7	2	2	1	3	1	1	4
2013	5,294	121	15	1	2	2	2	2	1	10
2014	12,767	45	30	2	10	8	3	2	2	9
2015	4,951	90	42	24	4	11	2	2	1	5
2016	2,203	75	55	5	4	23	1	1	0	4
2017	6,153	48	12	3	3	18	1	1	1	2
2018	8,252	52	25	1	9	4	1	1	0	4
2019	3,889	73	51	1	9	2	1	0	0	6
2020	3,150	94	61	1	43	6	1	6	1	26
2021	7,831	35	20	0	55	1	2	3	1	3
2022	7,609	23	185	1	11	0	1	1	1	11
2023	7,088	38	28	0	86	1	2	3	1	6
2024	4,402	43	16	3	18	3	1	4	0	3
2025	1,769	99	8	4	6	14	2	1	1	3
2026	4,418	68	5	3	1	1	2	0	1	2

Source: supplied AKFIN / NMFS Alaska Regional Office nontarget catch-accounting export. BSAI records (BS and AI) are included; GOA and inside-water records are excluded. Both pollock target categories, all reported gears, and retained and discarded records are combined using Nontarget Estimate (mt), not the species-count field. Squid, sculpins, and seabirds are outside the selected groups of this table and are excluded. MiscFish is the source Misc fish group. Osmerid combines capelin, eulachon, other osmerids, family Osmeridae smelts, and surf smelt; Grenadier combines all named grenadier groups. Other contains the remaining selected fish and invertebrate groups; the complete mapping accompanies the annual CSV. NA means no records for a group-year; a displayed zero may be a positive estimate below 0.5 t. Values are summed before rounding to whole metric tons. The 2026 estimates are preliminary through the week ending October 24. The November 2024 summary approximately doubled the 2019-2022 Jellies, MiscFish, SeaStar, Osmerid, Grenadier, and Eelpouts columns relative to September. Sea pen, Anemone, Snails, and Other changed non-uniformly, with cross-column numerical matches indicating changed alignment or allocation. These are comparisons of historical headers. The current table uses explicit source species names and both catch dispositions, with no scaling adjustment. The upstream preparation step remains unconfirmed. View the jellyfish source-history and filter audit.

Figures



Figure 1: Nominal seasonal catch per tow-hour for herring, sablefish, Chinook salmon, chum salmon, and pollock in available EBS pollock-dominated haul records, 1991-2026, with source vessel types 1-3 combined. Each rate is the sum of available extrapolated catch divided by the sum of positive valid haul hours in the same year and season; the calculation uses the ratio of totals. Weight rates are kg per tow-hour for herring and sablefish and tonnes per tow-hour for pollock. Salmon rates are extrapolated fish per tow-hour. A season is January-May and B season is June-December, retaining the original figure's calendar split. Selection requires recorded NMFS area 500-539, positive official total catch, and pollock kg / official total catch in tonnes >800 (a >80% pollock proxy). 2026 values are preliminary; B-season records extend through September 18, 2026 and are partial. Historical years use their full available seasons. Blank species entries are omitted from available catch sums, retained in the effort denominator, and counted in the plotting table. Entirely unreported species-season combinations are gaps, including sablefish in 2014. The supplied file has incomplete CV/EM coverage; no CV records pass the target filter in 2025-2026. Rates describe the available source records; fleet-wide catch accounting uses a separate dataset.

Source values and reproduction records are retained in the technical supplement.



Figure 2: Estimated EBS pollock catch numbers (thousands of fish) by sex for the A season (January-May), B season (June-December), and annual total, 1991-2025. Circles denote females and triangles denote males. B-season estimates combine the northwest and southeast strata, and annual totals equal A+B. Shading marks provisional 2024-2025 estimates on the original sampler age basis.

This provisional summary uses saved sampler point estimates, with ages 1-14 and 15+ included once. Catch in tonnes and fish weights in kg make raw N thousands of fish; the original scale is retained. Sex labels have been checked against the source records. The 1998 catch allocation agrees with the alternate historical three-stratum control and archived point outputs. Historical source records remain incomplete; the exact discrepancy is retained in the technical appendix. The 2024-2025 estimates retain original FTNIR ages; the incomplete 2024 TMA conversion is excluded. Both new years have all-12-month source coverage; their provisional status reflects the age basis and input lineage. This figure uses the saved sampler estimates.

Source values and reproduction records are retained in the technical supplement.



Figure 3: Cumulative EBS pollock catch as a proportion of annual specified TAC for weeks ending in January, February, March, and April, 2000–2026. Catch includes retained and discarded pollock from all gears and targets in the Bering Sea subarea, excluding Bogoslof reporting area 518. Each full reporting week is assigned to its week-ending month. The denominator is the annual stock TAC from harvest specifications, matching the original figure convention, before later AI-to-BS quota transfers. Amendment 110 shifted five percentage points of the directed fishery seasonal allocation from B to A to provide greater flexibility to avoid Chinook salmon bycatch; 2017 was the first A season after the rule took effect.

January–April 2026 has elapsed; the weekly observations remain revisable. The source's maximum week-ending label is October 17, 2026, while its file modification time is October 15; an authenticated extraction timestamp is unavailable. This figure summarizes cumulative January–April week-ending groups, within the regulatory A season extending to June 10. The denominator is specified annual stock TAC before quota transfers, with final transferred account quotas tracked separately. Allocation context comes from the Amendment 110 final rule and 2026 harvest specifications; annual differences may reflect several influences.

Source values and reproduction records are retained in the technical supplement.



Figure 4: Pollock roe production in the supplied EBS source series by calendar season group, A (January–May) and B (June–December), 2000–2026. Production combines at-sea records labeled Bering Sea (BS) and inshore records labeled Bering Sea and Aleutian Islands (BSAI). Values are roe product metric tonnes. Open markers and dashed links identify provisional 2026 values. The 2026 export contains January–October; the B group is partial and October completeness is unverified. Calendar groups follow the original figure definition; applying the June 10 regulatory season boundary requires daily production records.

The user-identified production export also supplies January–December 2025 coverage, with B-group production of 1,795.8 product tonnes, replacing the older partial 2025 estimate of 1,321.1 tonnes. Provisional 2026 totals are 19,092.9 product tonnes for A and 406.4 tonnes for the available June–October B records. The source combines offshore BS and inshore BSAI production; an exclusively EBS inshore subtotal remains unavailable. Calendar groups use production month; resolving the June 10 regulatory boundary or exact fish capture date requires finer records. October completeness and an authenticated query/extraction cutoff are unverified. Values retain product-tonne units, and unavailable months remain unobserved.

Source values and reproduction records are retained in the technical supplement.



Figure 5: Spatial distribution of observed pollock catch in the A season, 2024–2026, using January 20–June 9 in each year. Vertical bar height represents recorded catch tonnes within 0.5-degree longitude by 0.25-degree latitude cells; the height scale is fixed across years within each season, with separate A- and B-season scales and a 10,000-t reference column in each panel. Data are from inseas26.csv and include available catcher-processor, mothership and observed catcher-vessel records in NMFS areas below 540. Electronic-monitoring catcher-vessel records are absent. These are observed-record catch totals rather than fleet-wide expanded catches. The B-season comparison is restricted to the September 18 cutoff in all three years. Rendered with GMT using the AKsf coastline and the local bottom_topo.grd bathymetry grid. Blue shades show ocean depth and green shades show land elevation. Mercator projection viewed at azimuth 155 degrees and elevation 35 degrees, with increased column heights; vertical bars represent catch rather than seafloor height. Updated in the style of Figures 4 and 7 of the 2024 assessment.



Figure 6: Spatial distribution of observed pollock catch in the B season, 2024–2026, using June 10–September 18 in each year. Vertical bar height represents recorded catch tonnes within 0.5-degree longitude by 0.25-degree latitude cells; the height scale is fixed across years within each season, with separate A- and B-season scales and a 10,000-t reference column in each panel. Data are from inseas26.csv and include available catcher-processor, mothership and observed catcher-vessel records in NMFS areas below 540. Electronic-monitoring catcher-vessel records are absent. These are observed-record catch totals rather than fleet-wide expanded catches. The B-season comparison is restricted to the September 18 cutoff in all three years. Rendered with GMT using the AKsf coastline and the local bottom_topo.grd bathymetry grid. Blue shades show ocean depth and green shades show land elevation. Mercator projection viewed at azimuth 155 degrees and elevation 35 degrees, with increased column heights; vertical bars represent catch rather than seafloor height. Updated in the style of Figures 4 and 7 of the 2024 assessment.



Figure 7: Fleet dispersion in the A and B seasons, 2000–2026, recalculated from inseas26.csv using the 2024 assessment's daily pairwise-distance approach. Unique tow pairs are formed within each day and vessel group (catcher-processors and observed catcher vessels); seasonal means pool distance sums and pair counts across both groups. No cross-group pairs are included. Distances are great-circle nautical miles. The legacy boundary assigns day of year 161 and earlier to A and later days to B. Solid lines use available seasonal records; dashed lines restrict every year to September 18. The 2026 B season is partial. Records require more than 80% pollock by catch weight, valid positions, EBS reporting areas 500–539, and tow duration between zero and 40 hours. Duration is calculated from timestamps with recorded minutes as fallback. Vessel-type codes replace the unavailable historical interactive vessel lists; electronic-monitoring catcher-vessel records are absent.



Figure 8: Catch-weighted distributions of mean pollock weight per source record for A and B seasons, 1991–2026. Catch tonnes are grouped in 50-g bins; the vertical line marks 500 g. A is source week 24 or earlier and B is later than week 24. The extract ends September 18, 2026, so B-season coverage is partial. Available catcher-vessel records exclude electronic-monitoring records. Ridges describe the distribution of record means, with heights scaled for display.



Figure 9: Weekly A-season-window distributions of mean pollock weight, 2001–2026, for source weeks 2–15. Catch tonnes are grouped in approximately 66.7-g bins; the vertical line marks 500 g. Source weeks are seven-day periods beginning January 1. The window differs from the regulatory A season. Available catcher-vessel coverage is incomplete.



Figure 10: Weekly B-season-window distributions of mean pollock weight, 2001–2026, for source weeks 25–40. Catch tonnes are grouped in approximately 66.7-g bins; the vertical line marks 500 g. The 2026 extract ends September 18 and therefore covers only part of this window. Available catcher-vessel coverage is incomplete.



Figure 11: Eastern Bering Sea pollock catch, 1964–2026, in million tons. Catches for 1964–1990 are undifferentiated season-area catch; from 1991 the A-season combines northwest and southeast records, with separate NW and SE B-season catches. The source recipe classifies reported week-ending dates before June 1 as A and all remaining dates as B (June–December), including small November–December catches. NW comprises NMFS areas 520–529; SE comprises 500–517 and 519. Annual bars avoid interpolating splits at the 1990/1991 boundary. The 2026 bar includes 1.268 million tons of recorded catch through October 17 and is incomplete; it excludes the annual completion forecast.

Source values and reproduction records are retained in the technical supplement.



Figure 12: Bottom-trawl pollock CPUE and bottom-temperature contours in the eastern Bering Sea, 2018, 2019 and 2021–2026, reproduced from the AKsf pollock-mapping example (Ianelli 2026), snapshot dated 30 September 2026. Blue points mark sampled tows; contours are labeled 0, 2 and 4 °C. The common CPUE color scale saturates at 1,000 kg/ha. Surfaces are descriptive local interpolations within survey coverage, with no spatial prediction intervals. The supplied CPUE includes a density-dependent correction whose version and adoption status are unspecified in the example. The 2020 survey is absent.



Figure 13: BTS abundance at length on a relative scale, 1987–2026. Annual length distributions are arranged from top to bottom, with length in centimetres on the horizontal axis. The gap after 2019 marks the unsampled 2020 survey year.



Figure 14: BTS age composition by survey year, 1990–2026. Ages increase from left to right and years run from top to bottom. Highlighted bars trace the 1992 (green), 2012 (blue) and 2018 (red) year classes as they advance through successive ages. The blank 2020 row marks the unsampled survey year.



Figure 15: ATS age composition by survey year, 1994–2026. Ages increase from left to right, with ages 10 and older pooled as 10+. Blank rows indicate years lacking age-composition observations. Red bars follow the 2018 year class. The 2026 ages are provisional estimates derived by applying the BTS age–length key to ATS population length frequencies.



Figure 16: Provisional core-area 2026 ATS abundance at age after applying a BTS FT-NIRS length key (upper panel) and the screened 307-pair BTS conditional tester-age key (lower panel). Bars show billion fish over supported lengths only, with no uncertainty intervals. Released FT-NIRS age 15 is unpooled; the TMA terminal bar is 15+.



Figure 17: Provisional core-area 2026 ATS mean weight at TMA tester age, obtained by dividing biomass after the unchanged BTS length-key transfer and screened 307-pair tester-key transfer by the matching abundance estimates. The 15+ point uses pooled biomass and numbers for exact ages 15–18. These point estimates cover supported lengths only.



Figure 18: Historical pollock biomass between near surface and 3.0 m above the bottom in the summer eastern Bering Sea acoustic-trawl surveys of the U.S. EEZ, 1994–2026. Biomass between 3.0 m and the seafloor is excluded. The 2026 panel is preliminary; the legend gives biomass in kg.



Figure 19: ATS core-shelf biomass from the historical series. Bars show approximate normal 95% intervals using survey relative errors. The line connects observed survey estimates; missing calendar years have no observation. The orange triangle marks preliminary 2026.



Figure 20: Preliminary 2026 ATS pollock distribution, including the Northern Extension. Colors and GMT contours show gridded mean interval biomass (tons per EDSU), using all 10,924 observations. Approximately 10-nautical-mile cell means are smoothed and displayed within 30 km of sampled cell centers. Dashed gray lines mark 100 and 200 m depths. Coastline: AKsf.



Figure 21: Annual AVO relative backscatter index for 2009–2026, with the source 95% confidence intervals rescaled by the same mean. The dashed line marks a relative index of 1. The line has a gap at 2020; the orange triangle identifies 2026.



Figure 22: AVO pollock backscatter, 2006–2026 (m²/nmi²), mapped with common GMT gridding and contour levels. Twenty survey years are shown; AVO observations are unavailable for 2020. Cell means include observed zeros and use an approximately 20-nautical-mile grid. Surfaces are displayed within 45 km of sampled cell centers. Dashed gray lines mark 100 and 200 m depths. Coastline: AKsf. The 2026 observations are preliminary.



Figure 23: Cumulative input-update comparisons within Model 23.0, 2000–2026. Each fit includes the preceding updates. Biomass uses the same population-weight schedule in every comparison, and fishing mortality is the maximum realized rate across ages. ATS and AVO updates combine observations, historical revisions and the stated error calibrations. Lines show point estimates. Historical stages retain their actual source identities.



Figure 24: Projected 2027 spawning biomass and combined abundance at ages 1–3 across five selected cumulative Model 23.0 input-update stages. The final saved fits use estimated annual 2026 catch of 1,310,000 t. The displayed stages are cumulative; the omitted weight-at-age and A-season spawning-weight updates remain included in the fishery-age and later stages. Lower-panel columns stack ages 1, 2 and 3 at the start of 2027, before removals; spawning biomass includes mortality to spawning time. Saved diagnostic projection conventions are retained and documented in the technical supplement.



Figure 25: Approximate distributions of spawning-biomass change from 2026 to 2027 across five selected cumulative input-update stages, followed by Models 23.2 and 26.0 fitted to the same assessment data. The final saved fits use estimated annual 2026 catch of 1,310,000 t. The displayed stages are cumulative; the omitted weight-at-age and A-season spawning-weight updates remain included in the fishery-age and later stages. Green area represents an increase and orange area a decrease. Dots show estimated changes, bars show 95% local asymptotic ranges, and labels give approximate chances of an increase rounded to whole percentages. Matching fitted covariance and saved diagnostic projection conventions are retained; methods are documented in the technical supplement.



Figure 26: Approximate distributions of change from 2025 to 2026 in total numbers at ages 1–4. Five selected cumulative input-update stages are followed by Models 23.2 and 26.0 fitted to the same assessment data. The displayed stages are cumulative; the omitted weight-at-age and A-season spawning-weight updates remain included in the fishery-age and later stages. Annual totals sum ages 1–4 at the start of the year and retain both cross-age and cross-year fitted covariance. Green area represents an increase and orange area a decrease. Dots show estimates, bars show 95% uncertainty ranges and labels give approximate chances of an increase. This historical contrast retains its saved values.



Figure 27: Models 23.0 and 23.2 fitted to the same observations and supplied biological schedules. Lines show annual estimates and shading gives approximate 95% local asymptotic uncertainty ranges. Panels show biomass at common population weights, spawning biomass, age-1 recruitment and maximum realized fishing mortality across ages.



Figure 28: Models 23.2 and 26.0 fitted to the same observations and supplied biological schedules. Lines show annual estimates and shading gives approximate 95% local asymptotic uncertainty ranges. Panels show biomass at common population weights, spawning biomass, age-1 recruitment and maximum realized fishing mortality across ages. These estimation ranges are conditional on the stated model and supplied inputs.



Figure 29: Model 26.0 annual total biomass, female spawning biomass, age-1 recruitment and fishing mortality through 2026, with approximate 95% asymptotic estimation ranges. Total biomass uses population weights; spawning biomass uses mature female spawning weights and survival to spawning. Recruitment is labeled by the year fish enter age 1; fishing mortality is the maximum realized rate across ages. Supplied biological schedules are fixed.

Model 26.0 numbers at age

1964-1989 | abundance relative to each age's 1964-2026 mean
Cells: abundance / age-specific mean (CV%); CV = asymptotic SE as a percentage of the estimate



Colour denominators: each age's arithmetic mean over 1964-2026 (all 63 years).
Labels: multiples to 2 decimals; CV to whole %. 1 = age mean; 2 = twice the mean.
10+ sums ages 10-14 and 15+ once; its SE retains all cross-age covariance.

Figure 30: Model 26.0 start-of-year numbers at age, 1964–1989, followed by 1990–2026 as the same figure continued. The color legend shows multiples of the age-specific mean. Cells show abundance as multiples of each age-group mean, rounded to two decimals, with asymptotic CV (%) in parentheses. Each age-group mean uses the full 1964–2026 period. Ages 10–14 and 15+ are pooled once into 10+; its SE uses the summed saved Jacobian and full matching parameter covariance. Both pages use identical scales, denominators and cell sizing.

Figure 30 (continued): **1990–2026**. Age groups, full-period denominators, scales and row sizing match the preceding page.

Model 26.0 numbers at age (continued)

1990-2026 | abundance relative to each age's 1964-2026 mean
Cells: abundance / age-specific mean (CV%); CV = asymptotic SE as a percentage of the estimate

Year	Age at start of year 1	2	3	4	5	6	7	8	9	10+
1990	2.14 (4%)	0.51 (8%)	0.25 (12%)	0.34 (11%)	0.62 (10%)	1.74 (7%)	0.86 (12%)	3.52 (8%)	1.58 (13%)	2.85 (9%)
1991	1.16 (5%)	2.18 (4%)	0.53 (8%)	0.27 (12%)	0.35 (11%)	0.60 (10%)	1.77 (7%)	0.86 (12%)	3.69 (8%)	2.24 (9%)
1992	1.01 (6%)	1.18 (5%)	2.23 (4%)	0.56 (8%)	0.28 (12%)	0.35 (13%)	0.61 (12%)	1.69 (9%)	0.91 (14%)	2.55 (9%)
1993	1.96 (4%)	1.03 (6%)	1.21 (5%)	2.31 (4%)	0.58 (8%)	0.28 (13%)	0.33 (15%)	0.54 (15%)	1.48 (13%)	1.47 (11%)
1994	0.68 (6%)	2.00 (4%)	1.05 (6%)	1.28 (5%)	2.26 (4%)	0.62 (9%)	0.30 (14%)	0.33 (17%)	0.56 (17%)	1.39 (11%)
1995	0.47 (7%)	0.69 (6%)	2.05 (4%)	1.14 (6%)	1.34 (6%)	2.19 (5%)	0.66 (9%)	0.32 (15%)	0.37 (18%)	1.07 (12%)
1996	0.96 (5%)	0.48 (7%)	0.71 (6%)	2.24 (4%)	1.25 (6%)	1.40 (6%)	2.10 (6%)	0.68 (10%)	0.35 (16%)	0.80 (12%)
1997	1.29 (4%)	0.98 (5%)	0.49 (7%)	0.77 (6%)	2.50 (4%)	1.41 (6%)	1.45 (7%)	1.94 (8%)	0.66 (12%)	0.59 (12%)
1998	0.63 (5%)	1.31 (4%)	1.00 (5%)	0.54 (7%)	0.84 (6%)	2.78 (4%)	1.54 (7%)	1.47 (9%)	2.01 (10%)	0.58 (12%)
1999	0.68 (5%)	0.65 (5%)	1.34 (4%)	1.08 (5%)	0.58 (7%)	0.93 (7%)	2.97 (5%)	1.71 (7%)	1.61 (10%)	1.13 (11%)
2000	1.04 (4%)	0.69 (5%)	0.66 (5%)	1.42 (4%)	1.15 (5%)	0.65 (8%)	1.06 (8%)	3.22 (6%)	1.97 (8%)	1.39 (10%)
2001	1.46 (3%)	1.06 (4%)	0.71 (5%)	0.71 (5%)	1.47 (4%)	1.21 (6%)	0.72 (8%)	1.13 (9%)	3.44 (7%)	1.68 (8%)
2002	0.97 (4%)	1.49 (3%)	1.09 (4%)	0.77 (5%)	0.75 (6%)	1.45 (5%)	1.17 (7%)	0.74 (9%)	1.22 (10%)	2.42 (7%)
2003	0.61 (5%)	0.99 (4%)	1.53 (3%)	1.17 (4%)	0.80 (5%)	0.75 (6%)	1.29 (6%)	1.09 (8%)	0.74 (11%)	1.87 (7%)
2004	0.28 (6%)	0.62 (5%)	1.02 (4%)	1.60 (3%)	1.22 (4%)	0.77 (6%)	0.69 (8%)	1.15 (8%)	1.04 (10%)	1.36 (8%)
2005	0.20 (8%)	0.28 (6%)	0.64 (5%)	1.10 (4%)	1.54 (4%)	1.20 (5%)	0.76 (7%)	0.64 (9%)	1.13 (9%)	1.20 (8%)
2006	0.49 (5%)	0.20 (8%)	0.29 (6%)	0.69 (5%)	1.11 (4%)	1.38 (4%)	1.11 (6%)	0.75 (8%)	0.67 (10%)	1.15 (8%)
2007	1.06 (4%)	0.50 (5%)	0.21 (8%)	0.30 (7%)	0.68 (5%)	1.02 (5%)	1.19 (5%)	1.03 (7%)	0.73 (9%)	0.89 (8%)
2008	0.57 (6%)	1.08 (4%)	0.51 (5%)	0.22 (8%)	0.30 (7%)	0.62 (6%)	0.85 (6%)	1.07 (7%)	1.00 (8%)	0.75 (9%)
2009	2.10 (3%)	0.59 (6%)	1.11 (4%)	0.55 (5%)	0.22 (8%)	0.28 (8%)	0.52 (7%)	0.75 (8%)	1.00 (9%)	0.74 (9%)
2010	0.98 (5%)	2.14 (3%)	0.60 (6%)	1.18 (4%)	0.55 (6%)	0.21 (9%)	0.25 (11%)	0.48 (10%)	0.73 (9%)	0.74 (9%)
2011	0.60 (6%)	1.00 (5%)	2.20 (3%)	0.65 (6%)	1.16 (5%)	0.55 (7%)	0.22 (11%)	0.26 (13%)	0.49 (12%)	0.68 (10%)
2012	0.52 (6%)	0.61 (6%)	1.03 (5%)	2.38 (3%)	0.70 (6%)	0.96 (6%)	0.46 (9%)	0.20 (13%)	0.24 (16%)	0.51 (12%)
2013	1.84 (3%)	0.53 (6%)	0.63 (6%)	1.11 (5%)	2.39 (4%)	0.74 (7%)	0.81 (8%)	0.41 (11%)	0.18 (17%)	0.33 (15%)
2014	2.10 (3%)	1.87 (3%)	0.55 (6%)	0.68 (6%)	1.14 (5%)	2.36 (4%)	0.77 (8%)	0.78 (11%)	0.38 (16%)	0.22 (19%)
2015	0.90 (5%)	2.14 (3%)	1.92 (3%)	0.59 (6%)	0.71 (6%)	1.16 (6%)	2.32 (5%)	0.72 (10%)	0.77 (14%)	0.25 (18%)
2016	0.37 (9%)	0.92 (5%)	2.20 (3%)	1.98 (4%)	0.60 (6%)	0.73 (7%)	1.13 (7%)	2.19 (7%)	0.71 (12%)	0.40 (17%)
2017	0.33 (10%)	0.37 (9%)	0.94 (5%)	2.39 (4%)	1.79 (4%)	0.62 (7%)	0.77 (8%)	1.21 (9%)	2.24 (9%)	0.50 (14%)
2018	0.71 (8%)	0.34 (10%)	0.38 (9%)	1.03 (5%)	2.53 (4%)	1.64 (5%)	0.61 (8%)	0.79 (9%)	1.28 (10%)	1.15 (12%)
2019	3.72 (5%)	0.73 (8%)	0.35 (10%)	0.42 (9%)	1.13 (6%)	2.41 (4%)	1.61 (6%)	0.64 (9%)	0.86 (11%)	1.20 (11%)
2020	1.17 (9%)	3.79 (5%)	0.75 (8%)	0.38 (10%)	0.46 (9%)	1.27 (6%)	2.24 (6%)	1.40 (9%)	0.63 (11%)	1.02 (11%)
2021	0.55 (14%)	1.19 (9%)	3.87 (5%)	0.80 (8%)	0.40 (10%)	0.49 (10%)	1.16 (7%)	1.98 (8%)	1.36 (11%)	0.75 (14%)
2022	0.54 (17%)	0.56 (14%)	1.21 (9%)	3.99 (6%)	0.85 (8%)	0.43 (11%)	0.52 (11%)	1.08 (10%)	1.83 (11%)	0.88 (14%)
2023	0.49 (20%)	0.55 (17%)	0.57 (14%)	1.29 (9%)	4.18 (6%)	0.93 (9%)	0.48 (13%)	0.57 (13%)	1.13 (12%)	1.26 (12%)
2024	1.66 (23%)	0.50 (20%)	0.56 (17%)	0.61 (14%)	1.38 (10%)	4.26 (7%)	0.99 (10%)	0.54 (14%)	0.65 (15%)	1.21 (13%)
2025	1.19 (27%)	1.69 (23%)	0.52 (20%)	0.60 (18%)	0.65 (15%)	1.47 (11%)	4.33 (9%)	1.11 (12%)	0.64 (15%)	1.01 (13%)
2026	1.54 (53%)	1.21 (27%)	1.74 (23%)	0.55 (21%)	0.64 (19%)	0.71 (17%)	1.42 (14%)	4.37 (12%)	1.23 (14%)	0.88 (14%)



Colour denominators: each age's arithmetic mean over 1964-2026 (all 63 years).
Labels: multiples to 2 decimals; CV to whole %. 1 = age mean; 2 = twice the mean.
10+ sums ages 10-14 and 15+ once; its SE retains all cross-age covariance.





Figure 31: Age-1 recruitment from the accepted 2024 assessment, Model 23.0, compared with Models 26.0 (panel A) and 26.1.2 (panel B). Bars show point estimates for both sexes in billions of fish. Historical estimates are repeated unchanged without intervals. Whiskers show local 95% log-scale intervals: the native asymptotic covariance for Model 26.0 and the marginal fit's Louis observed information for Model 26.1.2, conditional on its integration samples and normalizer grid. Calendar year denotes age 1; the 2018 year class enters at age 1 in 2019.



Figure 32: Model 26.0 log recruits per spawner, updating the two-panel diagnostic from the 2024 assessment. Age-1 recruits in year t+1 (million fish) are divided by female spawning biomass in year t (thousand tonnes), then natural-log transformed. The numerical ratio is recruits per kilogram of female spawning biomass. Upper panel: birth-year series, 1964–2025. Lower panel: the same ratios against spawning biomass, with two-digit birth-year labels and lines joining consecutive years. Orange identifies the less certain 2023–2025 year classes. The figure shows point estimates of the ratio.



Mortality, maturity, sex proportions and weights are supplied inputs. Shaded selectivity bands show 95% uncertainty ranges.

Figure 33: Model 26.0 biological and selectivity schedules. Maturity, female fraction, natural mortality and weights are supplied inputs; displayed weights are for 2026. Fishing and survey selectivity curves are estimated, with approximate 95% asymptotic ranges. ATS and AVO share their estimated selectivity curve. Fixed supplied schedules have no fitted-parameter bands.



Figure 34: Model 26.0 instantaneous fishing mortality by age through 2026, using the verified fit. Tiles and contours show the same realized total fishing mortality rates; age 15+ remains the terminal group.



Figure 35: Model 26.0 spawning exploitation for spawning-to-spawning intervals starting in 1964–2026. Black lines show the revised production fit; dashed blue lines and shading show MCMC posterior medians and central 95% credible intervals from 5,000 existing draws. Each within-draw ratio compares next-spawning biomass with and without fishing from a shared spawning-date population. The terminal interval assumes 1.375 million t catch in 2027, within-draw mean 1978–2023 recruitment and mean 2023–2025 selectivity, and the revised deterministic fishery-weight schedule. Supplied biology is fixed. The terminal posterior median is 14.5333%, with interval 11.2075–18.3700%.



Figure 36: Observed indices and Model 26.0 fitted predictions. Black points show source observations, lines show fitted predictions, and bars show approximate 95% observation-error ranges. BTS uses its supplied covariance. The ATS age-1 panel uses a logarithmic scale in billions of fish; the retained 2020 age-1 placeholder is distinguished from the real 2020 USV biomass observation.

BTS errors account for the supplied covariance among survey years. Dashed lines: standard-normal 95% reference.
The gray 2020 ATS age-1 cross is an inherited placeholder, excluded from observed-index summaries.
The fitted 2020 USV biomass remains an observed ATS index.

Figure 37: Model 26.0 index residuals relative to the selected observation errors, with BTS accounting for supplied covariance among survey years. Dashed lines mark the standard-normal 95% reference range. The retained 2020 ATS age-1 placeholder is distinguished and excluded from observed-index summaries; the 2020 USV biomass remains an observation.



Figure 38: Annual fishery selectivity at ages 1–15+ from the Model 26.0 fit. Each ridge shows the estimated age pattern for that year, using the normalization recorded with the verified figure data.



Figure 39: Annual bottom-trawl survey selectivity at ages 1–15+ from the Model 26.0 fit. Each ridge is divided by its maximum over the displayed ages. The taller, narrower display compares annual age-pattern shape, including years without an age-composition sample.



Figure 40: Annual acoustic-trawl survey selectivity at ages 2–15+ from the Model 26.0 fit. ATS and AVO share the estimated age curve; structural age 1 is excluded. The taller, narrower display includes model curves for years without an age-composition sample.



Points: estimated annual recruitment relative to the fitted level; bars: approximate 95% uncertainty ranges.
The dashed line marks the fitted level. The vertical axis uses a logarithmic scale.

Figure 41: Annual recruitment relative to the fitted model recruitment level in the Model 26.0. Points show exp(d_y) = R_y / R0 and bars show exp(d_y ± 1.959964 SE(d_y)) from the matching fitted covariance. R0 is the fitted recruitment-level parameter. The intervals use the fitted covariance of the recruitment deviations.



Figure 42: Observed grouped mean age for ATS, BTS and fishery with Model 26.0 predictions and the Model 26.1.2 fishery prediction. Bars are the observed mean ±2 SE, where SE = sqrt(sum[p_obs × (age score − observed mean)^2] / N), using the selected observed proportions and likelihood input N. Ages 1–14 and 15+ are scored 1–15; ATS excludes structural age 1 and the retained 2020 substitute. The same fishery observations appear once. The alternative prediction is evaluated at its fitted conditional selectivity-state modes. Bars show uncertainty in the observations.



Figure 43: Fishery observed age proportions (cohort-colored bars) and Model 26.0 fitted proportions (points joined by lines), 1964–2008. Annual panels are arranged down columns and then across, with a common proportion scale. Each panel shows its year and selected likelihood N. Age 15+ combines older cohorts.



Bars: observed proportions, colored by birth cohort; red dots and lines: fitted proportions.
Cohort colors repeat every 15 birth years. Gray bars: age 15+, a mixture of cohorts. N: selected input sample size.
The same vertical scale is used in both fishery sheets; year panels read downward, then across.

Figure 44: Fishery observed age proportions (cohort-colored bars) and Model 26.0 fitted proportions (points joined by lines), 2009–2025. The proportion scale and cohort colors match the earlier-year figure. Each panel shows its year and selected likelihood N; age 15+ combines older cohorts.



Figure 45: Bottom-trawl survey observed age proportions and Model 26.0 fitted proportions, 1982–2026. Annual panels share a proportion scale; the 2020 panel is explicitly labeled No survey. Sampled panels retain the selected likelihood Ns. Age 15+ combines older cohorts.



Figure 46: Acoustic-trawl survey observed and Model 26.0 fitted age proportions, shown on a complete 1994–2026 calendar. Empty panels mark every year without observed age-composition data, including 2020; the retained 2020 pseudo composition is excluded. Sampled panels retain their original observations, fitted proportions, likelihood Ns and common scale.



Blue filled circles: negative; red open circles: positive. Circle size represents absolute residual magnitude.
ATS uses ages 2–15+ and excludes the 2020 pseudo sample. The final age bin has no separate OSA residual.
Aggregate/Pearson displays use afscOSA rounded counts; native OSA values and assessment weights are preserved.

Figure 47: Standard afscOSA v0.0.1 diagnostics for Model 26.0, with fishery, BTS and ATS columns. Rows show aggregate observed and fitted compositions with 95% conditional observation ranges, normal-reference QQ plots with SDNR and tail summaries, OSA residual bubbles, and Pearson residual bubbles. Native OSAs and selected likelihood Ns are retained; ATS structural age 1 and the 2020 pseudo composition are excluded. Aggregate ranges condition on the fit; parameter and posterior predictive uncertainty are outside their scope.



Figure 48: ATS catchability likelihood profile for Model 26.0. Seventeen fixed catchability values span 0.45–0.65, including the fitted value 0.51694. Panel A shows the change in total negative log likelihood; the dashed line marks 1.92. Panel B shows composition data, index data and selectivity-deviate contributions; Other pools the remaining components. All other active parameters were re-estimated at each fixed q.



Figure 49: ATS-q negative log marginal density and likelihood profile for Model 26.0. Panel A compares the negative log of the MCMC marginal density, normalized to integrate to one, with the optimized profile NLL. Each curve is measured relative to its own minimum over ATS q. The marginal density uses a kernel estimate from all 5,000 saved draws; equal-grid probabilities also sum to one. Panel B pairs weighted quadratic regressions fitted directly to all 5,000 MCMC component values with profile NLL contributions for composition data, index data and selectivity deviates. The smoothers use individual draws and are displayed over the central 99% of sampled ATS q. MCMC curves are centered at the ATS-q marginal mode; profile components retain the profile minimum as their reference. These smoothers describe conditional mean native objective components; the negative log marginal density has no unique additive decomposition into those components. Other is omitted.



Figure 50: MCMC pairs plot of ATS catchability, 2026 abundance at ages 2–5, age-1 abundance in 2019 (the 2018 year class), and projected 2027 female spawning biomass for Model 26.0. Lower-triangle panels show paired values from all 5,000 saved draws with point alpha 0.3; upper-triangle panels give posterior Pearson correlations. Diagonal panels show marginal densities. Abundances are in billion fish and female SSB in million tons. The 2027 projection uses 1,375 kt catch, mean 2023–2025 fishery selectivity and within-draw mean 1978–2023 recruitment, with biological schedules fixed.

Figure 51: retrospective trajectories for Model 26.0 (left) and Model 26.1.2 with fixed process SDs (right). Black lines show each analysis's matched full 2026 reference; colors identify the nine peels ending in 2017–2025, and symbols mark their endpoints. Female spawning biomass is in million tonnes, age-1 recruitment is in billion fish, and maximum fishing mortality across ages is per year. Recruitment uses the saved age-1 calendar year rather than birth year. Panels show point estimates.



Figure 52: Historical retrospectives of beginning-of-year age-3+ EBS pollock biomass, updating Figure 62 of the 2024 SAFE. Gray lines show archived 2007–2023 assessments; orange shows the official 2024 SAFE Table 23 estimates through 2024; blue shows Model 26.0 estimates through 2026. Biomass sums beginning-of-year numbers at ages 3–15+ multiplied by the corresponding annual fishery weights at age, using the Model 26.0 fit. Historical series retain their original definitions and available endpoints, which may include next-year estimates. The 2024 tabulated values are rounded. No new assessment fit was produced in 2025. These are point estimates from successive assessments, encompassing changes in data, methods and biological inputs.



Figure 53: Selected annual composition sample sizes and their Francis diagnostic equivalents for the Model 26.0 fishery, BTS and ATS groups. This is a diagnostic comparison; no Francis reweighting is applied to the assessment inputs.



Figure 54: Standardized mean-age residuals used in the Francis diagnostic comparison for Model 26.0, with the selected input sample sizes. Positive values indicate older observed than predicted mean age. The original selected-data treatment and diagnostic formulas are retained.



Figure 55: Annual fishery selectivity from Models 26.0 and 26.1.0. The split configuration separates Foreign and Modern domestic fisheries. Curves use the normalization recorded with the verified figure data.



Figure 56: Fishery selectivity before and after the fleet split, with signed 1991-minus-1990 differences by age for Models 26.0 and 26.1.0. The figure preserves the existing boundary-year and normalization definitions.



Figure 57: Model 26.0 fishery age-composition fits. We displayed the Foreign period (1964–1990, left) and Modern domestic period (1991–2025, right) separately for comparison with Model 26.1.2, although Model 26.0 fitted one fishery. Row A compared sample-weighted observed age proportions (bars) with fitted proportions (black lines). Row B compared observed annual mean ages and approximate 95% sampling intervals with fitted mean ages. Row C showed observed minus fitted proportions in percentage points; open red circles indicated observed proportions above the fit, and filled blue circles indicated proportions below it. Bubble areas represented difference magnitudes on a scale shared with the Model 26.1.2 figure. Mean ages treated the 15+ group as age 15.



Figure 58: Model 26.1.2 fishery age-composition fits, using the fixed-process-SD conditional fit. Columns showed the Foreign and Modern domestic fisheries. Row A compared sample-weighted observed and fitted age proportions; row B compared observed annual mean ages and approximate 95% sampling intervals with fitted mean ages; row C showed observed minus fitted proportions in percentage points. The figure used the same observations, likelihood weights, age ranges and bubble scale as the Model 26.0 figure. The mean-age intervals described observation sampling uncertainty, and the fitted curves showed point estimates. Mean ages treated the 15+ group as age 15.



Figure 59: Conditional simulation-retrospective revisions for Model 26.0. Columns show ten datasets generated at the fitted parameters (left) and ten generated from regularized posterior parameter draws (right). Rows show female spawning biomass and age-1 recruitment. Blue points identify individual simulated datasets, blue lines join their medians, and black diamonds and lines show observed-data revisions. For a peel ending in year y, revision is 100 × [peel estimate in year y / its own full 2026 fit estimate for year y − 1]. Both columns use the same vertical scale within a quantity.

Figure 60: Native SparseNUTS pairs diagnostic for the six parameters with the lowest bulk effective sample size in the newly fitted regularized Model 26.0 posterior. Lower panels show all 5,000 retained draws in black, the saved regularized native MAP as a red dot, and the 95% bivariate Normal approximation from the matching full inverse augmented Hessian as a red ellipse. Upper panels report pooled posterior Pearson correlations. Diagonal panels show four retained chain traces, bulk ESS and R-hat. Parameters retain their native coordinate scales. SparseNUTS names the reference field mle; here it contains the regularized MAP, separate from the production penalized-ML fit.



Figure 61: Matching native-coordinate local Hessian SE and posterior SD for all 1,266 active parameters under the same approved regularized target. Both axes use logarithmic scales; the dashed line denotes equality. Rings identify the six slowest parameters, with posterior-SD/local-SE ratios 1.28–1.52. Coordinate units differ among parameters. MCMC standard errors quantify simulation precision and are separate from the parameter uncertainties shown here.



Figure 62: MCMC posterior histograms for the six native coordinates with the lowest bulk ESS in the original chains. All 5,000 saved parameter vectors contribute. Bar heights show probability per bin; the panels retain parameter units. The identity of these coordinates follows the original-chain diagnostic.



Figure 63: Index checks compare observed survey and historical catch-rate indices with simulated observations generated from 1,000 MCMC draws. Each draw supplies a model expectation; random observation errors are then added using the same error assumptions used to fit each index, including the relationships among BTS errors. Lines show median expectations and shading covers the middle 95% of simulated observations at each year. Circles are observed indices and triangles identify existing substitutes. Some simulated values are negative because the fitted Normal error distributions allow them. The BTS overall mismatch measure exceeds its observed value in 0.2% of simulations, indicating that the model rarely reproduces this feature of the BTS data. Units differ among panels. Biological inputs remain fixed.



Figure 64: Age-composition checks compare the latest three observed samples from each data source with simulated samples generated from 1,000 MCMC draws. For each draw, fish are randomly assigned to age groups using the model-predicted age proportions and the sample size used in fitting, rounded to a whole number. Lines show median model proportions, shading covers the middle 95% of simulated sample proportions at each age, and points show observed proportions. The shading includes parameter uncertainty and the assumed sampling variation. This simulation approximates the age-composition fitting formula, which adds a small constant to age proportions when calculating fit. The 2020 ATS substitute is excluded. Age 15 includes all older fish. Age conversions, weights and other biology remain fixed.



Figure 65: Catch projections for Model 26.0 under Alternative 2 (expected catch, top) and Alternative 3 (recent mean F, bottom). Black lines show simulation means, ribbons show the 5th–95th percentiles from 1,000 future-recruitment simulations, and thin colored lines show ten matched trajectories. Colors identify the same recruitment realization in both alternatives and the spawning-biomass figure. Alternative 2 fixes catch at 1,375 kt in 2027 and 2028, then applies the maximum-ABC rule; its paths coincide in the prescribed-catch years. The ribbons represent future recruitment variation; the advice tables retain historical-mean recruitment.



Figure 66: Female spawning-biomass projections for Model 26.0 under Alternative 2 (top) and Alternative 3 (bottom). Black lines show simulation means, ribbons show the 5th–95th percentiles from 1,000 future-recruitment simulations, and thin colored lines show the same ten matched trajectories as the catch figure. Future recruitment follows an inverse-Gaussian distribution fitted to the 1978–2023 age-1 entries. These ranges include future recruitment variation only. Dashed B40% and dotted B35% lines mark the primary projection reference points.



Figure 67: Historical trajectory of Model 26.0 female spawning biomass (B) and maximum fishing mortality at age (F), 1981–2026, with deterministic Alternative 2 projections for 2027–2028. Ratios use the current primary Tier 3 reference points, B35% = 2.182 million t and F35% = 0.688 per year, held constant across years. Lines connect consecutive years; two-digit labels identify years, and boxed labels with yellow shading identify projections. Solid lines mark ratios of one; the dashed vertical line marks half B35%. Projected catch is 1.375 million t in each year.



Ages 9–15+ appear in the full annual-curve figure. 2026 is a model-inferred terminal pattern.

Figure 68: Recent normalized fishery selectivity curves from the Model 26.0 and the proposed raw-curve averaging reference. Half-maximum crossings and curve-shape summaries use the existing verified definitions.



Figure 69: Annual Model 26.0 fishery selectivity curves for 2007–2026, each compared with the proposed 2023–2025 raw-curve average rescaled to maximum one. The inferred 2026 curve retains its distinction from directly informed earlier-year curves.



Figure 70: Changes in maximum ABC and OFL under the annual future-selectivity curves relative to the matching Model 26.0 baseline. Rows retain the recorded 2027 maximum-ABC ordering. These are projection selectivity sensitivities conditional on the same fitted source and projection design.



Figure 71: Annual younger-age selectivity measures for 2007–2026 from the Model 26.0. The inferred 2026 point is distinguished. Curve-weighted mean age is a shape measure under equal age weights, distinct from a population-weighted selected age.



*2026 is an inferred terminal curve. Dashed line and diamond: current 2023–2025 average.
Grey path follows years; point labels identify the selectivity year.

Figure 72: Annual Model 26.0 fishery selectivity and fully selected F35%, a proxy for FMSY, for 2001–2026. Mean selected age uses ages 1–10; SPR uses ages 1–14 and 15+, fixed 2026 mortality, maturity and female spawning weights, and quarter-year spawning timing. Curves are normalized to maximum one. The 2023–2025 mean-curve reference and inferred 2026 curve remain separately identified. The historical 2024 figure used FMSY, a different quantity.

Appendices

Appendix A
SSC and Plan Team Requests and Responses

The Plan Team requests and assessment responses are summarized in Table A1.

Table A1 *Responses to SSC and Plan Team Requests*

SSC and Plan Team requests			
ID	Recommendation	Response	Response status
PT-01	Bring Model 26.0 forward for management use in November and present Model 23.0 for continuity and comparison during the transition to Rceattle.	Model 26.0 estimates, uncertainty, diagnostics, Tier 3 projections and Model 23.0 comparisons are presented. Regularized posterior summaries, predictive checks and observed and simulated-data retrospectives are included. Model 26.1.2 is a sensitivity comparison conditional on fixed Foreign and Modern process SDs of 0.1235 and 0.582. Risk categories are Level 1.	Assessment results and sensitivity comparisons presented
PT-02	Evaluate alternative treatments of age-1 BTS selectivity, including the selectivity penalty and a time-invariant alternative, to assess their influence on the retrospective pattern.	The author proposes no additional age-1 BTS selectivity sensitivities for this assessment. Observed and simulated-data retrospective results are presented; these results do not separately identify the effect of an alternative age-1 selectivity treatment.	No additional age-1 selectivity tests proposed
PT-03	Extend the Model 26.0 self-test with retrospective peels on simulated datasets generated from the fitted model, and examine retrospective patterns in time-varying selectivity estimates.	Two sets of ten simulated datasets use the fitted parameters and ten regularized posterior draws, respectively. Each dataset has a full-2026 fit and nine peels ending in 2017–2025. Observed spawning-biomass rho (+0.121) exceeds every simulated value in both sets. Posterior diagnostics and predictive checks are also presented. The simulations assess population revisions; they establish no single retrospective cause.	Paired simulation retrospectives and posterior predictive checks presented
PT-04	Present retrospective results from Models 23.0 and 26.0 side by side in November.	Matched retrospective runs for Models 23.0 and 26.0 were completed and compared side by side. Both showed similar spawning-biomass retrospective patterns, with nine-peel Mohn's rho of +0.156 and +0.143, respectively (positive values indicate higher estimates in the shorter fits). See the side-by-side comparison and numerical summaries.	Completed; requested comparison provided
PT-05	Further examine the influence of BTS index volatility and other potential data or ecosystem-related causes on the retrospective pattern.	The author proposes no additional BTS data-influence tests for this assessment. Model and cumulative-data comparisons, observed and simulated-data retrospectives, and posterior predictive checks are presented. The BTS mismatch measure is as large as observed in 0.2% of predictive simulations. These results identify model-data mismatch but establish no ecosystem cause.	Diagnostics presented; no additional BTS influence tests proposed

SSC and Plan Team requests			
ID	Recommendation	Response	Response status
PT-06	Clearly document age-composition weighting in Model 26.0 and prioritize further evaluation of data weighting given patterns in the OSA diagnostics.	Age-composition information weights and likelihood multipliers are documented by data source and year. Model 26.0 and fixed-SD conditional Model 26.1.2 OSA and mean-age comparisons retain the selected sample sizes. Model 26.1.2 residuals condition on fitted selectivity states and exclude parameter uncertainty. Diagnostic equivalent weights are not applied as new inputs. Alternative-weight fits are outside the scope of this comparison.	Weighting and conditional diagnostics presented; alternative-weight fits remain separate
PT-07	Continue development of material that could contribute to a future EBS pollock Ecosystem and Socioeconomic Profile, as resources allow.	A draft was developed in September and further work is included here following recommended format and guidance.	Draft has been developed and is included as Appendix B.

Note. Responses to the September 2026 BSAI Groundfish Plan Team recommendations. Status distinguishes completed work, decisions about further tests and development of the ESP draft.

Earlier SSC comments and response plan

The matrix below preserves the October 2025 SSC response plan as reproduced in the September 2026 development source. It records proposed actions at that stage; completion and outstanding work are identified in Section 1.4 and the response registers. Its RTMB and Model 25.0 entries are historical planning context rather than the software selection. The September 2026 Plan Team direction establishes Model 26.0 and Model 23.0 as the November main and comparison models; the RTMB and SPoRC work remains part of the development record.

The earlier SSC comments and response plan are reproduced in Table A2.

Table A2 *Earlier SSC Comments and Response Plan*

SSC comment	Response
Use the 2024 base model as the source for an RTMB implementation, with routine data updates.	Use the accepted Model 23.0 (ADMB) configuration as the scientific source. First check the direct RTMB translation as Model 23.1, then use Model 25.0 (RTMB) for the assessment line begun in 2025. List each new data set and each model change separately. Save the model version, run files, diagnostics, projections, and management quantities used in the report.
Use the updated Groundfish Assessment Program estimates of survey biomass and age composition.	Add the sdmTMB biomass estimates and tinyVAST age compositions as revised survey data. Record when the estimates were received, the software version, model settings, and input files. Compare the old and new survey series and show whether the change affects the assessment results.
Use FT-NIRS age compositions for the 2025 fishery and surveys.	Document how the FT-NIRS ages were produced, screened, grouped, and assigned to fishery and survey data. Record sample sizes and aging-error matrices. Check the final model input against the source data before fitting the assessment.
Compare traditional microscope ages with FT-NIRS ages and show the effects on model results and management quantities.	Run the same assessment with traditional ages, FT-NIRS ages with one aging-error matrix, and FT-NIRS ages with fleet-specific aging-error matrices. Compare cohort patterns, fits to fishery and survey ages, residuals, recruitment, spawning biomass, reference points, ABC, and OFL. Save the full comparison in the assessment appendix and at a permanent web address.
Revisit the revised AVO index after the new uncertainty analysis has been reviewed.	After review of the final uncertainty method, compare the revised point estimates and uncertainty with the AVO series. Show their effect on model fit, stock estimates, and management quantities before proposing a change to the base model.
Develop a plan with the BSAI Groundfish Plan Team for moving beyond ADMB.	Use the model history and numbering plan above. Compare RTMB, Rceattle, and SPoRC with the same data and a common set of tests. Ask the Plan Team to agree on the tests before selecting the model software for future assessments. Keep the accepted assessment model as the control until a replacement gives stable estimates, sound uncertainty, acceptable retrospective results, and reproducible projections.

SSC comment	Response
Examine the effect of including Russian catches.	First confirm the source, area, years, units, species identification, and uncertainty of the Russian catch estimates. Then run clearly defined catch scenarios. Explain the assumptions about stock movement and population boundaries and show the effects on stock size, reference points, ABC, and OFL.
Explore time-varying natural mortality, selectivity, age-composition weighting, and cohort and temperature effects on weight at age.	Study each topic separately. State the scientific question before fitting a model. For each change, show model fit, parameter uncertainty, retrospective results, projections, and effects on management advice. Keep poorly estimated or unstable models as research results.
Continue work with CEATTLE estimates of natural mortality and consider Rceattle for time-varying mortality.	Test time-varying natural mortality as a sensitivity analysis. Check whether the model can separate mortality changes from recruitment, selectivity, and survey effects. The full set of assessment needs will guide the choice of model software.
Explain how the recruitment penalty affects the estimated 2018 year class and why the likelihood profile favored a larger recruitment estimate.	Prepare a likelihood profile for the 2018 recruitment deviation. Show how the recruitment penalty and each data source change across the profile. Explain which parts of the model favor a larger year class and how the estimate affects spawning biomass, reference points, ABC, and OFL.
Clearly identify the models proposed for the assessment, including model numbers.	Use Model 23.0 (ADMB) for the accepted 2024 source, Model 23.1 (RTMB) for the direct translation check, Model 25.0 (RTMB) for the line begun in 2025, and Model 26.0 (Rceattle) for the line begun in 2026. Use the same short names in the report, presentations, run folders, tables, and archive. For every model, list its purpose, parent model, data set, model changes, software version, and review status. State clearly which model is recommended for management advice.
Archive the October and December SAFE documents and presentations.	Archive the October 2025 material that exists and add a short note explaining that the funding lapse prevented the November and December SAFE package. For future reviews, save the Quarto source, rendered reports, presentations, model files, data list, software versions, and file checksums in one dated release.

Note. Responses to the October 2025 SSC comments on EBS pollock. The SSC comments are summarized from pages 27 to 29 of the final October 2025 SSC report.

Appendix B
Ecosystem and Socioeconomic Profile of Eastern Bering Sea Walleye Pollock

This draft Ecosystem and Socioeconomic Profile (ESP) for EBS pollock brings together substantial work completed to date and provides a starting point for future ESP Team discussions. An AI agent helped compile and coordinate material from disparate information sources and research activities. Over the next year, the EBS pollock ESP Team will formalize its membership and begin meetings that engage a broader group of scientific experts to develop the profile. Please contact Jim Ianelli and Kalei Shotwell with questions or for further details about this document or the EBS pollock ESP.

Contents

1. Executive Summary
2. Responses to SSC and Plan Team Comments on ESPs in General
3. Responses to SSC and Plan Team Comments Specific to EBS Pollock
4. Introduction
5. Indicator Synthesis
6. Indicator Assessment
7. Data Gaps and Future Research Priorities
8. Conclusion
9. Literature Cited
10. Tables
11. Figures

Executive Summary

The Ecosystem and Socioeconomic Profile (ESP) connects stock-specific environmental, biological, fishery, and community information with assessment interpretation and management questions. Its sequence of focus, synthesis, analysis, and communication provides a structured way to evaluate proposed indicators (Shotwell et al. 2023). This draft focuses initially on recruitment to age 1, then identifies processes affecting juvenile survival to age 3, adult growth and condition, survey interpretation, and fishery performance.

The ESP distinguishes two DSEM applications. The archived Rceattle pilot evaluates birth-year SST and age-1-year cold-pool proxies (Ianelli et al. 2026). A separate age-1 DSEM working paper analyzes 18 ecosystem series alongside an age-1 response estimated by the assessment. These are recorded repository inputs with incomplete primary-source metadata. Regional age-0 energy density provides an additional measured-condition series (Page and Maselko 2026). Interpreting this series requires its sampling season and regional coverage.

Acceptable Biological Catch Considerations

Candidate Predictive Indicators

Fall age-0 total energy, prey quality and abundance, and juvenile–predator overlap are priorities for testing. Field studies connect fall total energy with age-1 recruits per spawner, and laboratory experiments identify body size and energy reserves as determinants of survival during food deprivation (Heintz et al. 2013; Sogard and Olla 2000). Studies of pollock recruitment and spatial predation identify juvenile overlap with adult pollock and arrowtooth flounder as another important pathway (Mueter et al. 2006; Spencer et al. 2016). These studies support biological hypotheses; performance in withheld cohorts will determine whether the resulting annual indicators improve assessment forecasts.

Age-1 response observations in the working-paper input end in 2024. Its 2025 and 2026 values are full-data conditional missing-year estimates, rather than observations or genuine forecasts. Rolling validation over seven observed age-1 years, 2018–2024, gives RMSE 1.924 for the configured DSEM versus 1.847 for AR1 in the working response units. The configured DSEM scores worse than the AR1 reference in this test. The historical Rceattle pilot has a different response, timing and scaling; its in-sample comparisons remain separate from that forecast test.

The regional energy-density series contain 21 southeastern Bering Sea (SEBS) and 15 northern Bering Sea (NBS) annual means within 2002–2025, measured in kJ/g dry mass. The latest values, for 2025, are 19.23 and 20.84 kJ/g dry mass for SEBS and NBS, respectively. Both are lower than their 2024 observations; a 2026 energy-density observation is absent. Sampling metadata, observation uncertainty, and forecast evaluation remain prerequisites for a predictive role (Page and Maselko 2026).

Contextual Indicators

Sea ice, seasonal temperature, cold-pool extent, plankton and euphausiids, age-0 abundance and condition, adult condition, weight at age, and predator overlap form the initial contextual set. The relevant biological question is whether each cohort encounters adequate prey, accumulates energy before winter, and survives subsequent predation. The poor 2007 cohort illustrates why a favorable temperature category alone can give an incomplete account: low summer production and poor age-0 condition occurred during a cold year (Gann et al. 2016).

The recorded 2026 physical and prey indicators show a mixed pattern relative to their own observed-period means. Sea ice and cold-pool extent are above those means; SST and bottom temperature are below. The open-water bloom indicator is lower, while chlorophyll a is higher. Spring small and large copepods and summer small copepods are below their respective means; summer large copepods are slightly above. These descriptive comparisons use variable-specific periods and gaps, as detailed under Recorded Annual Indicator Series. Metadata and approval checks remain necessary before ecological status categories are assigned.

Fishery Informed Indicators

Candidate indicators include age and size composition of catch, fishery weight at age, season and area of harvest, standardized fishery catch rates, catch and quota utilization, and salmon bycatch measures. Their interpretation must account for fishing choices, allocation and avoidance behavior, processing demand, and reporting coverage. These indicators may reveal operational changes or sampling differences that help interpret assessment inputs. Their proposed inclusion in this ESP requires a defined management question and documented denominators.

The draft supports evaluation of ecosystem and fishery concerns in the SAFE risk table. Any resulting change in risk classification or ABC must be documented by the assessment authors using the completed annual synthesis and assessment results. ABC and OFL remain unchanged by this contribution.

Total Allowable Catch Considerations

Verified economic context covers BSAI catch, participation, ex-vessel and first-wholesale statistics through 2024, and U.S.-national trade through 2025 (Catch and Market Context). These scopes differ from the BS assessment stock and from each other. A **speculative 2026 interpretation** is that firmer product unit values could support gross returns, while operating costs and product mix could offset those gains unevenly among vessels, processors and communities. 2026 margins and community outcomes remain uncertain. This interpretation is qualitative.

Responses to SSC and Plan Team Comments on ESPs in General

The template establishes useful review questions: distinguish predictive from contextual indicators, evaluate competing mechanistic explanations, document DSEM diagnostics, and coordinate economic and community information with existing Council products. This pollock contribution addresses those questions by defining indicator roles, documenting the fitted pilot and its diagnostic limitation, and proposing an explicit socioeconomic inventory. These are drafting choices informed by the template. Formal responses require the exact relevant comment, meeting date, source page, analysis, and review disposition for the EBS pollock submission.

Responses to SSC and Plan Team Comments Specific to EBS Pollock

December 2024 SSC, EBS Ecosystem Status Report (page 3): the SSC recommended pollock-focused DSEM work to reconcile contrasting larval, juvenile-condition, and adult signals, and to investigate changed recruitment relationships (North Pacific Fishery Management Council Scientific and Statistical Committee 2024). This recommendation supports exploration; acceptance of a particular model or assessment application requires subsequent review.

Contribution to that work: Elizabeth Siddon's 3 September 2026 coordination email describes an exploratory age-1 survival analysis for the 2026 EBS ESR, drawing on process studies, the January 2025 pollock workshop, and Ecosystem REFRESH/subject-matter-expert input (Siddon 2026). Johanna Page's reply supplies annual energy density and requests the DAG and DSEM for contributor review. This revision documents the data, presents the regional series, and specifies a candidate cohort-aligned pathway. Fitting the expanded DSEM and reviewing its results remain the next analytical steps. The ESR analysis and the broader SAFE ESP share this development effort while retaining their own review processes.

The assessment website records continuing work on contrasting early-life observations, cannibalism and biomass consumed, multispecies mortality, weight at age and condition, and stock distribution across U.S. and Russian waters (EBS Pollock Assessment Team 2026). Its retained earlier responses carry an explicit requirement for verification before use as final results. This draft translates those topics into proposed ESP tasks.

Pollock review topics and proposed ESP contributions are summarized in Table B1.

Introduction

Pollock recruitment reflects survival through several life stages with different food requirements, habitats, and predator exposure. Large differences in egg or larval abundance can be modified by later feeding, growth, transport, and mortality. The first-year review by Duffy-Anderson et al. provides the organizing synthesis for these processes (Duffy-Anderson et al. 2016). The local recruitment review extends that structure through ages 1 and 2 to survival at age 3. In this appendix, **age-1 recruitment** is the focal response used by the DSEM pilot; **survival to age 3** is a separate later-stage outcome that retains the identity of the same birth cohort.

Justification

The ESP creates an evidence trail from a biological or socioeconomic question to an indicator and its proposed use. For EBS pollock, the immediate question is whether early-life information can improve interpretation of uncertain recruitment and future biomass. Additional questions concern adult productivity, changing spatial availability to surveys, fishery operations, and community exposure. The September 2026 document provides the starting framework for developing this appendix (Ianelli et al. 2026).

The September 2026 development document recommends Model 26.0 for review in November and December 2026 and retains Model 23.0 for continuity while the review bodies consider that transition. Its operational specification keeps DSEM disabled, with environmental pathways evaluated as research. This ESP contribution retains that review status. Selection of an assessment model and acceptance of an environmental forecasting component are separate decisions.

Data

Each annual series needs a source, contributor, definition, units, season, spatial domain, observation years, uncertainty, revision history, and update schedule. Survey and assessment products require additional checks because several indicators may derive from the same samples. For example, assessment recruitment, survey age composition, and a juvenile index may contain overlapping evidence. A model that uses them together must account for that dependence.

The pilot's archived standardized covariates contain 42 cold-pool observations and 41 SST observations in recruitment-year rows ending in 2024. The broader local cohort table contains 15 prey-index values and retains empty age-0 energy, ice-retreat, and juvenile–predator overlap fields. The energy-density workbook provides a separate annual energy-density product; integrating it requires explicit regional and cohort definitions while preserving the historical fit archive.

Age-0 Energy Density

Annual mean energy density of age-0 walleye pollock is available for the northern and southeastern Bering Sea during 2002–2025, in **kJ/g dry mass** (Page and Maselko 2026). Johanna Page (Vollenweider) provided the data, with Jacek Maselko as a co-contact. The southeastern series contains 21 annual means and the northern series contains 15. Both regions lack observations for 2003, 2013 and 2020; the northern series also lacks 2008–2012 and 2014. Missing years remain gaps, and values retain the precision reported by the contributors.

The source describes annual means. Sampling season, exact regional boundaries, annual weighting, sample sizes, measurement consistency, and uncertainty require contributor confirmation. These definitions will determine whether the observations represent pre-winter condition and which region matches the recruitment response. Energy density measures energy per unit dry mass. Estimating total energy per fish requires compatible individual dry mass and aggregation methods. Regional means remain separate, and 2026 energy-density observations are pending.

The source inventory and remaining data-assembly needs are summarized in Table B2.

The 2026 NOAA temperature page reports an EBS survey domain, whereas the 2025 survey included both eastern and northern Bering Sea areas (Alaska Fisheries Science Center 2026a). Comparisons of area occupied, centers of distribution, temperature exposure, and predator overlap should therefore use a common domain or explicitly describe differences in coverage.

Indicator Synthesis

Ecosystem Processes

Spawning Supply and Transport

Spawning biomass and spawning location influence egg and early-larval supply. Smart et al. found associations between spawning biomass and early stages, with weaker correspondence after early larval development (Smart et al. 2012). Transport and the spatial arrangement of spawning also matter. The biophysical model of Petrik et al. reproduced warm–cold distribution differences most successfully through changes in spawning areas (Petrik et al. 2015). These results support separate hypotheses for cohort supply, transport, and access to nursery habitat.

Feeding and Fall Age-0 Condition

Prey availability includes amount, nutritional quality, and spatial match with fish. Larval copepod concentrations and frontal habitat illustrate why a broad regional biomass index can differ from prey encountered by an individual fish (Smart et al. 2012; Brodeur et al. 2000). Diet studies show that prey use varies with fish size and that pollock and Pacific cod can partition prey (Buckley et al. 2016; Strasburger et al. 2014).

Siddon et al. describe a seasonal shift from larval growth and development toward lipid storage after transformation to the juvenile stage (Siddon et al. 2013). Heintz et al. found higher diet lipid content and age-0 energy density during cold conditions in their study; fall total energy was positively associated with age-1 recruits per spawner (Heintz et al. 2013).

Moss et al. also documented differences in age-0 energetic condition and food habits across thermal conditions (Moss et al. 2009). These findings motivate indicators of energy density, body mass, and total energy with explicit measurement definitions. The annual series measures energy density in kJ/g dry mass. Total energy in kJ per fish additionally depends on fish dry mass and appropriate individual-level aggregation.

Eisner et al. evaluated large copepods as leading indicators of age-3 pollock abundance for the 2002–2015 year classes, excluding 2013 (Eisner et al. 2020). Their comparison reinforces the value of prey information while identifying a specific cohort range and recruitment age. Translation into an age-1 forecasting component requires testing the corresponding response and lag.

Overwinter Survival

Fall condition provides the starting reserves for the first winter. Sogard and Olla's food-deprivation experiments show that greater size, higher initial condition, and colder water can extend survival (Sogard and Olla 2000). Field survival also involves winter feeding, movement, and predators. Laboratory results therefore support a mechanism and its direction, while annual field data are needed to quantify its effect on recruitment.

Sea ice, bloom timing, summer mixing, and temperature influence prey production and energy demand. Sigler et al. connect annual production of copepods, euphausiids, and juvenile pollock with seasonal environmental conditions (Sigler et al. 2016). Gann et al.'s analysis of 2007 identifies a plausible production limitation during a cold year (Gann et al. 2016). The ESP should evaluate these intermediate processes before interpreting a simple warm–cold contrast as a cohort forecast.

Predation and Later Juvenile Survival

Adult pollock can increase juvenile mortality through cannibalism, while arrowtooth flounder and other predators create additional exposure. Mueter et al. identify both food-related and predation-related controls on pollock recruitment, including juvenile–adult overlap (Mueter et al. 2006). Spencer et al. model spatial predation by arrowtooth flounder and show how cold-pool extent can alter predator access to juvenile habitat (Spencer et al. 2016). Predator abundance, spatial overlap, diet, biomass consumed, and predation mortality describe successive parts of this process and require separate definitions.

The timing of predation matters for the response. A summer overlap or cold-pool observation in year $t+1$ occurs after the beginning of the cohort's age-1 year. Its use to explain an age-1 entry state requires explicit justification or a revised response definition. Such observations may be more directly aligned with survival during age 1 and later ages. The pilot's same-year cold-pool edge is retained as a research proxy, with this timing question identified for review.

Adult Productivity and Observation Processes

Adult growth, condition, maturity, distribution, and consumption connect recruitment to biomass production and future spawning output. Weight at age and condition should retain age, sex where available, season, gear, and location. Survey distribution indices also help evaluate changing availability to sampling. Changes in survey footprint or in the transformation of observations into annual estimates require documented comparison before they become evidence of a biological change.

Life stages, processes and priority observations are summarized in Table B3.

Workshop Conceptual Models for Recruitment and Growth

The January 2025 pollock biology and ecology workshop developed preliminary conceptual models for Bering Sea recruitment and growth (Monnahan et al. 2026). Figure B1 organizes reproductive output, seasonal juvenile stages, prey quality, energetic condition, predation, and cold-pool extent. Figure B2 connects population weight at age with somatic growth, size-selective mortality, prey, competitors, fishing, and physical conditions. Together they support an ESP that examines recruitment and growth. Their pathways are workshop hypotheses; annual indicator selection and estimation of pathway effects require subsequent evaluation.

Recruitment pathways: spawning biomass and fecundity inform reproductive output. Seasonal surveys track the cohort from spring age 0 through age 2. Diet, krill, copepods, and prey lipid measurements inform forage quality and juvenile

energetic condition; predation indices and cold-pool extent inform later survival. The diagram highlights both candidate observations and intermediate biological states.

Growth pathways: the model separates growth of surviving fish from changes in population weight at age associated with size-selective mortality. Prey availability, density, genetic diversity, and spatial history enter the growth network alongside physical conditions and zooplankton. This distinction supports age- and season-specific weight and condition indicators with explicit survey or fishery sampling definitions.

The cohort sequence in Figure B3 summarizes the recruitment mechanisms developed in this draft and extends the survival question to age 3. The workshop diagrams provide a broader set of hypotheses for future indicator development.

Socioeconomic Processes

Biological supply, access to fishing grounds, operating costs, product demand, and processing capacity can jointly affect realized harvest and its value. Those processes may alter where and when vessels fish, the size and age of fish landed, and the distribution of benefits. This component should therefore connect each indicator to a concrete question, such as whether lower quota utilization reflects reduced fish availability, market conditions, processing limits, or avoidance behavior.

The Economic SAFE provides source categories for catch, discards, prohibited species catch, production and value, participation, and effort (Alaska Fisheries Science Center 2026b). This draft proposes species- and sector-specific summaries where the source supports that detail. Community analysis requires clear distinctions among landing ports, vessel ownership, workforce residence, processor location, and CDQ beneficiaries. Each describes a different form of participation or exposure.

Proposed Indicators

The mechanism table (Table B4) links a driver, a measurable process, and the intended response. Expected directions are hypotheses for the stated pathway. Several upstream physical effects depend on season, prey response, and spatial coverage, so their net sign can change across conditions.

Indicator Assessment

Indicator Selection

Selection starts with a mechanistic explanation, the population or socioeconomic response, and the date when the information becomes available. Candidates then require reliable annual production, adequate spatial and temporal coverage, quantified uncertainty, and additional information beyond the other indicators.

Use four working roles: **candidate predictive**, for indicators under forecast evaluation; **contextual**, for interpreting ecosystem or assessment conditions; **fishery informed**, for operational observations that require fishing-behavior interpretation; and **socioeconomic monitoring**, for economic and community questions. Promotion to a predictive role requires reliable performance using information that would have been available at the historical forecast date. This is the proposed working classification for this appendix; the ESP team should reconcile its labels with review guidance.

Ecosystem Indicators

The initial set should remain small enough for scientific review: seasonal physical forcing; a documented prey index; fall age-0 size and energetic condition; juvenile–predator overlap or a reviewed mortality product; and adult growth and condition. Larval supply and transport add value when annual coverage and biological alignment support them. Indicator families should be assessed together so correlated temperature, ice, and prey signals receive an interpretation that accounts for shared processes.

The available age-0 energy-density series describe regional energetic condition (Figure B4). The recorded histories of the ecosystem inputs and age-1 response provide context for the DSEM analysis (Figure B5).

Socioeconomic Indicators

Proposed socioeconomic indicator families, measures and management questions are summarized in Table B5.

Catch and Market Context

The broader Bering Sea and Aleutian Islands (BSAI) pollock fishery reported 1.3163 million tons of total catch in 2024 (Table B6). Trawl catcher vessels retained 678.40 thousand tons, with an ex-vessel value of $213.43 million. These economic statistics cover the BSAI; their area and accounting definitions differ from the BS stock-assessment catch series. Published pollock-target vessel counts were 103 across harvesting sectors and 76 catcher vessels. The counts describe participation in the target fishery and include their source-specific gear definitions.

Published first-wholesale pollock production was 487.89 thousand tons of product weight and $1.4743 billion in 2024 (Table B7). The Economic SAFE excludes confidential components from some aggregates; the combined fillet quantities and derived values for 2021–2022 remain unavailable here. Product mix and market conditions both affect the observed unit values (Abelman et al. 2026).

Global Alaska pollock capture production was 3.573 million tons in 2024 (Table B8, panel A). The United States and Russian Federation accounted for 39.9% and 54.0%, respectively, and BSAI catch represented about 36.8% of the global total. U.S. Alaska-pollock domestic exports totaled 333 thousand tons of products valued at $1.00 billion in 2025, compared with 332 thousand tons and $0.946 billion in 2024 (panel B). These national trade statistics extend through 2025; global catch statistics extend through 2024. Their product weights, nominal export values and legacy import/destination baskets are defined in the table notes (FAO 2026; NOAA Fisheries 2026).

Among the three destination groups shown, selected European countries accounted for the largest export shares throughout 2020–2025 (Figure B6). Meat's share of export volume rose from 45.8% in 2020 to 53.3% in 2025, while its value share rose from 44.0% to 53.0% (Figure B7). The Russian Federation accounted for more than half of global catch in each year from 2020 to 2024 (Figure B8). First-wholesale surimi volume declined from 185.5 to 165.1 thousand tons between 2023 and 2024, while fillet volume rose from 142.9 to 158.6 thousand tons (Figure B9).

Annual Indicator Status

For each retained indicator, present the latest observation, historical baseline, trend, uncertainty, missing years, and mechanism-based interpretation. Define thresholds before assigning status categories. A high value describes an observation relative to its baseline; its biological meaning depends on the pathway. Missing observations should receive a missing-data label.

The completed annual status tables should report five recent observation years where the series supports them, with 2026 provisional data clearly marked. Comparisons of year-to-date catch, effort, value, or bycatch should use the same cutoff date in each comparison year. Data revisions and changes in survey coverage should accompany the plotted series.

Age-0 Energy Density Through 2025

The latest five available observation years are 2021–2025 (Table B9). In 2025, the SEBS annual mean was 19.23 kJ/g dry mass, 0.64 lower than in 2024. The NBS annual mean was 20.84, 0.09 lower than in 2024 (Page and Maselko 2026). These are descriptive differences between annual means. Their sampling uncertainty and biological status classification await contributor metadata and agreed baselines. The latest observation year remains 2025.

The plotted energy-density series preserves gaps and regional differences (Figure B4). Every plotted point corresponds to an annual mean. The series contains annual means without sampling-error estimates; regional pooling, interpolation, and a 2026 extrapolation remain outside this summary.

Recorded Annual Indicator Series

The age-1 DSEM input spans 1964–2026 and contains 18 ecosystem series plus the assessment-derived age-1 response (Figure B5). Seventeen ecosystem series have recorded 2026 values; the energy-density series ends in 2025, and age-1 observations end in 2024. These are working repository inputs, with primary-source attribution, units, seasons, spatial domains and transformations still partly unverified. Energy density has an independently corroborated SEBS identity and kJ/g dry-mass units; NBS remains a separate series.

Standardized departures below equal each recorded value minus that series' mean, divided by its sample standard deviation. Each baseline uses all available finite observations, including the latest recorded year, and retains the series'

gaps. These baselines are descriptive historical variability, rather than a common fixed climatology, sampling-error intervals or approved biological thresholds.

In 2026, recorded sea ice is +0.89 SD relative to its 1985–2026 baseline of 42 observations. Cold-pool extent is +1.12 SD, SST −0.39 SD and bottom temperature −0.79 SD relative to their separate 1982–2026 baselines of 44 available observations each. Sea ice and cold pool both exceed their recorded 2025 values. SST also increased from 2025 despite remaining below its observed-period mean, whereas bottom temperature fell.

The open-water bloom indicator is −1.54 SD and chlorophyll a is +0.77 SD relative to their separate 1998–2026 baselines of 29 observations each. Both declined from their recorded 2025 values. Spring small and large copepods are −0.45 and −0.66 SD using their separate 2000–2026 baselines of 25 observations; summer small and large copepods are −0.92 and +0.21 SD using their separate 2003–2026 baselines of 22 observations. The mixed physical and prey pattern supports contextual discussion rather than an overall favorable or adverse recruitment classification.

The repository energy series has 21 observations during 2002–2025 and numerically matches SEBS annual means rounded to two decimals. Its 2025 value is 19.23 kJ/g dry mass, compared with 19.87 in 2024 and an observed-period mean of 20.78. The separate NBS annual means are retained above. Missing 2026 energy and age-1 observations remain missing; the saved 2025–2026 age-1 estimates are conditional fitted states. The working paper's exact-observation assumption and ±1 estimated-state-SD ribbons describe uncertainty in the fitted states under the assumed model.

The configured DSEM's age-1 rolling-validation RMSE of 1.924 exceeds the AR1 value of 1.847 over the seven observed 2018–2024 targets. The input histories, fitted states and forecast comparisons serve different purposes and retain separate labels. The archived Rceattle pilot uses different timing, scaling and response definitions; its coefficients and fit comparisons apply to that pilot analysis.

Indicator Analysis

Analysis Workflow

The proposed workflow defines the response and cohort timing, assembles the literature-based mechanism table, draws a conceptual graph, and selects a small set of competing hypotheses. Statistical comparisons then use a common observation block and reference model. DSEM can organize these hypotheses while accounting for temporal dynamics (Thorson et al. 2024). Forecast evaluation and management strategy evaluation address whether environmentally informed recruitment methods improve decisions under uncertainty (Haltuch et al. 2019).

The competing hypotheses from the recruitment review are: prey and age-0 energy; predator overlap; spawning and transport; and interactions between prey limitation and predator exposure over time. The present pilot fits only two direct proxy edges. Expansion should prioritize measured intermediates while retaining a small set of graphs that can be identified from the available series.

Proposed Energy-Density Pathway

The new observations allow a measured-condition hypothesis alongside the SST and cold-pool proxies: prey and thermal conditions in birth year t affect age-0 energy density, which contributes to overwinter survival and age-1 recruitment in $t+1$. This is a candidate pathway derived from the literature and workshop synthesis (Heintz et al. 2013; Siddon et al. 2013; Monnahan et al. 2026). It advances the age-1 focus described in the ESR coordination email (Siddon 2026).

Confirm sampling season, response date, and regional overlap before assigning the observation to a pre-winter state. Evaluate SEBS and NBS individually before considering a combined regional index. Any regional aggregation requires documented weights and spatial coverage. The limited and different coverage supports a small set of competing pathways on a common observation block, with observation error and missing years handled explicitly. Preserve direct effects and intermediate-condition hypotheses as alternatives, and use withheld cohorts to evaluate prediction. The added data have descriptive and candidate explanatory roles in this revision; the expanded DSEM remains to be fitted.

Existing DSEM Pilot

The following tables reproduce the EBS pollock website data retrieved on 1 October 2026. The archived pilot used Rceattle 5.23.0 and dsem 3.0.0 with the simple assessment fit; the two-dimensional age-by-year AR1 selectivity sensitivity was excluded. This contribution reuses the archived assessment and DSEM fits.

The archived pilot comparisons are summarized in Table B10.

The SST model has the lowest AIC, while the combined model is 1.63 AIC units higher and the no-edge reference is 2.25 units higher. This is limited support from fit to the same historical data. The reduction in residual variance measures a component of the fitted recruitment model; it differs from the percentage of future recruitment variation predicted successfully. The cold-pool-only result provides little improvement in fitted recruitment variation.

Its standardized direct proxy coefficients and approximate intervals are shown in Table B11.

Both SST coefficients are negative, with their reported intervals below zero. That direction is consistent with the energetic hypothesis, while measured prey and age-0 condition are needed to identify the intermediate mechanism. The cold-pool intervals include zero and the point estimate changes sign in the combined fit. That pattern supports further testing of spatially explicit overlap and the biological timing of the pathway.

Separate age-1 DSEM working paper

A separate age-1 DSEM working paper documents the repository workflow, input versions and numerical checks. It retains its own review status and distinguishes terminal conditional estimates from forecasts. AR1 has lower RMSE in its saved forecast comparison; the archived Rceattle pilot remains a separate analysis.

The recorded input series and their available-record baselines are summarized here (Table B12). The working paper retains its fitted-state and forecast comparisons in the separate report. Its Figures 5–7 show IID, AR1 and recruitment-path state reconstructions. This appendix presents the observed input histories.

Numerical and Scientific Validation

The live CSV files label all four fits OK and record positive-definite Hessians. Recalculation of the diagnostic gate shows that the SST-only fit's maximum gradient, 1.52×10^{-4}, exceeds the required 10^{-4}. The other three pass that gradient threshold. Tightening and rechecking the SST-only fit is required before treating its numerical estimates as a finalized comparison. Compilation and warning logs were outside the archived CSV evidence reviewed here and require confirmation with the fit archive.

The remaining scientific checks are cohort and seasonal timing, alternative spatial domains, observation-error sensitivity, retrospective stability, residual patterns, competing mechanisms, and terminal-year or blocked prediction. Forecast tests must use the assessment estimate and environmental information available at each historical prediction date; later assessment revisions can otherwise introduce future information. Report both uncertainty calibration and improvement over the reference model. A common data block is required within each comparison, with shorter biological series evaluated on their explicitly defined subset.

Decision Support

A contextual indicator may inform the SAFE risk-table discussion when it identifies a documented condition relevant to recruitment, productivity, survey interpretation, or fishery operations. Its influence on a risk category requires an explanation of the evidence and uncertainty. A predictive indicator needs the additional forecast and numerical checks above. Incorporation into an assessment likelihood, projections, or a harvest rule requires a separate, reviewed implementation and evaluation of effects on biomass, uncertainty, ABC, and OFL.

Socioeconomic Monitoring

Socioeconomic monitoring should start with descriptive annual summaries and documented mechanisms. For each series, identify whether its interpretation concerns economic performance, participation, operating constraints, or community exposure. Anomaly scores may help display change once the baseline and coverage are stable. Combining indicators into a single score requires justification for weighting, dependence, and missing values and should remain a proposed analysis until reviewed.

Data Gaps and Future Research Priorities

Ecosystem Priorities

The first priority is to confirm the annual energy-density metadata and connect the regional observations to age-0 season, size, diet, prey, sampling uncertainty, and the appropriate age-1 response. Annual energy density is now available through 2025. Integrating the series into a cohort-aligned model and deriving compatible total-energy measures remain analytical tasks. The second priority is cohort-aligned predator overlap or mortality, with timing that matches the survival interval. Joint juvenile and predator distributions, diet and consumption estimates, and direct fall–spring cohort comparisons would strengthen interpretation.

Transport and nursery delivery, prey timing and lipid content, winter exposure, and annual observation-error estimates remain additional priorities. Climate relationships should be assessed for stability across recent warm conditions and historical regimes. Long-term climate projections in Mueter et al. and Spencer et al. describe conditional scenarios (Mueter et al. 2011; Spencer et al. 2016); their projected effects provide research context rather than annual 2026 indicator values.

Socioeconomic Priorities

Confirm a compact pollock-specific indicator set with economic and social science contributors. Resolve sector and community definitions, annual update capacity, disclosure rules, cost comparability, and a common price base. Evaluate whether incomplete quota utilization or altered fishing patterns have identifiable operational, market, processing, or biological causes. Document the availability date for each product so the ESP distinguishes information usable during the specification process from information released later.

Completion Priorities for the 2026 Appendix

Confirm the primary sources, units, seasonal and spatial definitions and observation uncertainty for the working annual series. Review the age-1 DAG and the proposed energy and predator pathways with the ESP and ecosystem teams. Preserve separate timing and scaling for the archived Rceattle pilot and standalone DSEM, and retain the AR1 forecast benchmark when evaluating any expanded model. Reconcile the historical pilot archive and its outstanding numerical checks before using the indicators to support an annual status or risk classification.

Evidence

The working indicator inventory, descriptive time series and economic context support the annual synthesis within their stated source limits. Primary indicator metadata remain incomplete. The standalone working paper retains its own inputs, uncertainty assumptions and forecast evaluation.

An updated fish-oil yield series, in tons of oil per 100 tons of retained catch, requires oil-specific production and matching retained-catch totals for shoreside and at-sea processing. The grouped production sources combine oil with other products, while the older oil-specific records end in 2021 and lack processing mode. Mode-specific oil yields therefore remain unavailable.

Conclusion

The literature supports following cohorts through spawning and delivery, prey and fall age-0 energy, first-winter survival, and later predator exposure. The ESP also considers adult productivity, survey availability and socioeconomic conditions. The regional energy-density series supplies a measured-condition indicator through 2025. The recorded working series add a mixed 2026 physical and prey context, while their metadata and biological interpretation remain subject to source and contributor checks.

The standalone DSEM's terminal age-1 states are conditional estimates for missing years, and AR1 has lower RMSE in the recorded forecast comparison. Predictive use therefore requires further evaluation against that benchmark. Economic observations extend through 2024 for the BSAI fishery and 2025 for national trade; the 2026 interpretation is qualitative.

Literature Cited

Abelman, Anna, Angela Abolhassani, Michael Dalton, Russel Dame, Ron Felthoven, Brian Garber-Yonts, Jean Lee, Chang Seung, Marysia Szymkowiak, and Sarah Wise. 2026. *Stock Assessment and Fishery Evaluation Report for the*

Groundfish Fisheries of the Gulf of Alaska and Bering Sea/Aleutian Islands Area: Economic Status of the Groundfish Fisheries off Alaska, 2025. Alaska Fisheries Science Center, National Marine Fisheries Service. Published 16 March 2026. Report.

Alaska Fisheries Science Center. 2026a. *Eastern Bering Sea Bottom Trawl Temperature Survey 2026*. NOAA Fisheries. https://www.fisheries.noaa.gov/alaska/science-data/near-real-time-temperatures-2026-bering-sea-bottom-trawl-survey.

Alaska Fisheries Science Center. 2026b. *Economic Status Reports for the Gulf of Alaska and Bering Sea/Aleutian Islands*. NOAA Fisheries report catalogue. https://www.fisheries.noaa.gov/alaska/ecosystems/economic-status-reports-gulf-alaska-and-bering-sea-aleutian-islands.

Alaska Fisheries Science Center. 2026c. *Ecosystem Status Reports for the Gulf of Alaska, Bering Sea and Aleutian Islands*. NOAA Fisheries report catalogue. https://www.fisheries.noaa.gov/alaska/ecosystems/ecosystem-status-reports-gulf-alaska-bering-sea-and-aleutian-islands.

Brodeur, Richard D., Matthew T. Wilson, and Lorenzo Ciannelli. 2000. "Spatial and Temporal Variability in Feeding and Condition of Age-0 Walleye Pollock (Theragra chalcogramma) in Frontal Regions of the Bering Sea." *ICES Journal of Marine Science* 57: 256–64. https://doi.org/10.1006/jmsc.1999.0525.

Buckley, Troy W., Ivonne Ortiz, Stan Kotwicki, and Kerim Aydin. 2016. "Summer Diet Composition of Walleye Pollock and Predator-Prey Relationships with Copepods and Euphausiids in the Eastern Bering Sea, 1987-2011." *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 302–11. https://doi.org/10.1016/j.dsr2.2015.10.009.

Duffy-Anderson, Janet T., Steven J. Barbeaux, Edward V. Farley, et al. 2016. "The Critical First Year of Life of Walleye Pollock (Gadus chalcogrammus) in the Eastern Bering Sea: Implications for Recruitment and Future Research." *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 283–301. https://doi.org/10.1016/j.dsr2.2015.02.001.

EBS Pollock Assessment Team. 2026. *Responses to SSC Comments*. Assessment preparation website. https://noaa-afsc.github.io/EBS_pollock/doc/ssc.html.

Eisner, Lisa B., Ellen M. Yasumiishi, Alexander G. Andrews, and Cecilia A. O'Leary. 2020. "Large Copepods as Leading Indicators of Walleye Pollock Recruitment in the Southeastern Bering Sea: Sample-Based and Spatio-Temporal Model (VAST) Results." *Fisheries Research* 232: 105720. https://doi.org/10.1016/j.fishres.2020.105720.

FAO. 2026. *Global Capture Production.* Release Capture_2026.1.0, March 2026, observations through 2024. Retrieved 30 October 2026. Data collection.

Gann, Jeanette C., Lisa B. Eisner, Steve Porter, et al. 2016. "Possible Mechanism Linking Ocean Conditions to Low Body Weight and Poor Recruitment of Age-0 Walleye Pollock (Gadus chalcogrammus) in the Southeast Bering Sea During 2007." *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 115–27. https://doi.org/10.1016/j.dsr2.2015.07.010.

Haltuch, M. A., E. N. Brooks, J. Brodziak, et al. 2019. "Unraveling the Recruitment Problem: A Review of Environmentally-Informed Forecasting and Management Strategy Evaluation." *Fisheries Research* 217: 198–216. https://doi.org/10.1016/j.fishres.2018.12.016.

Heintz, Ron A., Elizabeth C. Siddon, Edward V. Farley, and Jeffrey M. Napp. 2013. "Correlation Between Recruitment and Fall Condition of Age-0 Pollock (Theragra chalcogramma) from the Eastern Bering Sea Under Varying Climate Conditions." *Deep-Sea Research Part II: Topical Studies in Oceanography* 94: 150–56. https://doi.org/10.1016/j.dsr2.2013.04.006.

Ianelli, James, Sophia Wassermann, Carey McGilliard, et al. 2026. *September 2026 Assessment Developments: Eastern Bering Sea Walleye Pollock*. Draft assessment development document. https://noaa-afsc.github.io/EBS_pollock/doc/EBSpollock_Sept_2026.html.

Monnahan, Cole C., David W. McGowan, Jim Ianelli, Bridget E. Ferriss, and Lauren Rogers. 2026. *Proceedings of the 2025 Workshop on Walleye Pollock Biology and Ecology*. AFSC Processed Report Nos. 2026-09. Alaska Fisheries Science Center, NOAA, National Marine Fisheries Service. https://doi.org/10.25923/hkwv-2f57.

Moss, Jamal H., Edward V. Farley, Angela M. Feldmann, and James N. Ianelli. 2009. “Spatial Distribution, Energetic Status, and Food Habits of Eastern Bering Sea Age-0 Walleye Pollock.” *Transactions of the American Fisheries Society* 138 (3): 497–505. https://doi.org/10.1577/T08-126.1.

Mueter, Franz J., Carol Ladd, Michael C. Palmer, and Brenda L. Norcross. 2006. “Bottom-up and Top-down Controls of Walleye Pollock (Theragra chalcogramma) on the Eastern Bering Sea Shelf.” *Progress in Oceanography* 68 (2–4): 152–83. https://doi.org/10.1016/j.pocean.2006.02.012.

Mueter, Franz J., Nicholas A. Bond, James N. Ianelli, and Anne B. Hollowed. 2011. “Expected Declines in Recruitment of Walleye Pollock (Theragra chalcogramma) in the Eastern Bering Sea Under Future Climate Change.” *ICES Journal of Marine Science* 68 (6): 1284–96. https://doi.org/10.1093/icesjms/fsr022.

NOAA Fisheries. 2026. *Foreign Fishery Trade Data.* FOSS, calendar-year observations through 2025. Retrieved 30 October 2026. Data and documentation.

North Pacific Fishery Management Council Scientific and Statistical Committee. 2024. *SSC Meeting Final Report to the NPFMC Council, December 2024*. North Pacific Fishery Management Council. https://meetings.npfmc.org/CommentReview/DownloadFile?p=dc61bc73-be5c-492f-89b3-a723ea14311d.pdf&fileName=SSC%20Report%20Dec%202024_FINAL.pdf.

Page, Johanna, and Jacek Maselko. 2026. *Annual Age-0 Walleye Pollock Energy Density in the Northern and Southeastern Bering Seas, 2002–2025*. Unpublished contributor dataset, Age0PollockED_DSEM.xlsx, README and DATA sheets.

Petrik, Colleen M., Janet T. Duffy-Anderson, Franz J. Mueter, Katherine Hedstrom, and Enrique N. Curchitser. 2015. “Biophysical Transport Model Suggests Climate Variability Determines Distribution of Walleye Pollock Early Life Stages in the Eastern Bering Sea Through Effects on Spawning.” *Progress in Oceanography* 138: 459–74. https://doi.org/10.1016/j.pocean.2014.06.004.

Shotwell, S. K., K. Blackhart, C. Cunningham, et al. 2023. “Introducing the Ecosystem and Socioeconomic Profile, a Proving Ground for Next Generation Stock Assessments.” *Coastal Management* 51 (5–6): 319–52. https://doi.org/10.1080/08920753.2023.2291858.

Shotwell, S. Kalei, Jenny Bigman, and Russel Dame. 2026. *Appendix 7A. Ecosystem and Socioeconomic Profile of the Arrowtooth Flounder Stock in the Gulf of Alaska*. Draft ESP appendix; Word template.

Siddon, Elizabeth C., Ron A. Heintz, and Franz J. Mueter. 2013. “Conceptual Model of Energy Allocation in Walleye Pollock (Theragra chalcogramma) from Age-0 to Age-1 in the Southeastern Bering Sea.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 94: 140–49. https://doi.org/10.1016/j.dsr2.2012.12.007.

Siddon, Elizabeth. 2026. *EBS Pollock Age-1 Survival DAG and DSEM Coordination*. Email correspondence, 3 September 2026.

Sigler, Michael F., Jeffrey M. Napp, Phyllis J. Stabeno, Ronald A. Heintz, Michael W. Lomas, and George L. Hunt. 2016. “Variation in Annual Production of Copepods, Euphausiids, and Juvenile Walleye Pollock in the Southeastern Bering Sea.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 134: 223–34. https://doi.org/10.1016/j.dsr2.2016.01.003.

Smart, Tracey I., Janet T. Duffy-Anderson, John K. Horne, Edward V. Farley, Christopher D. Wilson, and Jeffrey M. Napp. 2012. “Influence of Environment on Walleye Pollock Eggs, Larvae, and Juveniles in the Southeastern Bering Sea.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 65–70: 196–207. https://doi.org/10.1016/j.dsr2.2012.02.018.

Sogard, Susan M., and Bori L. Olla. 2000. “Endurance of Simulated Winter Conditions by Age-0 Walleye Pollock: Effects of Body Size, Water Temperature and Energy Stores.” *Journal of Fish Biology* 56: 1–21. https://doi.org/10.1111/j.1095-8649.2000.tb02083.x.

Spencer, Paul D., Kirstin K. Holsman, Stephani Zador, et al. 2016. “Modelling Spatially Dependent Predation Mortality of Eastern Bering Sea Walleye Pollock, and Its Implications for Stock Dynamics Under Future Climate Scenarios.” *ICES Journal of Marine Science* 73 (5): 1330–42. https://doi.org/10.1093/icesjms/fsw040.

Strasburger, Wesley W., Nicola Hillgruber, Alexei I. Pinchuk, and Franz J. Mueter. 2014. “Feeding Ecology of Age-0 Walleye Pollock (Gadus chalcogrammus) and Pacific Cod (Gadus macrocephalus) in the Southeastern Bering Sea.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 109: 172–80. https://doi.org/10.1016/j.dsr2.2013.10.007.

Thorson, James T., Alexander G. Andrews III, Timothy E. Essington, and Scott I. Large. 2024. “Dynamic Structural Equation Models Synthesize Ecosystem Dynamics Constrained by Ecological Mechanisms.” *Methods in Ecology and Evolution* 15 (4): 744–55. https://doi.org/10.1111/2041-210X.14289.

Tables

Table B1 *Pollock Review Topics and Proposed ESP Contributions*

Review topic	Contribution in this draft	Completion needed
Contrasting larval, age-0 and adult signals	Follow cohorts through prey, condition and survival stages.	Verify source years, survey domains and cohort identity.
Cannibalism and biomass consumed	Distinguish overlap, diet occurrence, consumption and mortality.	Add reviewed consumption estimates with uncertainty.
Multispecies mortality	Treat age-specific predation as a candidate indicator.	Identify model version, predator coverage and sensitivity.
Weight at age and condition	Include growth and energetic context by age and season.	Update survey and fishery products and assess comparability.
U.S.–Russian distribution	Track survey domain and movement assumptions.	Use verified spatial products and explicit stock boundaries.

Note. Pollock review topics and proposed ESP contributions. Topics are paraphrased from the assessment website; final responses require the original review record.

Table B2 *Source Inventory for the EBS Pollock ESP*

Data group	Source and definition	Status in this contribution
Assessment states and survey inputs	EBS assessment; BTS, ATS and AVO products with age, season and spatial coverage.	Historical development material verified; align with the final 2026 SAFE run.
Temperature and cold pool	Seasonal SST; survey-derived cold-pool extent and temperatures.	Historical standardized pilot inputs archived; 2026 indicator update pending.
Prey and age-0 fish	BASIS/EcoFOCI and associated studies; size, biomass, diet and energetic measurements.	Literature reviewed; consolidate prey, size and sampling uncertainty.
Age-0 energy density	Page and Maselko annual NBS and SEBS means; 2002–2025; kJ/g dry mass.	Regional series received; season, weighting, uncertainty and cohort alignment require confirmation.
Predation	Diet observations, overlap products and multispecies model estimates.	Mechanisms synthesized; reviewed annual products required.
Catch and fishery behavior	Catch accounting, observer and fishery products; consistent season and area definitions.	Indicator specification proposed; extraction and quality review pending.
Economics and communities	Economic SAFE and approved species-level economic and community products.	Inventory proposed; latest complete years and disclosure rules require confirmation.

Note. Source inventory for the EBS pollock contribution. Availability describes this draft's verified inputs and proposed data assembly.

Table B3 *Life-Stage Synthesis for Eastern Bering Sea Pollock*

Stage and timing	Processes and evidence	Priority observations
Spawning and eggs, t	Spawner output, location and transport establish early supply.	Maturity, spawning timing, egg abundance and distribution.
Larvae, t	Prey encounter, growth and delivery to nursery habitat modify supply.	Local prey, larval size and growth, transport and habitat match.
Age 0, summer–fall t	Size and lipid storage determine energy available before winter.	Mass, energy density, total energy, diet quality and prey biomass.
First winter, t to t+1	Reserves, demand, feeding and predation determine early survival.	Fall–spring cohort comparison and winter exposure.
Ages 1–2, t+1 to t+2	Spatial overlap, cannibalism and other predation affect later survival.	Juvenile and predator distributions, diets and mortality.
Adults, later years	Growth, maturity, condition and distribution affect productivity and observation.	Weight at age, condition, maturity and comparable spatial indices.

Note. Life-stage synthesis for eastern Bering Sea pollock. The same cohort is followed from birth year t to ages 1 and 3.

Table B4 *Proposed Mechanisms and Responses*

Driver and mechanism	Expected relation and timing	Evidence and candidate measurement
Spawner output → early supply	Positive toward egg supply in birth year t; later survival modifies the effect.	Smart et al. 2012; SSB, eggs and spawning location.
Prey quality and match → fall energy	Positive toward age-0 energy in t.	Heintz 2013; Siddon 2013; prey lipid, biomass and fish–prey match.
Fall size and energy → winter survival	Positive toward age-1 survival in t+1.	Heintz 2013; Sogard and Olla 2000; mass, energy density and total energy.
Summer mixing → production and prey	Conditional through nutrient supply and prey production in t.	Gann 2016; wind mixing, stratification and nutrients.
Spawning and transport → habitat match	Spatially dependent effect before and during age 0.	Petrik 2015; Smart 2012; transport and nursery-delivery indices.
Predator overlap → juvenile mortality	Negative toward subsequent cohort survival in t+1 and t+2.	Mueter 2006; Spencer 2016; overlap, diet and mortality.
Cold pool → predator access	Hypothesized benefit through separation; depends on predator and stage.	Spencer 2016; common-domain cold pool and overlap.
Late-summer SST → energetic conditions	Hypothesized negative relation with age-1 recruitment after birth-year exposure.	Mueter 2011 and pilot; seasonal SST is a broad proxy.

Note. Proposed mechanism table. Direction refers to the stated response; evidence supports hypotheses rather than a universal coefficient.

Table B5 *Proposed Pollock Socioeconomic Indicators*

Indicator family	Proposed measure	Management question
Participation	Active vessels, processors and deliveries by sector and season.	Where is participation changing, and which operations face exposure?
Fishery performance	Retained catch, TAC utilization and standardized effort measures.	How do fishing conditions and operating choices affect realized harvest?
Value and product mix	Ex-vessel and first-wholesale value, prices and product shares.	How do product demand and prices affect value from the catch?
Costs	Fuel, labor or other comparable cost measures where available.	What evidence indicates changes in operating margins?
Community exposure	Pollock dependence, employment and participation under approved definitions.	Which communities and CDQ beneficiaries are exposed, and through which pathways?
Bycatch and access	Chinook and chum observations, defined rates, season and sector.	How do bycatch conditions and avoidance affect access and operations?

Note. Proposed pollock socioeconomic indicators and the questions they address. All require defined indicators and assessment data.

Table B6 *BSAI Pollock Catch, Participation and Ex-Vessel Economics, 2020–2024*

Measure	Unit	2020	2021	2022	2023	2024
Total catch, all harvesting sectors	1,000 t round weight	1,370.4	1,378.1	1,108.7	1,314.5	1,316.3
Retained catch, all harvesting sectors	1,000 t round weight	1,358.4	1,363.9	1,094.9	1,299.2	1,298.6
Pollock-target vessels (all harvesting sectors)	Vessels	117.0	116.0	114.0	105.0	103.0
Retained catch, trawl catcher vessels	1,000 t round weight	725.0	718.2	570.0	678.2	678.4
Ex-vessel value, trawl catcher vessels	US$ million (nominal)	233.7	242.8	242.0	244.0	213.4
Ex-vessel unit value, trawl catcher vessels	US$/lb (nominal)	0.1	0.2	0.2	0.2	0.1
Trawl catcher-vessel share of retained catch	%	53.4%	52.7%	52.1%	52.2%	52.2%
Pollock-target catcher vessels	Vessels	88.0	84.0	84.0	77.0	76.0

Note. BSAI denotes the Bering Sea and Aleutian Islands and differs from the BS-only assessment catch series. Total catch includes retained and discarded federal and state catch; retained catch and vessel counts cover federal TAC fisheries. The pollock target is assigned by vessel,

week, catching mode, NMFS area and gear. Published catcher-vessel counts describe the target-vessel group; only the catch and value rows explicitly marked trawl carry a separate trawl filter. Unit values and shares are derived from rounded published cells; catcher-vessel unit value combines the reported catcher-vessel catch and value across delivery modes. Monetary values are nominal. Values are rounded to one decimal place. Source: Abelman et al. (2026), Tables 4.1, 4.2, 4.9, 4.10 and 4.14 (printed pp. 34–35, 45–46, 48 and 59), using NMFS accounting and production reports and ADF&G COAR data compiled by AKFIN.

Table B7 *BSAI Pollock First-Wholesale Products and Values, 2020–2024*

Measure	Unit	2020	2021	2022	2023	2024
All products, published volume	1,000 t product weight	492.7	464.1	387.5	480.9	487.9
All products, published value	US$ million (nominal)	1,352.9	1,370.5	1,400.9	1,603.0	1,474.3
All products, unit value	US$/lb (nominal)	1.2	1.3	1.6	1.5	1.4
At-sea share of published value	%	60.4%	61.7%	61.6%	55.6%	57.9%
Fillets, volume	1,000 t product weight	149.4	Unavailable	Unavailable	142.9	158.6
Fillets, share of published value	%	37.4%	Unavailable	Unavailable	40.2%	40.6%
Fillets, unit value	US$/lb (nominal)	1.5	Unavailable	Unavailable	2.0	1.7
Surimi, volume	1,000 t product weight	171.8	187.8	154.8	185.5	165.1
Surimi, share of published value	%	35.2%	42.8%	37.8%	36.0%	32.2%
Surimi, unit value	US$/lb (nominal)	1.3	1.4	1.6	1.4	1.3
Roe, volume	1,000 t product weight	24.7	16.0	13.8	14.7	19.2
Roe, share of published value	%	8.2%	6.5%	7.1%	6.0%	7.5%
Roe, unit value	US$/lb (nominal)	2.0	2.5	3.3	3.0	2.6
At-sea minus shoreside all-product unit value	US$/lb (nominal)	0.4	0.4	0.4	0.1	0.2

Note. BSAI denotes the Bering Sea and Aleutian Islands. Product weight differs from round-weight retained catch. Published aggregates can be incomplete because confidential components are excluded within a table (Economic SAFE §2.2.6). Fillets combine deep-skin and other fillets and exclude minced fish. Fillet volume, value share and unit value are unavailable for 2021–2022 because shoreside deep-skin components are confidential; unavailable cells are not zeros. Unit values and shares use rounded published values. The at-sea-minus-shoreside unit-value difference is unadjusted for product quality and reflects product mix. All dollar values are nominal. Values are rounded to one decimal place. Source: Abelman et al. (2026), Tables 4.16–4.18 (printed pp. 65, 70 and 74), using NMFS production reports and ADF&G COAR data compiled by AKFIN.

Table B8 *Global Alaska Pollock Catch and U.S. Trade*

Panel A. Global Alaska Pollock Catch and Shares, 2020–2024

Metric	Unit	2020	2021	2022	2023	2024
Global Alaska pollock catch	kt live weight	3,544.3	3,484.2	3,380.4	3,543.5	3,573.5
U.S. share of global catch	percent	41.3%	42.0%	36.3%	40.5%	39.9%
Russian share of global catch	percent	51.6%	50.2%	56.3%	53.5%	54.0%
BSAI share of global catch	percent	38.7%	39.6%	32.8%	37.1%	36.8%

Panel B. U.S. Alaska-Pollock Trade, 2020–2025

Metric	Unit	2020	2021	2022	2023	2024	2025
Export volume	kt product weight	323.5	311.4	274.1	326.3	332.5	333.1
Export value	million nominal USD	941.9	898.9	885.6	1,004.4	945.9	1,002.4
Export unit value	USD/lb product weight	1.3	1.3	1.5	1.4	1.3	1.4
Import value (Alaska + unspecified pollock)	million nominal USD	99.7	96.2	141.3	105.5	65.3	48.7
Net exports (legacy basket)	million nominal USD	842.2	802.6	744.4	899.0	880.5	953.6
Japan export volume share	percent	19.3%	21.0%	21.2%	25.0%	19.7%	15.0%
Japan export value share	percent	20.2%	21.1%	20.1%	22.9%	20.7%	16.6%
China, Taiwan and Hong Kong (legacy aggregate) export volume share	percent	9.7%	6.8%	11.6%	7.7%	13.3%	10.6%
China, Taiwan and Hong Kong (legacy aggregate) export value share	percent	6.5%	5.3%	7.5%	5.2%	8.5%	6.7%
Selected European countries export volume share	percent	37.0%	32.7%	31.4%	32.1%	34.2%	37.1%
Selected European countries export value share	percent	39.4%	36.3%	35.3%	37.9%	38.6%	40.7%
Meat export volume share	percent	45.8%	41.1%	42.7%	40.6%	47.5%	53.3%
Meat export value share	percent	44.0%	40.8%	42.5%	43.4%	48.0%	53.0%
Surimi export volume share	percent	47.0%	53.8%	52.2%	54.3%	46.3%	40.4%
Surimi export value share	percent	42.2%	49.4%	49.1%	47.1%	39.1%	34.1%
Roe export volume share	percent	7.3%	5.1%	5.1%	5.2%	6.2%	6.3%
Roe export value share	percent	13.8%	9.8%	8.4%	9.5%	13.0%	13.0%

Note. These observed historical statistics cover global catch and shares through 2024 and U.S. trade through 2025. Trade statistics remain revisable. Global catch is in thousand tons of live weight. Trade volumes are thousand tons of product weight. BSAI catch is a subset of U.S. catch and includes federal and state retained plus discarded catch from both the Bering Sea and Aleutian Islands. Trade covers U.S.-national Alaska-pollock products across their source fisheries. Export values are nominal U.S. dollars on a free-alongside-ship (FAS) export-value basis, distinct from wholesale revenue. Unit values reflect product mix as well as market conditions. Domestic exports exclude re-exports. The legacy import basket combines Alaska and unspecified (NSPF) pollock; legacy net exports subtract that broader basket from Alaska-pollock domestic exports. The legacy statistical aggregate combines mainland China, Taiwan and Hong Kong. The selected European aggregate comprises Austria, Belgium, Denmark, France, Germany, Greece, Ireland, Italy, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, United Kingdom. It is neither the European Union nor all Europe. Meat combines Alaska-pollock products other than surimi and roe, including frozen whole/headed-and-gutted fish, fillets and other meat. Product shares sum to 100% before rounding. Values are rounded to one decimal place from the source extract. Displayed net exports are rounded from the saved source totals rather than recalculated from separately rounded export and import values. Sources: NOAA Fisheries FOSS, retrieved 30 October 2026; FAO Global Capture Production release Capture_2026.1.0 (March 2026); Economic SAFE 2025, Table 4.1, p. 34, published 16 March 2026. Source links: NOAA foreign fishery trade data; FAO Global Capture Production; Economic SAFE.

Table B9 *Annual Mean Age-0 Pollock Energy Density, 2021–2025*

Year	NBS (kJ/g dry mass)	SEBS (kJ/g dry mass)
2021	21.5	20.1
2022	21.5	19.7
2023	21.3	19.8
2024	20.9	19.9
2025	20.8	19.2

Note. Annual mean age-0 pollock energy density for 2021–2025, in kJ/g dry mass. Regions remain separate; sampling uncertainty is pending. Values are rounded to one decimal place. Source: Page and Maselko (2026), DATA sheet.

Table B10 *Archived DSEM Pilot Results*

Specification	Delta AIC	Residual variance reduction	Maximum gradient	Gradient below 1e-4
Age-0 late-summer SST	0.00	10.5%	1.52e-04	Requires refit
Cold pool + age-0 late-summer SST	1.63	12.0%	5.00e-05	Pass
No-edge reference	2.25	0.0%	5.61e-05	Pass
Cold pool	4.10	0.3%	5.69e-05	Pass

Note. Archived live DSEM pilot results. All Hessians are recorded as positive definite. The gradient column applies the strict threshold of 1e-4 independently of the saved OK status.

Table B11 *Standardized Direct Proxy Coefficients From the Archived DSEM Pilot*

Specification	Pathway	Estimate	95 percent interval
Cold pool	Cold pool	0.041	-0.170 to 0.251
Age-0 late-summer SST	Age-0 SST	-0.222	-0.426 to -0.017
Cold pool + age-0 late-summer SST	Cold pool	-0.069	-0.292 to 0.153
Cold pool + age-0 late-summer SST	Age-0 SST	-0.254	-0.484 to -0.024

Note. Standardized direct proxy coefficients from the archived live DSEM pilot, with approximate 95 percent intervals. These fitted paths remain provisional research estimates.

Table B12 *Recorded Annual Input Series and Available-Record Baselines*

Series	Recorded period	Observations	Latest year	Latest departure (SD)
Age-1 pollock response	1964–2024	61	2024	−0.27
Sea-ice extent	1985–2026	42	2026	+0.89
Open-water bloom indicator	1998–2026	29	2026	−1.54
Cold-pool extent	1982–2026	44	2026	+1.12
Sea-surface temperature	1982–2026	44	2026	−0.39
Bottom temperature	1982–2026	44	2026	−0.79
Chlorophyll a	1998–2026	29	2026	+0.77
Euphausiids	2004–2026	14	2026	−0.55
Spring small copepods	2000–2026	25	2026	−0.45
Spring large copepods	2000–2026	25	2026	−0.66
Summer small copepods	2003–2026	22	2026	−0.92
Summer large copepods	2003–2026	22	2026	+0.21
Jellyfish	1982–2026	44	2026	+0.16
Larval pollock abundance	2012–2026	6	2026	−0.10
Juvenile Age-0 Pollock	2003–2026	16	2026	−0.49
SEBS age-0 energy density	2002–2025	21	2025	−1.24
Arrowtooth flounder	1992–2026	34	2026	+0.25
Pacific cod	1982–2026	44	2026	−0.54
Adult pollock	1982–2026	44	2026	+0.82

Note. Recorded annual input series, coverage and latest departures. Source: the saved age-1 DSEM input matrix. Departures use all available observations separately for each series, including its latest year. Gaps remain missing. The full baseline year lists, means and sample SDs are in the indicator evidence table. Units of departure are historical sample SDs; these describe variability rather than sampling uncertainty or approved status thresholds. Primary-source, season, spatial-domain and release metadata require contributor verification. SEBS energy is corroborated in kJ/g dry mass; the age-1 response ends in 2024. Other inputs retain the units and definitions with the recorded data.

Figures



Note. Preliminary Bering Sea recruitment conceptual model reproduced from Monnahan et al. (2026), original Figure 1, printed page 19 (PDF page 27). Source: AFSC Processed Report 2026-09 (Monnahan et al. 2026).

Figure B1 *Preliminary Bering Sea Recruitment Conceptual Model*



Note. Preliminary Bering Sea growth conceptual model reproduced from Monnahan et al. (2026), original Figure 2, printed page 20 (PDF page 28). Source: AFSC Processed Report 2026-09 (Monnahan et al. 2026).

Figure B2 *Preliminary Bering Sea Growth Conceptual Model*



Note. Conceptual cohort sequence from physical conditions and spawning through prey, fall energy, first-winter survival and age-1 recruitment, with subsequent predation affecting survival to age 3. Arrows represent hypotheses supported by the cited literature; they are separate from the fitted direct proxy paths.

Figure B3 *Conceptual Cohort Sequence From Spawning Through Age-3 Survival*



Note. Annual mean age-0 pollock energy density in the northern and southeastern Bering Seas, 2002–2025, in kJ/g dry mass. Points show regional means; lines connect consecutive observed years and break at missing years. Source: Page and Maselko (2026).

Figure B4 *Annual Mean Age-0 Pollock Energy Density, 2002–2025*

Physical environment and primary production



Figure B5 (continued)

Zooplankton and prey



Figure B5 (continued)

Fish abundance, energy density and the age-1 response



Note. Recorded annual input histories grouped for readability. Each value is centered on that series' available-record arithmetic mean and divided by its sample SD, including the latest recorded year. The zero line denotes that mean; gaps are retained without interpolation. Symbols identify recorded values and highlighted endpoints identify each latest record. All 19 series and their reference periods are listed in Table B12. The energy record ends in 2025; the age-1 response ends in 2024. These descriptive histories contain no DSEM fitted states, forecast values or uncertainty bands. Primary-source metadata remain incomplete.

Figure B5 *Recorded Indicator Histories in Standardized Units*



Note. Destination shares of U.S. Alaska pollock domestic exports by (a) product volume and (b) nominal FAS value. The three groups shown cover part of total exports. Selected European countries are Austria, Belgium, Denmark, France, Germany, Greece, Ireland, Italy, Netherlands, Norway, Portugal, Spain, Sweden, Switzerland and the United Kingdom. China, Taiwan and Hong Kong form the legacy aggregate. Shares and definitions match Table B8, panel B. Source: (NOAA Fisheries 2026).

Figure B6 *U.S. Alaska Pollock Export Destination Shares, 2020–2025*



Note. Product shares of U.S. Alaska pollock domestic exports by (a) product volume and (b) nominal FAS value. Meat combines products other than surimi and roe, including frozen whole and headed-and-gutted fish and fillets. The one-decimal source shares are rescaled to sum to 100% within each year, as in the dashboard; the largest adjustment is 0.054 percentage points. Values and product definitions are in Table B8, panel B. Source: (NOAA Fisheries 2026).

Figure B7 *U.S. Alaska Pollock Export Product Mix, 2020–2025*



Note. Shares of global Alaska pollock capture production by live weight. Other U.S. is the U.S. share minus the BSAI share; other countries is the remainder after the United States and Russian Federation. BSAI includes Bering Sea and Aleutian Islands catch. FAO release Capture_2026.1.0 supplies the global totals; the Economic SAFE supplies BSAI catch. Definitions and totals are in Table B8, panel A. Sources: (FAO 2026); (Abelman et al. 2026).

Figure B8 *Global Alaska Pollock Catch by Origin, 2020–2024*



Note. BSAI pollock (a) nominal first-wholesale unit values, (b) production in thousand tons of product weight, and (c) shares of published first-wholesale value. Unit values equal total published value times the product value share, divided by product volume and converted to US$/lb; they reflect product mix as well as market conditions. Fillet observations for 2021–2022 are unavailable because confidential shoreside deep-skin components prevent their calculation. Gaps remain missing rather than zero. Other products account for the remaining value share. Values and definitions match Table B7. Source: (Abelman et al. 2026).

Figure B9 *BSAI Pollock First-Wholesale Product Indicators, 2020–2024*

Appendix C
Research and Other Noncommercial Pollock Removals

Reported noncommercial catch

Reported noncommercial pollock catches in the Bering Sea are summarized by collection in Table C1 and by year in Table C2. The AKR dataset covers 1982–2025 across the Gulf of Alaska, Bering Sea and Aleutian Islands; Bering Sea records begin in 1991, with broader multisurvey coverage from 2010. These are reported collection-year amounts, rather than verified complete EBS-stock totals. Missing years and collection cells mean no record in this extract. The extract ends in 2025.

The summaries use the reported Bering Sea FMP area, retaining stations from collections whose names mention other regions. Explicitly Bogoslof-named surveys are shown separately; other collections in reporting area 518 remain among the other reported Bering Sea catches. In 2024, the total was 216.280081 t, including 4.393640 t from the named Bogoslof survey and 211.886441 t from other reported Bering Sea collections. The corresponding 2025 reported total was 140.994428 t.

Experimental fishing permit (EFP) records account for 1,198.506090 t in 2011 and 771.516610 t in 2021. Their inclusion explains the large totals in those years. The sole 2020 record is a Bogoslof collection; other EBS research removals remain unverified for that year. Historical EFP inclusion in official catch accounting requires reconciliation. AKR metadata state that no 2025 EFP catches occurred outside CAS.

Weights are converted from kilograms to tons using the units corroborated by AKR dataset correspondence, dated 9 October and forwarded on 13 October 2026. Annual completeness of the AKR dataset received on 27 October remains unverified. The 2020–2023 totals agree, after rounding, with the accepted 2024 SAFE's research-catch table (Ianelli et al. 2024). Noncommercial removals are reported separately from assessment catch. The full annual summaries and source record retain the collection details and reporting qualifications.

Exempted fishing and other removal categories

The 2025 EFP 2025-01 permit and 2026 renewal allow up to **3,000 t of total groundfish**, rather than pollock alone, to be exempt from TAC during each permit's effective period; additional groundfish catch counts against TAC. This authorization ceiling is separate from observed removals. Actual pollock catch under these permits and its accounting treatment require permit-specific records.

EFP 2024-02 provides a footrope/gear exemption and otherwise retains applicable regulations. It supplies no basis for assuming a TAC exemption. The noncommercial dataset includes BS-labelled subsistence records in 1991 and 1994; complete recent stock-specific subsistence, recreational and other-FMP removals remain unverified.

Tables

Table C1 *Reported Bering Sea Noncommercial Pollock Catch by Collection and Year*

Panel A. 1991–2009

Collection (agency)	1991	1994	1997	1999	2001	2003	2005	2007	2009
AFSC Annual Longline Survey (NMFS)	NA	NA	1.0	2.1	4.0	3.1	1.1	0.9	1.4
Subsistence Fishery (ADFG)	<0.1	<0.1	NA	NA	NA	NA	NA	NA	NA

Panel B. 2010–2017

Collection (agency)	2010	2011	2012	2013	2014	2015	2016	2017
AFSC Annual Longline Survey (NMFS)	NA	3.9	NA	2.3	NA	3.4	NA	1.6
Aleutian Island Bottom Trawl Survey (NMFS)	27.0	NA	4.5	NA	12.3	NA	2.6	NA
Bering Sea Acoustic Survey (NMFS)	49.6	NA	NA	NA	NA	NA	NA	NA
Bering Sea Bottom Trawl Survey (NMFS)	140.7	NA	NA	NA	NA	NA	NA	NA
Bering Sea Slope Survey (NMFS)	3.3	NA	7.7	NA	NA	NA	12.4	NA
Bogoslof EIT Survey with Northern Extensions (NMFS)	NA	NA	4.1	NA	NA	NA	NA	NA
Eastern Bering Sea Bottom Trawl Survey (NMFS)	NA	116.5	140.0	178.7	293.4	249.7	177.4	181.1
Gulf of Alaska Bottom Trawl Survey (NMFS)	NA	<0.1	NA	<0.1	NA	<0.1	NA	NA
IPHC Annual Longline Survey (IPHC)	0.7	0.2	0.1	0.2	0.6	0.7	0.9	0.1
Large-Mesh Trawl Survey (ADFG)	3.7	3.2	2.4	2.2	1.4	2.1	4.6	5.0
Northern Bering Sea Bottom Trawl Survey (NMFS)	0.6	NA	NA	NA	NA	NA	NA	38.1
Pollock EFP 11-01 (NMFS)	NA	1,198.5	3.8	NA	NA	NA	NA	NA
Pribilof Islands Crab Survey (ADFG)	NA	0.1	NA	NA	NA	NA	0.3	NA
St. Matthews Crab Survey (ADFG)	0.2	NA	NA	NA	NA	NA	NA	0.2
Summer EBS Survey with Russia (NMFS)	NA	NA	45.2	NA	NA	NA	NA	NA

Panel C. 2018–2025

Collection (agency)	2018	2019	2020	2021	2022	2023	2024	2025
AFSC Annual Longline Survey (NMFS)	NA	1.3	NA	0.9	NA	0.7	NA	NA
Aleutian Island Bottom Trawl Survey (NMFS)	6.1	NA	NA	NA	9.7	NA	5.0	NA
BSAI Trawl Salmon Excluder Device EFP 2018-03-02 (NMFS)	NA	NA	NA	771.5	NA	NA	NA	NA
Blue King Crab Pot Survey (ADFG)	<0.1	NA	NA	NA	0.1	NA	NA	NA
Bogoslof Island Walleye Pollock Acoustic-Trawl Survey (NMFS)	6.0	NA	NA	NA	NA	NA	NA	NA
Eastern Bering Sea Bottom Trawl Survey (NMFS)	111.7	188.4	NA	107.8	168.1	118.0	163.2	120.4
Eastern Bering Sea Walleye Pollock Acoustic-Trawl Survey (NMFS)	36.7	NA	NA	NA	48.3	NA	NA	NA
Gulf of Alaska Bottom Trawl Survey (NMFS)	NA	<0.1	NA	NA	NA	<0.1	NA	NA
IPHC Annual Longline Survey (IPHC)	0.1	0.1	NA	<0.1	0.2	NA	0.1	NA
Large-Mesh Trawl Survey (ADFG)	1.8	1.8	NA	1.4	6.1	0.7	3.8	2.1
Northern Bering Sea Bottom Trawl Survey (NMFS)	15.0	38.6	NA	16.0	13.1	NA	NA	18.5
Summer Acoustic-Trawl Survey of Walleye Pollock in the Eastern Bering Sea (NMFS)	NA	NA	NA	NA	NA	NA	39.8	NA
Winter Acoustic-Trawl Survey of Walleye Pollock Near Bogoslof Island and vicinity (NMFS)	NA	NA	12.9	NA	NA	NA	4.4	NA

Note. Values are metric tons (source kilograms divided by 1,000), rounded to one decimal place. Source: received Noncommercial Fishery Catch (1).csv, 27 October 2026 received-date field. Reported records have FMP Area = BS and Species Group Code = PLCK. All reporting areas are retained, including 518. NA indicates missing records; 0.0 indicates a reported zero. Early panels show observed BS years only, and collection rows with no records in a panel are omitted from that panel. All collection-year cells and full source precision are retained in the accompanying CSV. Scope follows the FMP field, including collections whose names mention other regions. Positive amounts that round to 0.0 t are shown as <0.1; 0.0 denotes a source-recorded zero.

Table C2 *Annual Reported Bering Sea Pollock Catch by Source Category*

Collection year	Reported BS total (t)	EFP (t)	Explicitly Bogoslof-named (t)	Other reported BS (t)	Source records
1991	<0.1	NA	NA	<0.1	1
1992	NA	NA	NA	NA	NA
1993	NA	NA	NA	NA	NA
1994	<0.1	NA	NA	<0.1	1
1995	NA	NA	NA	NA	NA
1996	NA	NA	NA	NA	NA
1997	1.0	NA	NA	1.0	4
1998	NA	NA	NA	NA	NA
1999	2.1	NA	NA	2.1	4
2000	NA	NA	NA	NA	NA
2001	4.0	NA	NA	4.0	4
2002	NA	NA	NA	NA	NA
2003	3.1	NA	NA	3.1	4
2004	NA	NA	NA	NA	NA
2005	1.1	NA	NA	1.1	3
2006	NA	NA	NA	NA	NA
2007	0.9	NA	NA	0.9	4
2008	NA	NA	NA	NA	NA
2009	1.4	NA	NA	1.4	4
2010	226.0	NA	NA	226.0	65
2011	1,322.5	1,198.5	NA	124.0	30
2012	207.9	3.8	4.1	200.0	38
2013	183.4	NA	NA	183.4	56
2014	307.7	NA	NA	307.7	26
2015	255.9	NA	NA	255.9	30
2016	198.1	NA	NA	198.1	32
2017	226.1	NA	NA	226.1	29
2018	177.4	NA	6.0	171.4	38
2019	230.1	NA	NA	230.1	29
2020	12.9	NA	12.9	NA	1
2021	897.6	771.5	NA	126.1	34
2022	245.7	NA	NA	245.7	40
2023	119.4	NA	NA	119.4	19
2024	216.3	NA	4.4	211.9	33
2025	141.0	NA	NA	141.0	20
2026	NA	NA	NA	NA	NA

Note. Weights are metric tons, rounded to one decimal place. Categories are mutually exclusive source-label summaries. EFP comprises the two explicitly EFP-named collections; the Bogoslof column includes collections explicitly named for Bogoslof. Other reported BS includes all remaining BS-labelled records, including area-518 records from other collections. NA indicates missing records; 0.0 indicates a reported zero. The row count measures records. Totals cover the reported records; annual coverage and overlap with official catch accounting remain uncertain.

Appendix D
Detailed Current-Model Estimates

All estimated parameters and their asymptotic uncertainty are listed below (Table D1), followed by the largest parameter correlations (Table D2). These estimates use the selected model and fixed biological schedules. The full numerical products also retain natural-scale transformations and all shared parameter positions.

Table D1 *Model 26.0 Estimated Parameters and Approximate 95% Confidence Intervals*

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0001	Recruitment level	R0	9.6728	0.1212	9.4352	9.9104
P0002	Recruitment deviation	1964	−1.0198	0.4681	−1.9373	−0.1023
P0003	Recruitment deviation	1965	0.3917	0.2916	−0.1798	0.9632
P0004	Recruitment deviation	1966	0.0001	0.3593	−0.7041	0.7042
P0005	Recruitment deviation	1967	0.5387	0.2917	−0.0330	1.1104
P0006	Recruitment deviation	1968	0.3459	0.3165	−0.2745	0.9662
P0007	Recruitment deviation	1969	0.5038	0.2931	−0.0705	1.0782
P0008	Recruitment deviation	1970	0.3846	0.3052	−0.2136	0.9828
P0009	Recruitment deviation	1971	−0.1528	0.3739	−0.8856	0.5800
P0010	Recruitment deviation	1972	−0.3766	0.3851	−1.1314	0.3782
P0011	Recruitment deviation	1973	0.5213	0.2325	0.0655	0.9770
P0012	Recruitment deviation	1974	0.1941	0.2371	−0.2706	0.6589
P0013	Recruitment deviation	1975	0.0354	0.2205	−0.3967	0.4675
P0014	Recruitment deviation	1976	−0.2044	0.2104	−0.6169	0.2081
P0015	Recruitment deviation	1977	−0.1243	0.1911	−0.4989	0.2503
P0016	Recruitment deviation	1978	0.5126	0.1555	0.2078	0.8175
P0017	Recruitment deviation	1979	1.3692	0.1375	1.0997	1.6387
P0018	Recruitment deviation	1980	0.6249	0.1492	0.3324	0.9174
P0019	Recruitment deviation	1981	0.7685	0.1463	0.4817	1.0553
P0020	Recruitment deviation	1982	0.1889	0.1614	−0.1274	0.5052
P0021	Recruitment deviation	1983	1.1863	0.1357	0.9204	1.4521
P0022	Recruitment deviation	1984	−0.1097	0.1634	−0.4299	0.2104
P0023	Recruitment deviation	1985	0.7176	0.1390	0.4450	0.9901
P0024	Recruitment deviation	1986	−0.2200	0.1558	−0.5254	0.0853
P0025	Recruitment deviation	1987	−0.7740	0.1648	−1.0971	−0.4510
P0026	Recruitment deviation	1988	−1.0224	0.1658	−1.3474	−0.6974
P0027	Recruitment deviation	1989	−0.2878	0.1424	−0.5669	−0.0086
P0028	Recruitment deviation	1990	1.1562	0.1262	0.9090	1.4035
P0029	Recruitment deviation	1991	0.5442	0.1307	0.2881	0.8003
P0030	Recruitment deviation	1992	0.4032	0.1325	0.1436	0.6628
P0031	Recruitment deviation	1993	1.0698	0.1269	0.8211	1.3186
P0032	Recruitment deviation	1994	0.0032	0.1356	−0.2625	0.2690
P0033	Recruitment deviation	1995	−0.3517	0.1389	−0.6239	−0.0794

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0034	Recruitment deviation	1996	0.3553	0.1292	0.1020	0.6086
P0035	Recruitment deviation	1997	0.6464	0.1269	0.3977	0.8950
P0036	Recruitment deviation	1998	−0.0602	0.1317	−0.3183	0.1978
P0037	Recruitment deviation	1999	0.0094	0.1306	−0.2466	0.2654
P0038	Recruitment deviation	2000	0.4371	0.1273	0.1876	0.6866
P0039	Recruitment deviation	2001	0.7735	0.1252	0.5281	1.0189
P0040	Recruitment deviation	2002	0.3678	0.1265	0.1199	0.6157
P0041	Recruitment deviation	2003	−0.1019	0.1286	−0.3540	0.1502
P0042	Recruitment deviation	2004	−0.8894	0.1362	−1.1564	−0.6224
P0043	Recruitment deviation	2005	−1.2337	0.1415	−1.5110	−0.9563
P0044	Recruitment deviation	2006	−0.3192	0.1309	−0.5758	−0.0627
P0045	Recruitment deviation	2007	0.4533	0.1271	0.2041	0.7025
P0046	Recruitment deviation	2008	−0.1584	0.1330	−0.4192	0.1024
P0047	Recruitment deviation	2009	1.1374	0.1249	0.8925	1.3823
P0048	Recruitment deviation	2010	0.3763	0.1287	0.1240	0.6287
P0049	Recruitment deviation	2011	−0.1110	0.1328	−0.3713	0.1493
P0050	Recruitment deviation	2012	−0.2546	0.1337	−0.5166	0.0074
P0051	Recruitment deviation	2013	1.0028	0.1249	0.7580	1.2475
P0052	Recruitment deviation	2014	1.1366	0.1252	0.8912	1.3819
P0053	Recruitment deviation	2015	0.2917	0.1313	0.0344	0.5490
P0054	Recruitment deviation	2016	−0.6103	0.1475	−0.8995	−0.3212
P0055	Recruitment deviation	2017	−0.6994	0.1519	−0.9971	−0.4018
P0056	Recruitment deviation	2018	0.0578	0.1410	−0.2187	0.3342
P0057	Recruitment deviation	2019	1.7104	0.1302	1.4552	1.9656
P0058	Recruitment deviation	2020	0.5518	0.1454	0.2667	0.8368
P0059	Recruitment deviation	2021	−0.2033	0.1786	−0.5534	0.1467
P0060	Recruitment deviation	2022	−0.2247	0.2049	−0.6263	0.1768
P0061	Recruitment deviation	2023	−0.3126	0.2286	−0.7606	0.1353
P0062	Recruitment deviation	2024	0.9050	0.2517	0.4118	1.3983
P0063	Recruitment deviation	2025	0.5663	0.2920	−0.0060	1.1387
P0064	Recruitment deviation	2026	0.8281	0.5378	−0.2260	1.8822
P0065	Initial-age deviation	Age1	−0.3949	0.3509	−1.0827	0.2928
P0066	Initial-age deviation	Age2	−0.4683	0.3455	−1.1455	0.2089
P0067	Initial-age deviation	Age3	−1.2468	0.4532	−2.1351	−0.3586

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0068	Initial-age deviation	Age4	−1.4797	0.5231	−2.5049	−0.4545
P0069	Initial-age deviation	Age5	−0.9725	0.4745	−1.9026	−0.0424
P0070	Initial-age deviation	Age6	−1.1558	0.5508	−2.2353	−0.0762
P0071	Initial-age deviation	Age7	−1.3132	0.6310	−2.5499	−0.0765
P0072	Initial-age deviation	Age8	−1.2359	0.6710	−2.5510	0.0791
P0073	Initial-age deviation	Age9	−1.0702	0.6957	−2.4338	0.2934
P0074	Initial-age deviation	Age10	−0.9172	0.7220	−2.3322	0.4979
P0075	Initial-age deviation	Age11	−0.7788	0.7491	−2.2470	0.6894
P0076	Initial-age deviation	Age12	−0.6579	0.7760	−2.1789	0.8631
P0077	Initial-age deviation	Age13	−0.5491	0.8035	−2.1239	1.0256
P0078	Initial-age deviation	Age14	−1.0041	0.7100	−2.3957	0.3874
P0079	Log fishing scale	Fishery; 1964	−2.9386	0.2174	−3.3647	−2.5125
P0080	Log fishing scale	Fishery; 1965	−2.7303	0.2116	−3.1450	−2.3155
P0081	Log fishing scale	Fishery; 1966	−2.6737	0.2155	−3.0960	−2.2513
P0082	Log fishing scale	Fishery; 1967	−2.0362	0.2270	−2.4811	−1.5913
P0083	Log fishing scale	Fishery; 1968	−1.9723	0.2260	−2.4153	−1.5292
P0084	Log fishing scale	Fishery; 1969	−1.9071	0.2317	−2.3611	−1.4530
P0085	Log fishing scale	Fishery; 1970	−1.5364	0.2309	−1.9890	−1.0838
P0086	Log fishing scale	Fishery; 1971	−1.1811	0.2325	−1.6368	−0.7254
P0087	Log fishing scale	Fishery; 1972	−1.0495	0.2265	−1.4935	−0.6055
P0088	Log fishing scale	Fishery; 1973	−0.9062	0.2295	−1.3561	−0.4563
P0089	Log fishing scale	Fishery; 1974	−0.8209	0.2280	−1.2679	−0.3740
P0090	Log fishing scale	Fishery; 1975	−0.9945	0.2055	−1.3973	−0.5916
P0091	Log fishing scale	Fishery; 1976	−1.0737	0.2009	−1.4673	−0.6800
P0092	Log fishing scale	Fishery; 1977	−1.1394	0.2153	−1.5614	−0.7174
P0093	Log fishing scale	Fishery; 1978	−1.0271	0.2171	−1.4525	−0.6017
P0094	Log fishing scale	Fishery; 1979	−0.9874	0.2107	−1.4003	−0.5745
P0095	Log fishing scale	Fishery; 1980	−1.0814	0.1983	−1.4701	−0.6927
P0096	Log fishing scale	Fishery; 1981	−1.4230	0.1935	−1.8023	−1.0438
P0097	Log fishing scale	Fishery; 1982	−1.8679	0.1901	−2.2404	−1.4954
P0098	Log fishing scale	Fishery; 1983	−2.0507	0.1833	−2.4100	−1.6914
P0099	Log fishing scale	Fishery; 1984	−2.0663	0.1527	−2.3655	−1.7671
P0100	Log fishing scale	Fishery; 1985	−2.0559	0.1408	−2.3318	−1.7801
P0101	Log fishing scale	Fishery; 1986	−2.1239	0.1347	−2.3878	−1.8599

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0102	Log fishing scale	Fishery; 1987	−2.4130	0.1266	−2.6611	−2.1649
P0103	Log fishing scale	Fishery; 1988	−2.0653	0.1178	−2.2962	−1.8344
P0104	Log fishing scale	Fishery; 1989	−2.0069	0.1006	−2.2040	−1.8098
P0105	Log fishing scale	Fishery; 1990	−1.5822	0.0944	−1.7672	−1.3972
P0106	Log fishing scale	Fishery; 1991	−1.4032	0.0849	−1.5696	−1.2368
P0107	Log fishing scale	Fishery; 1992	−1.1200	0.0920	−1.3004	−0.9396
P0108	Log fishing scale	Fishery; 1993	−1.5672	0.1086	−1.7800	−1.3543
P0109	Log fishing scale	Fishery; 1994	−1.8109	0.1176	−2.0413	−1.5804
P0110	Log fishing scale	Fishery; 1995	−1.8155	0.1118	−2.0347	−1.5964
P0111	Log fishing scale	Fishery; 1996	−1.6567	0.1114	−1.8751	−1.4383
P0112	Log fishing scale	Fishery; 1997	−1.6452	0.1244	−1.8890	−1.4014
P0113	Log fishing scale	Fishery; 1998	−1.8130	0.1174	−2.0431	−1.5830
P0114	Log fishing scale	Fishery; 1999	−2.0245	0.1131	−2.2463	−1.8028
P0115	Log fishing scale	Fishery; 2000	−1.9078	0.0941	−2.0923	−1.7232
P0116	Log fishing scale	Fishery; 2001	−1.7581	0.0844	−1.9236	−1.5926
P0117	Log fishing scale	Fishery; 2002	−1.5902	0.0776	−1.7424	−1.4381
P0118	Log fishing scale	Fishery; 2003	−1.5098	0.0772	−1.6611	−1.3585
P0119	Log fishing scale	Fishery; 2004	−1.5953	0.0838	−1.7596	−1.4311
P0120	Log fishing scale	Fishery; 2005	−1.6359	0.0829	−1.7985	−1.4734
P0121	Log fishing scale	Fishery; 2006	−1.4267	0.0820	−1.5873	−1.2660
P0122	Log fishing scale	Fishery; 2007	−1.3555	0.0809	−1.5141	−1.1970
P0123	Log fishing scale	Fishery; 2008	−1.2726	0.0841	−1.4374	−1.1078
P0124	Log fishing scale	Fishery; 2009	−1.3188	0.0943	−1.5036	−1.1341
P0125	Log fishing scale	Fishery; 2010	−1.5006	0.1034	−1.7033	−1.2978
P0126	Log fishing scale	Fishery; 2011	−1.1973	0.1005	−1.3944	−1.0003
P0127	Log fishing scale	Fishery; 2012	−1.2027	0.1268	−1.4512	−0.9542
P0128	Log fishing scale	Fishery; 2013	−1.2580	0.1457	−1.5435	−0.9725
P0129	Log fishing scale	Fishery; 2014	−1.3178	0.1579	−1.6272	−1.0085
P0130	Log fishing scale	Fishery; 2015	−1.2383	0.1649	−1.5615	−0.9152
P0131	Log fishing scale	Fishery; 2016	−1.5367	0.1456	−1.8220	−1.2513
P0132	Log fishing scale	Fishery; 2017	−1.5614	0.1284	−1.8130	−1.3097
P0133	Log fishing scale	Fishery; 2018	−1.7471	0.1197	−1.9818	−1.5124
P0134	Log fishing scale	Fishery; 2019	−1.6372	0.1073	−1.8475	−1.4269
P0135	Log fishing scale	Fishery; 2020	−1.3656	0.1186	−1.5980	−1.1331

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0136	Log fishing scale	Fishery; 2021	−1.4188	0.1220	−1.6579	−1.1798
P0137	Log fishing scale	Fishery; 2022	−1.7708	0.1385	−2.0422	−1.4993
P0138	Log fishing scale	Fishery; 2023	−1.8165	0.1494	−2.1092	−1.5238
P0139	Log fishing scale	Fishery; 2024	−1.9059	0.1439	−2.1878	−1.6239
P0140	Log fishing scale	Fishery; 2025	−1.8131	0.1338	−2.0753	−1.5508
P0141	Log fishing scale	Fishery; 2026	−1.6691	0.1614	−1.9853	−1.3528
P0142	Log catchability	AVO	−8.2249	0.0711	−8.3642	−8.0855
P0143	Log catchability	ATS	−0.6599	0.0620	−0.7814	−0.5384
P0144	Log catchability	CPUE	−0.2481	0.1849	−0.6104	0.1143
P0145	Selectivity increment	Fishery; Bin1; 1964 *	1.2875	0.5969	0.1177	2.4573
P0146	Selectivity increment	Fishery; Bin2; 1964 *	0.7997	0.4928	−0.1663	1.7656
P0147	Selectivity increment	Fishery; Bin3; 1964 *	0.4574	0.4383	−0.4016	1.3164
P0148	Selectivity increment	Fishery; Bin4; 1964 *	0.3677	0.3979	−0.4122	1.1476
P0149	Selectivity increment	Fishery; Bin5; 1964 *	−0.1820	0.3965	−0.9591	0.5952
P0150	Selectivity increment	Fishery; Bin6; 1964 *	−0.4579	0.2958	−1.0377	0.1219
P0151	Selectivity increment	Fishery; Bin7; 1964 *	−0.5719	0.2747	−1.1103	−0.0335
P0152	Selectivity increment	Fishery; Bin8; 1964 *	−0.5861	0.2680	−1.1114	−0.0608
P0153	Selectivity increment	Fishery; Bin9; 1964 *	−0.4992	0.2677	−1.0239	0.0255
P0154	Selectivity increment	Fishery; Bin10; 1964 *	−0.2953	0.2716	−0.8277	0.2371
P0155	Selectivity increment	Fishery; Bin11; 1964 *	−0.2831	0.2799	−0.8317	0.2654
P0156	Selectivity increment	Fishery; Bin12; 1964 *	−0.0367	0.2934	−0.6117	0.5383
P0157	Selectivity increment	Fishery; Bin1; 1965 *	0.3571	0.4151	−0.4565	1.1708
P0158	Selectivity increment	Fishery; Bin2; 1965 *	0.1619	0.4026	−0.6272	0.9511
P0159	Selectivity increment	Fishery; Bin3; 1965 *	0.1307	0.3944	−0.6423	0.9037
P0160	Selectivity increment	Fishery; Bin4; 1965 *	0.0902	0.3777	−0.6500	0.8304
P0161	Selectivity increment	Fishery; Bin5; 1965 *	−0.0379	0.3792	−0.7810	0.7053
P0162	Selectivity increment	Fishery; Bin6; 1965 *	−0.1784	0.3055	−0.7771	0.4203
P0163	Selectivity increment	Fishery; Bin7; 1965 *	−0.1639	0.2839	−0.7203	0.3926
P0164	Selectivity increment	Fishery; Bin8; 1965 *	−0.1318	0.2797	−0.6799	0.4163
P0165	Selectivity increment	Fishery; Bin9; 1965 *	−0.0992	0.2797	−0.6475	0.4491
P0166	Selectivity increment	Fishery; Bin10; 1965 *	−0.0684	0.2813	−0.6198	0.4830
P0167	Selectivity increment	Fishery; Bin11; 1965 *	−0.0439	0.2832	−0.5990	0.5111
P0168	Selectivity increment	Fishery; Bin12; 1965 *	−0.0165	0.2786	−0.5626	0.5295
P0169	Selectivity increment	Fishery; Bin1; 1966 *	0.3379	0.4159	−0.4772	1.1530

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0170	Selectivity increment	Fishery; Bin2; 1966 *	0.1962	0.4019	−0.5915	0.9838
P0171	Selectivity increment	Fishery; Bin3; 1966 *	0.1357	0.3933	−0.6351	0.9065
P0172	Selectivity increment	Fishery; Bin4; 1966 *	−0.0528	0.3090	−0.6585	0.5528
P0173	Selectivity increment	Fishery; Bin5; 1966 *	−0.0653	0.2844	−0.6227	0.4921
P0174	Selectivity increment	Fishery; Bin6; 1966 *	−0.0912	0.2786	−0.6373	0.4549
P0175	Selectivity increment	Fishery; Bin7; 1966 *	−0.0904	0.2774	−0.6341	0.4534
P0176	Selectivity increment	Fishery; Bin8; 1966 *	−0.0915	0.2773	−0.6350	0.4519
P0177	Selectivity increment	Fishery; Bin9; 1966 *	−0.0850	0.2779	−0.6298	0.4597
P0178	Selectivity increment	Fishery; Bin10; 1966 *	−0.0744	0.2795	−0.6221	0.4733
P0179	Selectivity increment	Fishery; Bin11; 1966 *	−0.0638	0.2811	−0.6149	0.4872
P0180	Selectivity increment	Fishery; Bin12; 1966 *	−0.0553	0.2749	−0.5940	0.4834
P0181	Selectivity increment	Fishery; Bin1; 1967 *	0.3044	0.4161	−0.5112	1.1200
P0182	Selectivity increment	Fishery; Bin2; 1967 *	0.1602	0.3988	−0.6214	0.9418
P0183	Selectivity increment	Fishery; Bin3; 1967 *	0.2062	0.3907	−0.5595	0.9719
P0184	Selectivity increment	Fishery; Bin4; 1967 *	−0.1132	0.3085	−0.7179	0.4915
P0185	Selectivity increment	Fishery; Bin5; 1967 *	−0.1023	0.2828	−0.6566	0.4520
P0186	Selectivity increment	Fishery; Bin6; 1967 *	−0.0914	0.2781	−0.6364	0.4536
P0187	Selectivity increment	Fishery; Bin7; 1967 *	−0.0794	0.2773	−0.6229	0.4641
P0188	Selectivity increment	Fishery; Bin8; 1967 *	−0.0677	0.2772	−0.6110	0.4757
P0189	Selectivity increment	Fishery; Bin9; 1967 *	−0.0629	0.2779	−0.6075	0.4818
P0190	Selectivity increment	Fishery; Bin10; 1967 *	−0.0573	0.2794	−0.6049	0.4904
P0191	Selectivity increment	Fishery; Bin11; 1967 *	−0.0511	0.2811	−0.6020	0.4998
P0192	Selectivity increment	Fishery; Bin12; 1967 *	−0.0455	0.2747	−0.5838	0.4928
P0193	Selectivity increment	Fishery; Bin1; 1968 *	0.3127	0.4163	−0.5033	1.1286
P0194	Selectivity increment	Fishery; Bin2; 1968 *	0.1419	0.3978	−0.6377	0.9215
P0195	Selectivity increment	Fishery; Bin3; 1968 *	0.0120	0.3855	−0.7436	0.7675
P0196	Selectivity increment	Fishery; Bin4; 1968 *	−0.0108	0.3097	−0.6179	0.5963
P0197	Selectivity increment	Fishery; Bin5; 1968 *	−0.0366	0.2850	−0.5952	0.5220
P0198	Selectivity increment	Fishery; Bin6; 1968 *	−0.0514	0.2793	−0.5988	0.4960
P0199	Selectivity increment	Fishery; Bin7; 1968 *	−0.0611	0.2781	−0.6061	0.4839
P0200	Selectivity increment	Fishery; Bin8; 1968 *	−0.0644	0.2790	−0.6113	0.4824
P0201	Selectivity increment	Fishery; Bin9; 1968 *	−0.0626	0.2800	−0.6114	0.4863
P0202	Selectivity increment	Fishery; Bin10; 1968 *	−0.0612	0.2825	−0.6149	0.4925
P0203	Selectivity increment	Fishery; Bin11; 1968 *	−0.0596	0.2834	−0.6150	0.4958

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0204	Selectivity increment	Fishery; Bin12; 1968 *	−0.0589	0.2764	−0.6007	0.4829
P0205	Selectivity increment	Fishery; Bin1; 1969 *	0.2851	0.4165	−0.5312	1.1014
P0206	Selectivity increment	Fishery; Bin2; 1969 *	0.0924	0.3968	−0.6852	0.8701
P0207	Selectivity increment	Fishery; Bin3; 1969 *	0.1579	0.3857	−0.5981	0.9139
P0208	Selectivity increment	Fishery; Bin4; 1969 *	−0.0960	0.3101	−0.7037	0.5117
P0209	Selectivity increment	Fishery; Bin5; 1969 *	−0.0947	0.2880	−0.6590	0.4697
P0210	Selectivity increment	Fishery; Bin6; 1969 *	−0.0822	0.2892	−0.6490	0.4846
P0211	Selectivity increment	Fishery; Bin7; 1969 *	−0.0710	0.3110	−0.6805	0.5385
P0212	Selectivity increment	Fishery; Bin8; 1969 *	−0.0543	0.3363	−0.7134	0.6047
P0213	Selectivity increment	Fishery; Bin9; 1969 *	−0.0484	0.3372	−0.7094	0.6125
P0214	Selectivity increment	Fishery; Bin10; 1969 *	−0.0360	0.4001	−0.8201	0.7481
P0215	Selectivity increment	Fishery; Bin11; 1969 *	−0.0272	0.3244	−0.6629	0.6085
P0216	Selectivity increment	Fishery; Bin12; 1969 *	−0.0257	0.2858	−0.5858	0.5345
P0217	Selectivity increment	Fishery; Bin1; 1970 *	0.3011	0.4170	−0.5161	1.1183
P0218	Selectivity increment	Fishery; Bin2; 1970 *	0.1488	0.3965	−0.6283	0.9259
P0219	Selectivity increment	Fishery; Bin3; 1970 *	−0.0316	0.3840	−0.7843	0.7211
P0220	Selectivity increment	Fishery; Bin4; 1970 *	−0.2072	0.3635	−0.9196	0.5052
P0221	Selectivity increment	Fishery; Bin5; 1970 *	−0.1218	0.3161	−0.7413	0.4976
P0222	Selectivity increment	Fishery; Bin6; 1970 *	−0.1011	0.3166	−0.7217	0.5194
P0223	Selectivity increment	Fishery; Bin7; 1970 *	−0.0529	0.3685	−0.7751	0.6693
P0224	Selectivity increment	Fishery; Bin8; 1970 *	−0.0096	0.3991	−0.7920	0.7727
P0225	Selectivity increment	Fishery; Bin9; 1970 *	−0.0009	0.3998	−0.7845	0.7828
P0226	Selectivity increment	Fishery; Bin10; 1970 *	0.0133	0.4076	−0.7855	0.8122
P0227	Selectivity increment	Fishery; Bin11; 1970 *	0.0230	0.4049	−0.7706	0.8167
P0228	Selectivity increment	Fishery; Bin12; 1970 *	0.0389	0.3034	−0.5557	0.6335
P0229	Selectivity increment	Fishery; Bin1; 1971 *	0.0965	0.4172	−0.7211	0.9142
P0230	Selectivity increment	Fishery; Bin2; 1971 *	0.0278	0.3955	−0.7474	0.8031
P0231	Selectivity increment	Fishery; Bin3; 1971 *	−0.0921	0.3808	−0.8385	0.6544
P0232	Selectivity increment	Fishery; Bin4; 1971 *	−0.0308	0.3714	−0.7587	0.6972
P0233	Selectivity increment	Fishery; Bin5; 1971 *	0.0085	0.3196	−0.6179	0.6350
P0234	Selectivity increment	Fishery; Bin6; 1971 *	−0.0046	0.3151	−0.6222	0.6129
P0235	Selectivity increment	Fishery; Bin7; 1971 *	−0.0485	0.3697	−0.7730	0.6761
P0236	Selectivity increment	Fishery; Bin8; 1971 *	0.0021	0.4013	−0.7845	0.7887
P0237	Selectivity increment	Fishery; Bin9; 1971 *	0.0038	0.4016	−0.7833	0.7910

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0238	Selectivity increment	Fishery; Bin10; 1971 *	0.0043	0.4080	−0.7953	0.8040
P0239	Selectivity increment	Fishery; Bin11; 1971 *	0.0104	0.4051	−0.7835	0.8044
P0240	Selectivity increment	Fishery; Bin12; 1971 *	0.0224	0.3007	−0.5671	0.6118
P0241	Selectivity increment	Fishery; Bin1; 1972 *	−0.1540	0.4171	−0.9715	0.6635
P0242	Selectivity increment	Fishery; Bin2; 1972 *	0.1905	0.3966	−0.5868	0.9678
P0243	Selectivity increment	Fishery; Bin3; 1972 *	0.0918	0.3791	−0.6513	0.8349
P0244	Selectivity increment	Fishery; Bin4; 1972 *	0.0742	0.3605	−0.6324	0.7808
P0245	Selectivity increment	Fishery; Bin5; 1972 *	−0.0186	0.3073	−0.6208	0.5836
P0246	Selectivity increment	Fishery; Bin6; 1972 *	−0.0055	0.2955	−0.5847	0.5738
P0247	Selectivity increment	Fishery; Bin7; 1972 *	−0.0047	0.3201	−0.6322	0.6227
P0248	Selectivity increment	Fishery; Bin8; 1972 *	−0.0323	0.4015	−0.8192	0.7546
P0249	Selectivity increment	Fishery; Bin9; 1972 *	−0.0229	0.4016	−0.8100	0.7641
P0250	Selectivity increment	Fishery; Bin10; 1972 *	−0.0315	0.4079	−0.8311	0.7680
P0251	Selectivity increment	Fishery; Bin11; 1972 *	−0.0396	0.4049	−0.8332	0.7540
P0252	Selectivity increment	Fishery; Bin12; 1972 *	−0.0473	0.3006	−0.6364	0.5419
P0253	Selectivity increment	Fishery; Bin1; 1973 *	−0.1854	0.4165	−1.0018	0.6309
P0254	Selectivity increment	Fishery; Bin2; 1973 *	0.1006	0.3966	−0.6767	0.8779
P0255	Selectivity increment	Fishery; Bin3; 1973 *	0.0425	0.3793	−0.7010	0.7860
P0256	Selectivity increment	Fishery; Bin4; 1973 *	0.0386	0.3526	−0.6525	0.7298
P0257	Selectivity increment	Fishery; Bin5; 1973 *	0.0410	0.3027	−0.5522	0.6342
P0258	Selectivity increment	Fishery; Bin6; 1973 *	0.0434	0.2924	−0.5296	0.6165
P0259	Selectivity increment	Fishery; Bin7; 1973 *	0.0391	0.3170	−0.5822	0.6604
P0260	Selectivity increment	Fishery; Bin8; 1973 *	0.0045	0.3969	−0.7735	0.7825
P0261	Selectivity increment	Fishery; Bin9; 1973 *	−0.0153	0.3987	−0.7967	0.7660
P0262	Selectivity increment	Fishery; Bin10; 1973 *	−0.0262	0.4063	−0.8225	0.7701
P0263	Selectivity increment	Fishery; Bin11; 1973 *	−0.0364	0.4038	−0.8279	0.7550
P0264	Selectivity increment	Fishery; Bin12; 1973 *	−0.0463	0.3011	−0.6364	0.5438
P0265	Selectivity increment	Fishery; Bin1; 1974 *	−0.1610	0.4155	−0.9753	0.6534
P0266	Selectivity increment	Fishery; Bin2; 1974 *	0.1384	0.3933	−0.6324	0.9092
P0267	Selectivity increment	Fishery; Bin3; 1974 *	0.1528	0.3689	−0.5702	0.8759
P0268	Selectivity increment	Fishery; Bin4; 1974 *	0.0118	0.3473	−0.6690	0.6926
P0269	Selectivity increment	Fishery; Bin5; 1974 *	0.0042	0.3027	−0.5891	0.5975
P0270	Selectivity increment	Fishery; Bin6; 1974 *	−0.0021	0.2923	−0.5750	0.5709
P0271	Selectivity increment	Fishery; Bin7; 1974 *	−0.0010	0.3158	−0.6200	0.6180

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0272	Selectivity increment	Fishery; Bin8; 1974 *	−0.0203	0.3456	−0.6978	0.6571
P0273	Selectivity increment	Fishery; Bin9; 1974 *	−0.0200	0.3422	−0.6907	0.6507
P0274	Selectivity increment	Fishery; Bin10; 1974 *	−0.0257	0.4021	−0.8137	0.7624
P0275	Selectivity increment	Fishery; Bin11; 1974 *	−0.0326	0.4002	−0.8171	0.7518
P0276	Selectivity increment	Fishery; Bin12; 1974 *	−0.0446	0.3017	−0.6360	0.5468
P0277	Selectivity increment	Fishery; Bin1; 1975 *	−0.1574	0.4148	−0.9705	0.6557
P0278	Selectivity increment	Fishery; Bin2; 1975 *	−0.1858	0.3879	−0.9460	0.5745
P0279	Selectivity increment	Fishery; Bin3; 1975 *	0.1323	0.3078	−0.4710	0.7357
P0280	Selectivity increment	Fishery; Bin4; 1975 *	0.0781	0.2842	−0.4789	0.6351
P0281	Selectivity increment	Fishery; Bin5; 1975 *	0.0430	0.2808	−0.5074	0.5933
P0282	Selectivity increment	Fishery; Bin6; 1975 *	0.0290	0.2859	−0.5313	0.5893
P0283	Selectivity increment	Fishery; Bin7; 1975 *	0.0231	0.3077	−0.5801	0.6262
P0284	Selectivity increment	Fishery; Bin8; 1975 *	0.0028	0.3377	−0.6589	0.6646
P0285	Selectivity increment	Fishery; Bin9; 1975 *	0.0035	0.3448	−0.6722	0.6792
P0286	Selectivity increment	Fishery; Bin10; 1975 *	0.0104	0.3139	−0.6049	0.6257
P0287	Selectivity increment	Fishery; Bin11; 1975 *	0.0105	0.2910	−0.5598	0.5807
P0288	Selectivity increment	Fishery; Bin12; 1975 *	0.0104	0.2783	−0.5351	0.5558
P0289	Selectivity increment	Fishery; Bin1; 1976 *	−0.1591	0.4145	−0.9714	0.6533
P0290	Selectivity increment	Fishery; Bin2; 1976 *	−0.0964	0.3884	−0.8577	0.6650
P0291	Selectivity increment	Fishery; Bin3; 1976 *	−0.0776	0.3491	−0.7618	0.6066
P0292	Selectivity increment	Fishery; Bin4; 1976 *	0.0621	0.3017	−0.5292	0.6534
P0293	Selectivity increment	Fishery; Bin5; 1976 *	0.0496	0.2832	−0.5055	0.6048
P0294	Selectivity increment	Fishery; Bin6; 1976 *	0.0379	0.2801	−0.5111	0.5870
P0295	Selectivity increment	Fishery; Bin7; 1976 *	0.0310	0.2819	−0.5214	0.5835
P0296	Selectivity increment	Fishery; Bin8; 1976 *	0.0269	0.2950	−0.5513	0.6051
P0297	Selectivity increment	Fishery; Bin9; 1976 *	0.0252	0.3416	−0.6443	0.6947
P0298	Selectivity increment	Fishery; Bin10; 1976 *	0.0334	0.3214	−0.5965	0.6633
P0299	Selectivity increment	Fishery; Bin11; 1976 *	0.0339	0.2922	−0.5388	0.6066
P0300	Selectivity increment	Fishery; Bin12; 1976 *	0.0329	0.2785	−0.5129	0.5787
P0301	Selectivity increment	Fishery; Bin1; 1977 *	−0.1490	0.4140	−0.9604	0.6624
P0302	Selectivity increment	Fishery; Bin2; 1977 *	−0.0115	0.3875	−0.7709	0.7479
P0303	Selectivity increment	Fishery; Bin3; 1977 *	−0.1853	0.3694	−0.9094	0.5387
P0304	Selectivity increment	Fishery; Bin4; 1977 *	−0.0719	0.3558	−0.7692	0.6254
P0305	Selectivity increment	Fishery; Bin5; 1977 *	0.0195	0.3047	−0.5777	0.6167

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0306	Selectivity increment	Fishery; Bin6; 1977 *	0.0338	0.2861	−0.5269	0.5945
P0307	Selectivity increment	Fishery; Bin7; 1977 *	0.0379	0.2880	−0.5266	0.6024
P0308	Selectivity increment	Fishery; Bin8; 1977 *	0.0395	0.3108	−0.5697	0.6487
P0309	Selectivity increment	Fishery; Bin9; 1977 *	0.0514	0.3946	−0.7220	0.8247
P0310	Selectivity increment	Fishery; Bin10; 1977 *	0.0697	0.3962	−0.7069	0.8463
P0311	Selectivity increment	Fishery; Bin11; 1977 *	0.0816	0.3251	−0.5556	0.7189
P0312	Selectivity increment	Fishery; Bin12; 1977 *	0.0844	0.2885	−0.4811	0.6498
P0313	Selectivity increment	Fishery; Bin1; 1978 *	−0.1935	0.4131	−1.0032	0.6162
P0314	Selectivity increment	Fishery; Bin2; 1978 *	−0.1580	0.3803	−0.9035	0.5874
P0315	Selectivity increment	Fishery; Bin3; 1978 *	−0.0660	0.3645	−0.7803	0.6483
P0316	Selectivity increment	Fishery; Bin4; 1978 *	−0.0381	0.3640	−0.7515	0.6753
P0317	Selectivity increment	Fishery; Bin5; 1978 *	0.0269	0.3143	−0.5891	0.6429
P0318	Selectivity increment	Fishery; Bin6; 1978 *	0.0307	0.3083	−0.5736	0.6349
P0319	Selectivity increment	Fishery; Bin7; 1978 *	0.0457	0.3154	−0.5725	0.6639
P0320	Selectivity increment	Fishery; Bin8; 1978 *	0.0496	0.3259	−0.5891	0.6883
P0321	Selectivity increment	Fishery; Bin9; 1978 *	0.0673	0.3989	−0.7144	0.8491
P0322	Selectivity increment	Fishery; Bin10; 1978 *	0.0741	0.3989	−0.7076	0.8559
P0323	Selectivity increment	Fishery; Bin11; 1978 *	0.0807	0.3341	−0.5741	0.7354
P0324	Selectivity increment	Fishery; Bin12; 1978 *	0.0808	0.2890	−0.4856	0.6472
P0325	Selectivity increment	Fishery; Bin1; 1979 *	−0.1527	0.4072	−0.9509	0.6455
P0326	Selectivity increment	Fishery; Bin2; 1979 *	−0.3807	0.3608	−1.0879	0.3265
P0327	Selectivity increment	Fishery; Bin3; 1979 *	−0.1744	0.3505	−0.8614	0.5126
P0328	Selectivity increment	Fishery; Bin4; 1979 *	−0.0494	0.3568	−0.7488	0.6499
P0329	Selectivity increment	Fishery; Bin5; 1979 *	0.0853	0.3491	−0.5990	0.7695
P0330	Selectivity increment	Fishery; Bin6; 1979 *	0.1164	0.3292	−0.5289	0.7617
P0331	Selectivity increment	Fishery; Bin7; 1979 *	0.0994	0.3399	−0.5669	0.7656
P0332	Selectivity increment	Fishery; Bin8; 1979 *	0.1032	0.3687	−0.6194	0.8257
P0333	Selectivity increment	Fishery; Bin9; 1979 *	0.1059	0.4008	−0.6796	0.8914
P0334	Selectivity increment	Fishery; Bin10; 1979 *	0.0896	0.3983	−0.6910	0.8703
P0335	Selectivity increment	Fishery; Bin11; 1979 *	0.0792	0.3286	−0.5649	0.7232
P0336	Selectivity increment	Fishery; Bin12; 1979 *	0.0784	0.2878	−0.4856	0.6424
P0337	Selectivity increment	Fishery; Bin1; 1980 *	−0.1661	0.4085	−0.9667	0.6344
P0338	Selectivity increment	Fishery; Bin2; 1980 *	−0.4732	0.3535	−1.1660	0.2197
P0339	Selectivity increment	Fishery; Bin3; 1980 *	−0.2547	0.3436	−0.9282	0.4188

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0340	Selectivity increment	Fishery; Bin4; 1980 *	0.0750	0.3564	−0.6234	0.7734
P0341	Selectivity increment	Fishery; Bin5; 1980 *	0.1095	0.3585	−0.5930	0.8121
P0342	Selectivity increment	Fishery; Bin6; 1980 *	0.1351	0.3138	−0.4800	0.7502
P0343	Selectivity increment	Fishery; Bin7; 1980 *	0.1332	0.3106	−0.4757	0.7420
P0344	Selectivity increment	Fishery; Bin8; 1980 *	0.1202	0.3686	−0.6023	0.8427
P0345	Selectivity increment	Fishery; Bin9; 1980 *	0.0907	0.3988	−0.6909	0.8722
P0346	Selectivity increment	Fishery; Bin10; 1980 *	0.0826	0.3937	−0.6891	0.8542
P0347	Selectivity increment	Fishery; Bin11; 1980 *	0.0751	0.3175	−0.5472	0.6975
P0348	Selectivity increment	Fishery; Bin12; 1980 *	0.0726	0.2867	−0.4893	0.6345
P0349	Selectivity increment	Fishery; Bin1; 1981 *	−0.1370	0.4123	−0.9450	0.6710
P0350	Selectivity increment	Fishery; Bin2; 1981 *	−0.2530	0.3700	−0.9782	0.4722
P0351	Selectivity increment	Fishery; Bin3; 1981 *	−0.2579	0.3394	−0.9230	0.4072
P0352	Selectivity increment	Fishery; Bin4; 1981 *	0.0107	0.3493	−0.6739	0.6953
P0353	Selectivity increment	Fishery; Bin5; 1981 *	0.1230	0.3471	−0.5573	0.8033
P0354	Selectivity increment	Fishery; Bin6; 1981 *	0.0783	0.3012	−0.5121	0.6686
P0355	Selectivity increment	Fishery; Bin7; 1981 *	0.0847	0.2895	−0.4827	0.6522
P0356	Selectivity increment	Fishery; Bin8; 1981 *	0.0882	0.3089	−0.5172	0.6936
P0357	Selectivity increment	Fishery; Bin9; 1981 *	0.0812	0.3921	−0.6872	0.8497
P0358	Selectivity increment	Fishery; Bin10; 1981 *	0.0626	0.3238	−0.5721	0.6973
P0359	Selectivity increment	Fishery; Bin11; 1981 *	0.0595	0.2952	−0.5191	0.6382
P0360	Selectivity increment	Fishery; Bin12; 1981 *	0.0596	0.2817	−0.4925	0.6117
P0361	Selectivity increment	Fishery; Bin1; 1982 *	−0.1040	0.4154	−0.9182	0.7102
P0362	Selectivity increment	Fishery; Bin2; 1982 *	−0.0931	0.3823	−0.8423	0.6562
P0363	Selectivity increment	Fishery; Bin3; 1982 *	−0.3008	0.3536	−0.9937	0.3922
P0364	Selectivity increment	Fishery; Bin4; 1982 *	−0.0092	0.3403	−0.6762	0.6579
P0365	Selectivity increment	Fishery; Bin5; 1982 *	0.0660	0.3012	−0.5244	0.6565
P0366	Selectivity increment	Fishery; Bin6; 1982 *	0.0694	0.2805	−0.4804	0.6193
P0367	Selectivity increment	Fishery; Bin7; 1982 *	0.0670	0.2772	−0.4763	0.6104
P0368	Selectivity increment	Fishery; Bin8; 1982 *	0.0685	0.2771	−0.4746	0.6116
P0369	Selectivity increment	Fishery; Bin9; 1982 *	0.0662	0.2794	−0.4813	0.6137
P0370	Selectivity increment	Fishery; Bin10; 1982 *	0.0600	0.2812	−0.4911	0.6111
P0371	Selectivity increment	Fishery; Bin11; 1982 *	0.0560	0.2836	−0.5000	0.6119
P0372	Selectivity increment	Fishery; Bin12; 1982 *	0.0538	0.2777	−0.4905	0.5981
P0373	Selectivity increment	Fishery; Bin1; 1983 *	−0.1223	0.4165	−0.9386	0.6940

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0374	Selectivity increment	Fishery; Bin2; 1983 *	0.1032	0.3892	−0.6597	0.8660
P0375	Selectivity increment	Fishery; Bin3; 1983 *	−0.0970	0.3645	−0.8113	0.6174
P0376	Selectivity increment	Fishery; Bin4; 1983 *	−0.1890	0.3452	−0.8656	0.4876
P0377	Selectivity increment	Fishery; Bin5; 1983 *	−0.0624	0.3207	−0.6909	0.5661
P0378	Selectivity increment	Fishery; Bin6; 1983 *	0.0676	0.2926	−0.5059	0.6410
P0379	Selectivity increment	Fishery; Bin7; 1983 *	0.0552	0.2786	−0.4908	0.6012
P0380	Selectivity increment	Fishery; Bin8; 1983 *	0.0486	0.2766	−0.4935	0.5906
P0381	Selectivity increment	Fishery; Bin9; 1983 *	0.0482	0.2775	−0.4956	0.5921
P0382	Selectivity increment	Fishery; Bin10; 1983 *	0.0497	0.2797	−0.4985	0.5978
P0383	Selectivity increment	Fishery; Bin11; 1983 *	0.0494	0.2820	−0.5033	0.6021
P0384	Selectivity increment	Fishery; Bin12; 1983 *	0.0488	0.2760	−0.4922	0.5898
P0385	Selectivity increment	Fishery; Bin1; 1984 *	−0.1099	0.4185	−0.9301	0.7102
P0386	Selectivity increment	Fishery; Bin2; 1984 *	−0.1521	0.3905	−0.9174	0.6132
P0387	Selectivity increment	Fishery; Bin3; 1984 *	0.0489	0.3700	−0.6763	0.7742
P0388	Selectivity increment	Fishery; Bin4; 1984 *	−0.0874	0.3505	−0.7743	0.5996
P0389	Selectivity increment	Fishery; Bin5; 1984 *	0.0264	0.3403	−0.6405	0.6933
P0390	Selectivity increment	Fishery; Bin6; 1984 *	0.0367	0.2951	−0.5417	0.6151
P0391	Selectivity increment	Fishery; Bin7; 1984 *	0.0512	0.2785	−0.4946	0.5970
P0392	Selectivity increment	Fishery; Bin8; 1984 *	0.0411	0.2762	−0.5002	0.5824
P0393	Selectivity increment	Fishery; Bin9; 1984 *	0.0380	0.2771	−0.5051	0.5811
P0394	Selectivity increment	Fishery; Bin10; 1984 *	0.0360	0.2792	−0.5113	0.5833
P0395	Selectivity increment	Fishery; Bin11; 1984 *	0.0358	0.2814	−0.5158	0.5874
P0396	Selectivity increment	Fishery; Bin12; 1984 *	0.0354	0.2754	−0.5043	0.5751
P0397	Selectivity increment	Fishery; Bin1; 1985 *	−0.1272	0.4190	−0.9484	0.6940
P0398	Selectivity increment	Fishery; Bin2; 1985 *	0.0402	0.3949	−0.7338	0.8141
P0399	Selectivity increment	Fishery; Bin3; 1985 *	0.1307	0.3677	−0.5900	0.8513
P0400	Selectivity increment	Fishery; Bin4; 1985 *	−0.0902	0.3563	−0.7885	0.6081
P0401	Selectivity increment	Fishery; Bin5; 1985 *	−0.2409	0.3470	−0.9210	0.4392
P0402	Selectivity increment	Fishery; Bin6; 1985 *	0.0564	0.2947	−0.5213	0.6340
P0403	Selectivity increment	Fishery; Bin7; 1985 *	0.0519	0.2754	−0.4879	0.5917
P0404	Selectivity increment	Fishery; Bin8; 1985 *	0.0500	0.2751	−0.4892	0.5891
P0405	Selectivity increment	Fishery; Bin9; 1985 *	0.0345	0.2765	−0.5074	0.5765
P0406	Selectivity increment	Fishery; Bin10; 1985 *	0.0325	0.2789	−0.5141	0.5790
P0407	Selectivity increment	Fishery; Bin11; 1985 *	0.0318	0.2811	−0.5191	0.5827

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0408	Selectivity increment	Fishery; Bin12; 1985 *	0.0304	0.2749	−0.5084	0.5693
P0409	Selectivity increment	Fishery; Bin1; 1986 *	−0.1344	0.4196	−0.9569	0.6881
P0410	Selectivity increment	Fishery; Bin2; 1986 *	−0.0383	0.3972	−0.8168	0.7401
P0411	Selectivity increment	Fishery; Bin3; 1986 *	0.0606	0.3691	−0.6628	0.7840
P0412	Selectivity increment	Fishery; Bin4; 1986 *	0.1677	0.3543	−0.5267	0.8622
P0413	Selectivity increment	Fishery; Bin5; 1986 *	−0.0207	0.3550	−0.7164	0.6751
P0414	Selectivity increment	Fishery; Bin6; 1986 *	−0.0709	0.3319	−0.7214	0.5796
P0415	Selectivity increment	Fishery; Bin7; 1986 *	0.0143	0.2940	−0.5620	0.5905
P0416	Selectivity increment	Fishery; Bin8; 1986 *	−0.0173	0.2782	−0.5626	0.5279
P0417	Selectivity increment	Fishery; Bin9; 1986 *	0.0089	0.2789	−0.5378	0.5555
P0418	Selectivity increment	Fishery; Bin10; 1986 *	0.0072	0.2788	−0.5393	0.5537
P0419	Selectivity increment	Fishery; Bin11; 1986 *	0.0093	0.2807	−0.5408	0.5594
P0420	Selectivity increment	Fishery; Bin12; 1986 *	0.0136	0.2745	−0.5244	0.5517
P0421	Selectivity increment	Fishery; Bin1; 1987 *	−0.1333	0.4198	−0.9561	0.6895
P0422	Selectivity increment	Fishery; Bin2; 1987 *	−0.0846	0.4008	−0.8702	0.7009
P0423	Selectivity increment	Fishery; Bin3; 1987 *	0.1086	0.3713	−0.6192	0.8364
P0424	Selectivity increment	Fishery; Bin4; 1987 *	−0.1061	0.3582	−0.8082	0.5959
P0425	Selectivity increment	Fishery; Bin5; 1987 *	0.0218	0.3538	−0.6717	0.7153
P0426	Selectivity increment	Fishery; Bin6; 1987 *	−0.0462	0.3557	−0.7433	0.6509
P0427	Selectivity increment	Fishery; Bin7; 1987 *	−0.0955	0.3359	−0.7538	0.5628
P0428	Selectivity increment	Fishery; Bin8; 1987 *	0.1044	0.2962	−0.4762	0.6850
P0429	Selectivity increment	Fishery; Bin9; 1987 *	0.0984	0.2793	−0.4491	0.6459
P0430	Selectivity increment	Fishery; Bin10; 1987 *	0.0660	0.2781	−0.4791	0.6112
P0431	Selectivity increment	Fishery; Bin11; 1987 *	0.0418	0.2801	−0.5072	0.5908
P0432	Selectivity increment	Fishery; Bin12; 1987 *	0.0248	0.2743	−0.5127	0.5624
P0433	Selectivity increment	Fishery; Bin1; 1988 *	−0.1650	0.4198	−0.9878	0.6577
P0434	Selectivity increment	Fishery; Bin2; 1988 *	−0.1229	0.4025	−0.9118	0.6660
P0435	Selectivity increment	Fishery; Bin3; 1988 *	0.4321	0.3737	−0.3003	1.1645
P0436	Selectivity increment	Fishery; Bin4; 1988 *	0.0242	0.3611	−0.6835	0.7320
P0437	Selectivity increment	Fishery; Bin5; 1988 *	0.0879	0.3542	−0.6062	0.7820
P0438	Selectivity increment	Fishery; Bin6; 1988 *	−0.0565	0.3542	−0.7508	0.6377
P0439	Selectivity increment	Fishery; Bin7; 1988 *	0.1187	0.3368	−0.5414	0.7788
P0440	Selectivity increment	Fishery; Bin8; 1988 *	−0.0812	0.2935	−0.6565	0.4941
P0441	Selectivity increment	Fishery; Bin9; 1988 *	−0.0785	0.2772	−0.6218	0.4648

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0442	Selectivity increment	Fishery; Bin10; 1988 *	−0.0796	0.2776	−0.6236	0.4644
P0443	Selectivity increment	Fishery; Bin11; 1988 *	−0.0497	0.2829	−0.6042	0.5048
P0444	Selectivity increment	Fishery; Bin12; 1988 *	−0.0295	0.2753	−0.5690	0.5100
P0445	Selectivity increment	Fishery; Bin1; 1989 *	−0.1307	0.4196	−0.9531	0.6916
P0446	Selectivity increment	Fishery; Bin2; 1989 *	−0.0816	0.4021	−0.8696	0.7065
P0447	Selectivity increment	Fishery; Bin3; 1989 *	−0.3948	0.3821	−1.1437	0.3540
P0448	Selectivity increment	Fishery; Bin4; 1989 *	0.1355	0.3658	−0.5814	0.8525
P0449	Selectivity increment	Fishery; Bin5; 1989 *	0.0220	0.3542	−0.6723	0.7163
P0450	Selectivity increment	Fishery; Bin6; 1989 *	0.0546	0.3531	−0.6374	0.7466
P0451	Selectivity increment	Fishery; Bin7; 1989 *	0.1293	0.2978	−0.4543	0.7130
P0452	Selectivity increment	Fishery; Bin8; 1989 *	0.1041	0.2772	−0.4392	0.6473
P0453	Selectivity increment	Fishery; Bin9; 1989 *	0.0309	0.2754	−0.5088	0.5706
P0454	Selectivity increment	Fishery; Bin10; 1989 *	0.0641	0.2773	−0.4795	0.6076
P0455	Selectivity increment	Fishery; Bin11; 1989 *	0.0445	0.2815	−0.5072	0.5962
P0456	Selectivity increment	Fishery; Bin12; 1989 *	0.0221	0.2747	−0.5163	0.5605
P0457	Selectivity increment	Fishery; Bin1; 1990 *	−0.1246	0.4192	−0.9462	0.6971
P0458	Selectivity increment	Fishery; Bin2; 1990 *	−0.0556	0.3998	−0.8392	0.7279
P0459	Selectivity increment	Fishery; Bin3; 1990 *	−0.3454	0.3826	−1.0953	0.4046
P0460	Selectivity increment	Fishery; Bin4; 1990 *	0.0258	0.3704	−0.7002	0.7518
P0461	Selectivity increment	Fishery; Bin5; 1990 *	0.1451	0.3528	−0.5464	0.8366
P0462	Selectivity increment	Fishery; Bin6; 1990 *	0.0542	0.3493	−0.6303	0.7387
P0463	Selectivity increment	Fishery; Bin7; 1990 *	−0.0416	0.3272	−0.6830	0.5997
P0464	Selectivity increment	Fishery; Bin8; 1990 *	0.0504	0.2894	−0.5169	0.6176
P0465	Selectivity increment	Fishery; Bin9; 1990 *	0.0621	0.2804	−0.4875	0.6116
P0466	Selectivity increment	Fishery; Bin10; 1990 *	0.0002	0.2925	−0.5731	0.5736
P0467	Selectivity increment	Fishery; Bin11; 1990 *	0.1328	0.2978	−0.4508	0.7164
P0468	Selectivity increment	Fishery; Bin12; 1990 *	0.0966	0.2719	−0.4364	0.6295
P0469	Selectivity increment	Fishery; Bin1; 1991 *	−0.0571	0.4181	−0.8765	0.7624
P0470	Selectivity increment	Fishery; Bin2; 1991 *	−0.0210	0.3915	−0.7883	0.7462
P0471	Selectivity increment	Fishery; Bin3; 1991 *	−0.3596	0.3748	−1.0942	0.3750
P0472	Selectivity increment	Fishery; Bin4; 1991 *	−0.2984	0.3583	−1.0006	0.4038
P0473	Selectivity increment	Fishery; Bin5; 1991 *	−0.1882	0.3484	−0.8710	0.4946
P0474	Selectivity increment	Fishery; Bin6; 1991 *	−0.0522	0.3230	−0.6853	0.5809
P0475	Selectivity increment	Fishery; Bin7; 1991 *	0.1413	0.3233	−0.4924	0.7750

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0476	Selectivity increment	Fishery; Bin8; 1991 *	−0.0373	0.2924	−0.6104	0.5359
P0477	Selectivity increment	Fishery; Bin9; 1991 *	0.1423	0.2774	−0.4014	0.6859
P0478	Selectivity increment	Fishery; Bin10; 1991 *	0.0714	0.2959	−0.5085	0.6513
P0479	Selectivity increment	Fishery; Bin11; 1991 *	0.3360	0.2985	−0.2491	0.9211
P0480	Selectivity increment	Fishery; Bin12; 1991 *	0.3228	0.2650	−0.1966	0.8421
P0481	Selectivity increment	Fishery; Bin1; 1992 *	−0.1291	0.4153	−0.9430	0.6848
P0482	Selectivity increment	Fishery; Bin2; 1992 *	−0.0575	0.3344	−0.7129	0.5978
P0483	Selectivity increment	Fishery; Bin3; 1992 *	0.5222	0.3061	−0.0779	1.1222
P0484	Selectivity increment	Fishery; Bin4; 1992 *	−0.2344	0.3156	−0.8530	0.3842
P0485	Selectivity increment	Fishery; Bin5; 1992 *	−0.3277	0.3180	−0.9509	0.2955
P0486	Selectivity increment	Fishery; Bin6; 1992 *	−0.0479	0.3127	−0.6607	0.5649
P0487	Selectivity increment	Fishery; Bin7; 1992 *	−0.0629	0.2927	−0.6365	0.5108
P0488	Selectivity increment	Fishery; Bin8; 1992 *	0.2020	0.2872	−0.3608	0.7648
P0489	Selectivity increment	Fishery; Bin9; 1992 *	0.0885	0.2802	−0.4607	0.6377
P0490	Selectivity increment	Fishery; Bin10; 1992 *	0.2149	0.2878	−0.3491	0.7790
P0491	Selectivity increment	Fishery; Bin11; 1992 *	−0.1178	0.2882	−0.6827	0.4471
P0492	Selectivity increment	Fishery; Bin12; 1992 *	−0.0503	0.2607	−0.5612	0.4607
P0493	Selectivity increment	Fishery; Bin1; 1993 *	−0.1187	0.4147	−0.9315	0.6940
P0494	Selectivity increment	Fishery; Bin2; 1993 *	−0.4719	0.3522	−1.1621	0.2184
P0495	Selectivity increment	Fishery; Bin3; 1993 *	−0.0290	0.2736	−0.5654	0.5073
P0496	Selectivity increment	Fishery; Bin4; 1993 *	0.8219	0.2800	0.2731	1.3708
P0497	Selectivity increment	Fishery; Bin5; 1993 *	0.0377	0.3059	−0.5619	0.6373
P0498	Selectivity increment	Fishery; Bin6; 1993 *	0.0368	0.3239	−0.5980	0.6717
P0499	Selectivity increment	Fishery; Bin7; 1993 *	0.0738	0.3239	−0.5609	0.7086
P0500	Selectivity increment	Fishery; Bin8; 1993 *	−0.0135	0.2905	−0.5830	0.5559
P0501	Selectivity increment	Fishery; Bin9; 1993 *	−0.0312	0.2806	−0.5812	0.5188
P0502	Selectivity increment	Fishery; Bin10; 1993 *	−0.0706	0.2792	−0.6178	0.4766
P0503	Selectivity increment	Fishery; Bin11; 1993 *	−0.1520	0.3012	−0.7423	0.4384
P0504	Selectivity increment	Fishery; Bin12; 1993 *	−0.0833	0.2738	−0.6200	0.4534
P0505	Selectivity increment	Fishery; Bin1; 1994 *	−0.0143	0.4166	−0.8309	0.8023
P0506	Selectivity increment	Fishery; Bin2; 1994 *	0.0152	0.3582	−0.6868	0.7172
P0507	Selectivity increment	Fishery; Bin3; 1994 *	−0.7774	0.3087	−1.3825	−0.1722
P0508	Selectivity increment	Fishery; Bin4; 1994 *	−0.3070	0.2656	−0.8275	0.2136
P0509	Selectivity increment	Fishery; Bin5; 1994 *	0.9075	0.2652	0.3877	1.4273

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0510	Selectivity increment	Fishery; Bin6; 1994 *	0.2876	0.3068	−0.3137	0.8890
P0511	Selectivity increment	Fishery; Bin7; 1994 *	0.0218	0.3155	−0.5966	0.6402
P0512	Selectivity increment	Fishery; Bin8; 1994 *	−0.0271	0.2831	−0.5820	0.5277
P0513	Selectivity increment	Fishery; Bin9; 1994 *	−0.0449	0.2690	−0.5721	0.4824
P0514	Selectivity increment	Fishery; Bin10; 1994 *	−0.0192	0.2727	−0.5538	0.5153
P0515	Selectivity increment	Fishery; Bin11; 1994 *	0.0142	0.2870	−0.5483	0.5767
P0516	Selectivity increment	Fishery; Bin12; 1994 *	−0.0564	0.2782	−0.6018	0.4889
P0517	Selectivity increment	Fishery; Bin1; 1995 *	0.0094	0.4170	−0.8078	0.8267
P0518	Selectivity increment	Fishery; Bin2; 1995 *	0.1923	0.3679	−0.5288	0.9134
P0519	Selectivity increment	Fishery; Bin3; 1995 *	−0.2946	0.3221	−0.9259	0.3368
P0520	Selectivity increment	Fishery; Bin4; 1995 *	−0.7007	0.2923	−1.2736	−0.1279
P0521	Selectivity increment	Fishery; Bin5; 1995 *	−0.3795	0.2662	−0.9013	0.1423
P0522	Selectivity increment	Fishery; Bin6; 1995 *	0.4455	0.2661	−0.0760	0.9671
P0523	Selectivity increment	Fishery; Bin7; 1995 *	0.1832	0.2782	−0.3621	0.7285
P0524	Selectivity increment	Fishery; Bin8; 1995 *	0.1784	0.2707	−0.3522	0.7090
P0525	Selectivity increment	Fishery; Bin9; 1995 *	0.1368	0.2714	−0.3951	0.6687
P0526	Selectivity increment	Fishery; Bin10; 1995 *	0.1109	0.2724	−0.4231	0.6449
P0527	Selectivity increment	Fishery; Bin11; 1995 *	0.0758	0.2748	−0.4628	0.6143
P0528	Selectivity increment	Fishery; Bin12; 1995 *	0.0424	0.2681	−0.4831	0.5680
P0529	Selectivity increment	Fishery; Bin1; 1996 *	−0.0097	0.4173	−0.8277	0.8083
P0530	Selectivity increment	Fishery; Bin2; 1996 *	0.5075	0.3722	−0.2219	1.2369
P0531	Selectivity increment	Fishery; Bin3; 1996 *	0.2794	0.3348	−0.3768	0.9357
P0532	Selectivity increment	Fishery; Bin4; 1996 *	−0.7854	0.3054	−1.3840	−0.1867
P0533	Selectivity increment	Fishery; Bin5; 1996 *	−0.7408	0.2866	−1.3026	−0.1790
P0534	Selectivity increment	Fishery; Bin6; 1996 *	−0.3569	0.2674	−0.8811	0.1672
P0535	Selectivity increment	Fishery; Bin7; 1996 *	0.2781	0.2585	−0.2285	0.7847
P0536	Selectivity increment	Fishery; Bin8; 1996 *	0.3081	0.2639	−0.2091	0.8253
P0537	Selectivity increment	Fishery; Bin9; 1996 *	0.2215	0.2701	−0.3080	0.7510
P0538	Selectivity increment	Fishery; Bin10; 1996 *	0.1476	0.2737	−0.3889	0.6841
P0539	Selectivity increment	Fishery; Bin11; 1996 *	0.0906	0.2752	−0.4489	0.6301
P0540	Selectivity increment	Fishery; Bin12; 1996 *	0.0601	0.2678	−0.4648	0.5849
P0541	Selectivity increment	Fishery; Bin1; 1997 *	−0.1094	0.4177	−0.9282	0.7093
P0542	Selectivity increment	Fishery; Bin2; 1997 *	0.3323	0.3622	−0.3776	1.0422
P0543	Selectivity increment	Fishery; Bin3; 1997 *	0.1555	0.3403	−0.5114	0.8224

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0544	Selectivity increment	Fishery; Bin4; 1997 *	0.6559	0.3059	0.0563	1.2556
P0545	Selectivity increment	Fishery; Bin5; 1997 *	−0.0012	0.2806	−0.5511	0.5488
P0546	Selectivity increment	Fishery; Bin6; 1997 *	−0.3984	0.2751	−0.9376	0.1409
P0547	Selectivity increment	Fishery; Bin7; 1997 *	−0.4036	0.2694	−0.9315	0.1243
P0548	Selectivity increment	Fishery; Bin8; 1997 *	−0.2889	0.2735	−0.8249	0.2471
P0549	Selectivity increment	Fishery; Bin9; 1997 *	−0.0227	0.2787	−0.5690	0.5235
P0550	Selectivity increment	Fishery; Bin10; 1997 *	0.0186	0.2746	−0.5197	0.5569
P0551	Selectivity increment	Fishery; Bin11; 1997 *	0.0284	0.2758	−0.5121	0.5689
P0552	Selectivity increment	Fishery; Bin12; 1997 *	0.0335	0.2671	−0.4900	0.5570
P0553	Selectivity increment	Fishery; Bin1; 1998 *	−0.0772	0.4183	−0.8971	0.7427
P0554	Selectivity increment	Fishery; Bin2; 1998 *	−0.3764	0.3344	−1.0318	0.2791
P0555	Selectivity increment	Fishery; Bin3; 1998 *	0.2778	0.3214	−0.3521	0.9077
P0556	Selectivity increment	Fishery; Bin4; 1998 *	0.2308	0.3062	−0.3693	0.8309
P0557	Selectivity increment	Fishery; Bin5; 1998 *	0.1763	0.2739	−0.3605	0.7132
P0558	Selectivity increment	Fishery; Bin6; 1998 *	0.2773	0.2583	−0.2290	0.7837
P0559	Selectivity increment	Fishery; Bin7; 1998 *	−0.2341	0.2704	−0.7642	0.2960
P0560	Selectivity increment	Fishery; Bin8; 1998 *	−0.0717	0.2888	−0.6378	0.4944
P0561	Selectivity increment	Fishery; Bin9; 1998 *	−0.0713	0.2812	−0.6223	0.4798
P0562	Selectivity increment	Fishery; Bin10; 1998 *	−0.0478	0.2738	−0.5845	0.4889
P0563	Selectivity increment	Fishery; Bin11; 1998 *	−0.0464	0.2758	−0.5870	0.4942
P0564	Selectivity increment	Fishery; Bin12; 1998 *	−0.0374	0.2688	−0.5643	0.4895
P0565	Selectivity increment	Fishery; Bin1; 1999 *	−0.1247	0.4186	−0.9452	0.6958
P0566	Selectivity increment	Fishery; Bin2; 1999 *	−0.3584	0.3509	−1.0462	0.3294
P0567	Selectivity increment	Fishery; Bin3; 1999 *	0.8134	0.2823	0.2601	1.3666
P0568	Selectivity increment	Fishery; Bin4; 1999 *	0.4525	0.2907	−0.1172	1.0221
P0569	Selectivity increment	Fishery; Bin5; 1999 *	0.1329	0.2861	−0.4279	0.6938
P0570	Selectivity increment	Fishery; Bin6; 1999 *	−0.2625	0.2614	−0.7749	0.2499
P0571	Selectivity increment	Fishery; Bin7; 1999 *	0.2338	0.2526	−0.2613	0.7289
P0572	Selectivity increment	Fishery; Bin8; 1999 *	−0.1112	0.2746	−0.6493	0.4269
P0573	Selectivity increment	Fishery; Bin9; 1999 *	−0.2599	0.2671	−0.7833	0.2636
P0574	Selectivity increment	Fishery; Bin10; 1999 *	−0.2373	0.2691	−0.7648	0.2901
P0575	Selectivity increment	Fishery; Bin11; 1999 *	−0.1704	0.2922	−0.7431	0.4023
P0576	Selectivity increment	Fishery; Bin12; 1999 *	−0.1081	0.2841	−0.6649	0.4487
P0577	Selectivity increment	Fishery; Bin1; 2000 *	−0.0558	0.4185	−0.8761	0.7645

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0578	Selectivity increment	Fishery; Bin2; 2000 *	−0.1443	0.3623	−0.8544	0.5658
P0579	Selectivity increment	Fishery; Bin3; 2000 *	−0.5590	0.2797	−1.1072	−0.0108
P0580	Selectivity increment	Fishery; Bin4; 2000 *	0.2946	0.2566	−0.2083	0.7976
P0581	Selectivity increment	Fishery; Bin5; 2000 *	0.4186	0.2725	−0.1155	0.9528
P0582	Selectivity increment	Fishery; Bin6; 2000 *	−0.0055	0.2800	−0.5544	0.5433
P0583	Selectivity increment	Fishery; Bin7; 2000 *	−0.0436	0.2605	−0.5542	0.4669
P0584	Selectivity increment	Fishery; Bin8; 2000 *	0.2266	0.2481	−0.2597	0.7129
P0585	Selectivity increment	Fishery; Bin9; 2000 *	0.0273	0.2558	−0.4741	0.5287
P0586	Selectivity increment	Fishery; Bin10; 2000 *	−0.0436	0.2647	−0.5624	0.4753
P0587	Selectivity increment	Fishery; Bin11; 2000 *	−0.0517	0.2881	−0.6163	0.5129
P0588	Selectivity increment	Fishery; Bin12; 2000 *	−0.0637	0.2858	−0.6239	0.4965
P0589	Selectivity increment	Fishery; Bin1; 2001 *	−0.0146	0.4182	−0.8342	0.8051
P0590	Selectivity increment	Fishery; Bin2; 2001 *	−0.2128	0.3639	−0.9260	0.5005
P0591	Selectivity increment	Fishery; Bin3; 2001 *	−0.4501	0.3107	−1.0591	0.1588
P0592	Selectivity increment	Fishery; Bin4; 2001 *	−0.3103	0.2707	−0.8409	0.2202
P0593	Selectivity increment	Fishery; Bin5; 2001 *	0.2544	0.2536	−0.2428	0.7515
P0594	Selectivity increment	Fishery; Bin6; 2001 *	0.6106	0.2699	0.0815	1.1397
P0595	Selectivity increment	Fishery; Bin7; 2001 *	0.1237	0.2651	−0.3958	0.6432
P0596	Selectivity increment	Fishery; Bin8; 2001 *	−0.1064	0.2514	−0.5990	0.3863
P0597	Selectivity increment	Fishery; Bin9; 2001 *	0.0296	0.2532	−0.4666	0.5259
P0598	Selectivity increment	Fishery; Bin10; 2001 *	0.0467	0.2626	−0.4679	0.5614
P0599	Selectivity increment	Fishery; Bin11; 2001 *	0.0368	0.2697	−0.4918	0.5654
P0600	Selectivity increment	Fishery; Bin12; 2001 *	−0.0076	0.2700	−0.5368	0.5215
P0601	Selectivity increment	Fishery; Bin1; 2002 *	−0.0584	0.4185	−0.8785	0.7618
P0602	Selectivity increment	Fishery; Bin2; 2002 *	0.1591	0.3560	−0.5386	0.8567
P0603	Selectivity increment	Fishery; Bin3; 2002 *	0.0693	0.3069	−0.5323	0.6709
P0604	Selectivity increment	Fishery; Bin4; 2002 *	−0.0089	0.2773	−0.5524	0.5345
P0605	Selectivity increment	Fishery; Bin5; 2002 *	−0.2313	0.2558	−0.7326	0.2699
P0606	Selectivity increment	Fishery; Bin6; 2002 *	0.0666	0.2451	−0.4137	0.5470
P0607	Selectivity increment	Fishery; Bin7; 2002 *	0.0798	0.2495	−0.4092	0.5689
P0608	Selectivity increment	Fishery; Bin8; 2002 *	0.0548	0.2547	−0.4443	0.5539
P0609	Selectivity increment	Fishery; Bin9; 2002 *	0.0383	0.2538	−0.4592	0.5357
P0610	Selectivity increment	Fishery; Bin10; 2002 *	−0.0247	0.2576	−0.5297	0.4802
P0611	Selectivity increment	Fishery; Bin11; 2002 *	−0.0785	0.2672	−0.6022	0.4452

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0612	Selectivity increment	Fishery; Bin12; 2002 *	−0.0660	0.2655	−0.5864	0.4544
P0613	Selectivity increment	Fishery; Bin1; 2003 *	−0.0709	0.4187	−0.8916	0.7499
P0614	Selectivity increment	Fishery; Bin2; 2003 *	−0.3090	0.3568	−1.0082	0.3902
P0615	Selectivity increment	Fishery; Bin3; 2003 *	0.6699	0.2727	0.1354	1.2045
P0616	Selectivity increment	Fishery; Bin4; 2003 *	−0.0671	0.2628	−0.5821	0.4478
P0617	Selectivity increment	Fishery; Bin5; 2003 *	0.0369	0.2581	−0.4690	0.5428
P0618	Selectivity increment	Fishery; Bin6; 2003 *	−0.1845	0.2516	−0.6775	0.3085
P0619	Selectivity increment	Fishery; Bin7; 2003 *	0.0299	0.2463	−0.4528	0.5126
P0620	Selectivity increment	Fishery; Bin8; 2003 *	0.0377	0.2530	−0.4581	0.5336
P0621	Selectivity increment	Fishery; Bin9; 2003 *	−0.0010	0.2590	−0.5085	0.5066
P0622	Selectivity increment	Fishery; Bin10; 2003 *	−0.0130	0.2594	−0.5213	0.4954
P0623	Selectivity increment	Fishery; Bin11; 2003 *	−0.0731	0.2643	−0.5912	0.4449
P0624	Selectivity increment	Fishery; Bin12; 2003 *	−0.0559	0.2639	−0.5732	0.4614
P0625	Selectivity increment	Fishery; Bin1; 2004 *	−0.0277	0.4190	−0.8489	0.7936
P0626	Selectivity increment	Fishery; Bin2; 2004 *	−0.1140	0.3769	−0.8527	0.6247
P0627	Selectivity increment	Fishery; Bin3; 2004 *	−0.6762	0.2837	−1.2323	−0.1201
P0628	Selectivity increment	Fishery; Bin4; 2004 *	0.7169	0.2537	0.2196	1.2142
P0629	Selectivity increment	Fishery; Bin5; 2004 *	0.0216	0.2568	−0.4817	0.5249
P0630	Selectivity increment	Fishery; Bin6; 2004 *	−0.1190	0.2671	−0.6425	0.4044
P0631	Selectivity increment	Fishery; Bin7; 2004 *	−0.0233	0.2567	−0.5265	0.4799
P0632	Selectivity increment	Fishery; Bin8; 2004 *	0.0195	0.2529	−0.4762	0.5151
P0633	Selectivity increment	Fishery; Bin9; 2004 *	0.0478	0.2620	−0.4656	0.5613
P0634	Selectivity increment	Fishery; Bin10; 2004 *	0.0528	0.2673	−0.4710	0.5766
P0635	Selectivity increment	Fishery; Bin11; 2004 *	0.0774	0.2682	−0.4484	0.6031
P0636	Selectivity increment	Fishery; Bin12; 2004 *	0.0243	0.2639	−0.4929	0.5415
P0637	Selectivity increment	Fishery; Bin1; 2005 *	−0.0551	0.4191	−0.8765	0.7662
P0638	Selectivity increment	Fishery; Bin2; 2005 *	0.1003	0.3875	−0.6593	0.8598
P0639	Selectivity increment	Fishery; Bin3; 2005 *	−0.0105	0.3079	−0.6139	0.5929
P0640	Selectivity increment	Fishery; Bin4; 2005 *	−0.3443	0.2523	−0.8388	0.1502
P0641	Selectivity increment	Fishery; Bin5; 2005 *	0.4211	0.2497	−0.0682	0.9104
P0642	Selectivity increment	Fishery; Bin6; 2005 *	0.2411	0.2598	−0.2682	0.7503
P0643	Selectivity increment	Fishery; Bin7; 2005 *	−0.1040	0.2576	−0.6089	0.4009
P0644	Selectivity increment	Fishery; Bin8; 2005 *	−0.1580	0.2557	−0.6591	0.3431
P0645	Selectivity increment	Fishery; Bin9; 2005 *	−0.0584	0.2607	−0.5694	0.4527

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0646	Selectivity increment	Fishery; Bin10; 2005 *	0.0069	0.2686	−0.5195	0.5333
P0647	Selectivity increment	Fishery; Bin11; 2005 *	0.0034	0.2723	−0.5304	0.5371
P0648	Selectivity increment	Fishery; Bin12; 2005 *	−0.0424	0.2634	−0.5586	0.4738
P0649	Selectivity increment	Fishery; Bin1; 2006 *	−0.0831	0.4190	−0.9043	0.7381
P0650	Selectivity increment	Fishery; Bin2; 2006 *	0.0868	0.3896	−0.6768	0.8504
P0651	Selectivity increment	Fishery; Bin3; 2006 *	0.5507	0.3137	−0.0642	1.1655
P0652	Selectivity increment	Fishery; Bin4; 2006 *	−0.0815	0.2585	−0.5881	0.4251
P0653	Selectivity increment	Fishery; Bin5; 2006 *	−0.3062	0.2441	−0.7846	0.1723
P0654	Selectivity increment	Fishery; Bin6; 2006 *	−0.0503	0.2432	−0.5269	0.4264
P0655	Selectivity increment	Fishery; Bin7; 2006 *	−0.0551	0.2479	−0.5410	0.4308
P0656	Selectivity increment	Fishery; Bin8; 2006 *	−0.0252	0.2559	−0.5268	0.4763
P0657	Selectivity increment	Fishery; Bin9; 2006 *	−0.0731	0.2605	−0.5836	0.4375
P0658	Selectivity increment	Fishery; Bin10; 2006 *	−0.0319	0.2663	−0.5540	0.4901
P0659	Selectivity increment	Fishery; Bin11; 2006 *	0.0189	0.2723	−0.5148	0.5526
P0660	Selectivity increment	Fishery; Bin12; 2006 *	0.0499	0.2638	−0.4671	0.5670
P0661	Selectivity increment	Fishery; Bin1; 2007 *	−0.0442	0.4186	−0.8647	0.7763
P0662	Selectivity increment	Fishery; Bin2; 2007 *	−0.0142	0.3863	−0.7714	0.7429
P0663	Selectivity increment	Fishery; Bin3; 2007 *	−0.0473	0.3156	−0.6659	0.5713
P0664	Selectivity increment	Fishery; Bin4; 2007 *	0.0167	0.2701	−0.5127	0.5461
P0665	Selectivity increment	Fishery; Bin5; 2007 *	−0.0412	0.2468	−0.5249	0.4425
P0666	Selectivity increment	Fishery; Bin6; 2007 *	0.0189	0.2405	−0.4525	0.4904
P0667	Selectivity increment	Fishery; Bin7; 2007 *	0.0371	0.2432	−0.4395	0.5138
P0668	Selectivity increment	Fishery; Bin8; 2007 *	−0.0466	0.2513	−0.5391	0.4459
P0669	Selectivity increment	Fishery; Bin9; 2007 *	−0.0207	0.2599	−0.5301	0.4886
P0670	Selectivity increment	Fishery; Bin10; 2007 *	0.0048	0.2650	−0.5145	0.5241
P0671	Selectivity increment	Fishery; Bin11; 2007 *	0.0473	0.2699	−0.4817	0.5762
P0672	Selectivity increment	Fishery; Bin12; 2007 *	0.0894	0.2630	−0.4261	0.6048
P0673	Selectivity increment	Fishery; Bin1; 2008 *	−0.0079	0.4177	−0.8266	0.8108
P0674	Selectivity increment	Fishery; Bin2; 2008 *	−0.1140	0.3624	−0.8243	0.5962
P0675	Selectivity increment	Fishery; Bin3; 2008 *	−0.5247	0.3047	−1.1219	0.0724
P0676	Selectivity increment	Fishery; Bin4; 2008 *	−0.1226	0.2806	−0.6725	0.4274
P0677	Selectivity increment	Fishery; Bin5; 2008 *	−0.0705	0.2571	−0.5745	0.4335
P0678	Selectivity increment	Fishery; Bin6; 2008 *	0.0489	0.2404	−0.4224	0.5201
P0679	Selectivity increment	Fishery; Bin7; 2008 *	0.0738	0.2405	−0.3975	0.5452

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0680	Selectivity increment	Fishery; Bin8; 2008 *	0.1378	0.2459	−0.3442	0.6198
P0681	Selectivity increment	Fishery; Bin9; 2008 *	0.1536	0.2555	−0.3472	0.6544
P0682	Selectivity increment	Fishery; Bin10; 2008 *	0.0828	0.2751	−0.4564	0.6220
P0683	Selectivity increment	Fishery; Bin11; 2008 *	0.1590	0.2830	−0.3956	0.7137
P0684	Selectivity increment	Fishery; Bin12; 2008 *	0.1838	0.2618	−0.3292	0.6969
P0685	Selectivity increment	Fishery; Bin1; 2009 *	−0.0405	0.4164	−0.8567	0.7757
P0686	Selectivity increment	Fishery; Bin2; 2009 *	−0.3417	0.3505	−1.0287	0.3453
P0687	Selectivity increment	Fishery; Bin3; 2009 *	0.1995	0.2750	−0.3395	0.7385
P0688	Selectivity increment	Fishery; Bin4; 2009 *	0.0605	0.2709	−0.4704	0.5914
P0689	Selectivity increment	Fishery; Bin5; 2009 *	−0.1362	0.2743	−0.6738	0.4015
P0690	Selectivity increment	Fishery; Bin6; 2009 *	−0.0916	0.2610	−0.6032	0.4199
P0691	Selectivity increment	Fishery; Bin7; 2009 *	−0.0530	0.2484	−0.5399	0.4339
P0692	Selectivity increment	Fishery; Bin8; 2009 *	−0.0082	0.2472	−0.4927	0.4763
P0693	Selectivity increment	Fishery; Bin9; 2009 *	0.0108	0.2611	−0.5009	0.5224
P0694	Selectivity increment	Fishery; Bin10; 2009 *	0.1422	0.2987	−0.4433	0.7277
P0695	Selectivity increment	Fishery; Bin11; 2009 *	0.1431	0.2927	−0.4305	0.7167
P0696	Selectivity increment	Fishery; Bin12; 2009 *	0.1152	0.2631	−0.4006	0.6309
P0697	Selectivity increment	Fishery; Bin1; 2010 *	0.0380	0.4155	−0.7764	0.8524
P0698	Selectivity increment	Fishery; Bin2; 2010 *	0.0713	0.3478	−0.6103	0.7529
P0699	Selectivity increment	Fishery; Bin3; 2010 *	−0.6053	0.2855	−1.1648	−0.0458
P0700	Selectivity increment	Fishery; Bin4; 2010 *	0.4401	0.2487	−0.0475	0.9276
P0701	Selectivity increment	Fishery; Bin5; 2010 *	0.1763	0.2702	−0.3533	0.7059
P0702	Selectivity increment	Fishery; Bin6; 2010 *	−0.2436	0.2851	−0.8024	0.3152
P0703	Selectivity increment	Fishery; Bin7; 2010 *	−0.0793	0.2864	−0.6407	0.4821
P0704	Selectivity increment	Fishery; Bin8; 2010 *	0.0697	0.2604	−0.4407	0.5800
P0705	Selectivity increment	Fishery; Bin9; 2010 *	0.0800	0.2622	−0.4339	0.5940
P0706	Selectivity increment	Fishery; Bin10; 2010 *	0.0262	0.2937	−0.5494	0.6018
P0707	Selectivity increment	Fishery; Bin11; 2010 *	−0.0324	0.2989	−0.6181	0.5533
P0708	Selectivity increment	Fishery; Bin12; 2010 *	0.0589	0.2752	−0.4806	0.5984
P0709	Selectivity increment	Fishery; Bin1; 2011 *	0.0295	0.4160	−0.7858	0.8448
P0710	Selectivity increment	Fishery; Bin2; 2011 *	−0.0180	0.3414	−0.6872	0.6511
P0711	Selectivity increment	Fishery; Bin3; 2011 *	0.0445	0.2837	−0.5116	0.6005
P0712	Selectivity increment	Fishery; Bin4; 2011 *	−1.0769	0.2551	−1.5769	−0.5770
P0713	Selectivity increment	Fishery; Bin5; 2011 *	0.4190	0.2469	−0.0649	0.9030

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0714	Selectivity increment	Fishery; Bin6; 2011 *	0.3585	0.2725	−0.1757	0.8927
P0715	Selectivity increment	Fishery; Bin7; 2011 *	0.1392	0.2841	−0.4176	0.6961
P0716	Selectivity increment	Fishery; Bin8; 2011 *	−0.0029	0.2677	−0.5275	0.5217
P0717	Selectivity increment	Fishery; Bin9; 2011 *	0.0017	0.2713	−0.5299	0.5334
P0718	Selectivity increment	Fishery; Bin10; 2011 *	0.0820	0.2732	−0.4534	0.6174
P0719	Selectivity increment	Fishery; Bin11; 2011 *	0.0533	0.2812	−0.4978	0.6044
P0720	Selectivity increment	Fishery; Bin12; 2011 *	−0.0299	0.2765	−0.5719	0.5121
P0721	Selectivity increment	Fishery; Bin1; 2012 *	−0.0139	0.4163	−0.8298	0.8021
P0722	Selectivity increment	Fishery; Bin2; 2012 *	0.2336	0.3613	−0.4745	0.9417
P0723	Selectivity increment	Fishery; Bin3; 2012 *	0.1890	0.2724	−0.3450	0.7229
P0724	Selectivity increment	Fishery; Bin4; 2012 *	0.6247	0.2561	0.1228	1.1266
P0725	Selectivity increment	Fishery; Bin5; 2012 *	−1.0798	0.2559	−1.5813	−0.5782
P0726	Selectivity increment	Fishery; Bin6; 2012 *	−0.0236	0.2490	−0.5115	0.4644
P0727	Selectivity increment	Fishery; Bin7; 2012 *	0.0121	0.2637	−0.5047	0.5289
P0728	Selectivity increment	Fishery; Bin8; 2012 *	0.0237	0.2771	−0.5194	0.5668
P0729	Selectivity increment	Fishery; Bin9; 2012 *	0.0051	0.2797	−0.5432	0.5533
P0730	Selectivity increment	Fishery; Bin10; 2012 *	−0.0627	0.2976	−0.6459	0.5205
P0731	Selectivity increment	Fishery; Bin11; 2012 *	0.0107	0.3307	−0.6375	0.6590
P0732	Selectivity increment	Fishery; Bin12; 2012 *	0.0811	0.2825	−0.4726	0.6347
P0733	Selectivity increment	Fishery; Bin1; 2013 *	0.0205	0.4165	−0.7958	0.8368
P0734	Selectivity increment	Fishery; Bin2; 2013 *	−0.1915	0.3668	−0.9104	0.5275
P0735	Selectivity increment	Fishery; Bin3; 2013 *	0.0270	0.2910	−0.5434	0.5974
P0736	Selectivity increment	Fishery; Bin4; 2013 *	−0.1384	0.2482	−0.6249	0.3481
P0737	Selectivity increment	Fishery; Bin5; 2013 *	0.5116	0.2562	0.0094	1.0138
P0738	Selectivity increment	Fishery; Bin6; 2013 *	−0.6014	0.2588	−1.1087	−0.0941
P0739	Selectivity increment	Fishery; Bin7; 2013 *	−0.1571	0.2700	−0.6863	0.3720
P0740	Selectivity increment	Fishery; Bin8; 2013 *	0.0768	0.3021	−0.5153	0.6688
P0741	Selectivity increment	Fishery; Bin9; 2013 *	0.1177	0.2916	−0.4538	0.6892
P0742	Selectivity increment	Fishery; Bin10; 2013 *	0.1362	0.2948	−0.4415	0.7139
P0743	Selectivity increment	Fishery; Bin11; 2013 *	0.1077	0.3419	−0.5625	0.7778
P0744	Selectivity increment	Fishery; Bin12; 2013 *	0.0911	0.2877	−0.4728	0.6550
P0745	Selectivity increment	Fishery; Bin1; 2014 *	−0.0208	0.4159	−0.8359	0.7943
P0746	Selectivity increment	Fishery; Bin2; 2014 *	0.1368	0.3573	−0.5635	0.8372
P0747	Selectivity increment	Fishery; Bin3; 2014 *	−0.0265	0.3022	−0.6189	0.5659

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0748	Selectivity increment	Fishery; Bin4; 2014 *	−0.1390	0.2622	−0.6529	0.3749
P0749	Selectivity increment	Fishery; Bin5; 2014 *	−0.0989	0.2478	−0.5845	0.3868
P0750	Selectivity increment	Fishery; Bin6; 2014 *	0.3036	0.2553	−0.1967	0.8038
P0751	Selectivity increment	Fishery; Bin7; 2014 *	0.1480	0.2667	−0.3747	0.6708
P0752	Selectivity increment	Fishery; Bin8; 2014 *	−0.1281	0.2924	−0.7011	0.4449
P0753	Selectivity increment	Fishery; Bin9; 2014 *	−0.1334	0.3287	−0.7777	0.5110
P0754	Selectivity increment	Fishery; Bin10; 2014 *	−0.0146	0.3099	−0.6219	0.5927
P0755	Selectivity increment	Fishery; Bin11; 2014 *	−0.0181	0.3208	−0.6467	0.6106
P0756	Selectivity increment	Fishery; Bin12; 2014 *	−0.0092	0.2918	−0.5810	0.5627
P0757	Selectivity increment	Fishery; Bin1; 2015 *	−0.0049	0.4173	−0.8228	0.8129
P0758	Selectivity increment	Fishery; Bin2; 2015 *	−0.4664	0.3403	−1.1335	0.2006
P0759	Selectivity increment	Fishery; Bin3; 2015 *	1.0864	0.2825	0.5327	1.6401
P0760	Selectivity increment	Fishery; Bin4; 2015 *	0.1881	0.2654	−0.3321	0.7083
P0761	Selectivity increment	Fishery; Bin5; 2015 *	−0.1869	0.2562	−0.6890	0.3151
P0762	Selectivity increment	Fishery; Bin6; 2015 *	−0.1074	0.2478	−0.5930	0.3783
P0763	Selectivity increment	Fishery; Bin7; 2015 *	−0.1479	0.2510	−0.6399	0.3441
P0764	Selectivity increment	Fishery; Bin8; 2015 *	−0.1207	0.2715	−0.6527	0.4113
P0765	Selectivity increment	Fishery; Bin9; 2015 *	−0.1278	0.3310	−0.7765	0.5209
P0766	Selectivity increment	Fishery; Bin10; 2015 *	−0.0646	0.3500	−0.7505	0.6213
P0767	Selectivity increment	Fishery; Bin11; 2015 *	−0.0234	0.3684	−0.7454	0.6986
P0768	Selectivity increment	Fishery; Bin12; 2015 *	−0.0245	0.2928	−0.5983	0.5493
P0769	Selectivity increment	Fishery; Bin1; 2016 *	0.0709	0.4184	−0.7491	0.8909
P0770	Selectivity increment	Fishery; Bin2; 2016 *	0.0205	0.3679	−0.7006	0.7417
P0771	Selectivity increment	Fishery; Bin3; 2016 *	−1.4595	0.2654	−1.9797	−0.9393
P0772	Selectivity increment	Fishery; Bin4; 2016 *	1.0623	0.2498	0.5726	1.5519
P0773	Selectivity increment	Fishery; Bin5; 2016 *	0.2840	0.2593	−0.2243	0.7922
P0774	Selectivity increment	Fishery; Bin6; 2016 *	−0.0027	0.2623	−0.5167	0.5114
P0775	Selectivity increment	Fishery; Bin7; 2016 *	−0.0982	0.2597	−0.6071	0.4107
P0776	Selectivity increment	Fishery; Bin8; 2016 *	0.2058	0.2624	−0.3085	0.7201
P0777	Selectivity increment	Fishery; Bin9; 2016 *	0.1100	0.3019	−0.4817	0.7018
P0778	Selectivity increment	Fishery; Bin10; 2016 *	−0.0858	0.3388	−0.7498	0.5782
P0779	Selectivity increment	Fishery; Bin11; 2016 *	−0.0413	0.3728	−0.7720	0.6894
P0780	Selectivity increment	Fishery; Bin12; 2016 *	−0.0660	0.2958	−0.6458	0.5138
P0781	Selectivity increment	Fishery; Bin1; 2017 *	0.0527	0.4179	−0.7664	0.8719

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0782	Selectivity increment	Fishery; Bin2; 2017 *	0.2806	0.3829	−0.4698	1.0311
P0783	Selectivity increment	Fishery; Bin3; 2017 *	−0.2525	0.3120	−0.8641	0.3590
P0784	Selectivity increment	Fishery; Bin4; 2017 *	−1.1177	0.2428	−1.5936	−0.6418
P0785	Selectivity increment	Fishery; Bin5; 2017 *	0.5963	0.2497	0.1068	1.0857
P0786	Selectivity increment	Fishery; Bin6; 2017 *	0.3303	0.2619	−0.1831	0.8436
P0787	Selectivity increment	Fishery; Bin7; 2017 *	0.1867	0.2691	−0.3407	0.7141
P0788	Selectivity increment	Fishery; Bin8; 2017 *	−0.0775	0.2733	−0.6132	0.4583
P0789	Selectivity increment	Fishery; Bin9; 2017 *	0.0698	0.2694	−0.4582	0.5978
P0790	Selectivity increment	Fishery; Bin10; 2017 *	0.0476	0.2886	−0.5180	0.6133
P0791	Selectivity increment	Fishery; Bin11; 2017 *	−0.0586	0.3038	−0.6541	0.5369
P0792	Selectivity increment	Fishery; Bin12; 2017 *	−0.0577	0.2800	−0.6065	0.4910
P0793	Selectivity increment	Fishery; Bin1; 2018 *	0.0269	0.4176	−0.7916	0.8454
P0794	Selectivity increment	Fishery; Bin2; 2018 *	0.3297	0.3844	−0.4237	1.0831
P0795	Selectivity increment	Fishery; Bin3; 2018 *	0.1383	0.3454	−0.5386	0.8153
P0796	Selectivity increment	Fishery; Bin4; 2018 *	−0.5920	0.2685	−1.1182	−0.0658
P0797	Selectivity increment	Fishery; Bin5; 2018 *	0.0098	0.2416	−0.4637	0.4833
P0798	Selectivity increment	Fishery; Bin6; 2018 *	0.1787	0.2486	−0.3085	0.6660
P0799	Selectivity increment	Fishery; Bin7; 2018 *	0.1098	0.2599	−0.3996	0.6191
P0800	Selectivity increment	Fishery; Bin8; 2018 *	0.0802	0.2732	−0.4553	0.6156
P0801	Selectivity increment	Fishery; Bin9; 2018 *	−0.0363	0.2658	−0.5572	0.4847
P0802	Selectivity increment	Fishery; Bin10; 2018 *	−0.0785	0.2671	−0.6021	0.4451
P0803	Selectivity increment	Fishery; Bin11; 2018 *	−0.0815	0.2758	−0.6221	0.4591
P0804	Selectivity increment	Fishery; Bin12; 2018 *	−0.0851	0.2740	−0.6221	0.4518
P0805	Selectivity increment	Fishery; Bin1; 2019 *	0.0466	0.4171	−0.7710	0.8641
P0806	Selectivity increment	Fishery; Bin2; 2019 *	0.4753	0.3768	−0.2631	1.2138
P0807	Selectivity increment	Fishery; Bin3; 2019 *	0.2133	0.3498	−0.4724	0.8989
P0808	Selectivity increment	Fishery; Bin4; 2019 *	−0.3828	0.3044	−0.9794	0.2138
P0809	Selectivity increment	Fishery; Bin5; 2019 *	−1.2476	0.2535	−1.7444	−0.7507
P0810	Selectivity increment	Fishery; Bin6; 2019 *	0.0949	0.2440	−0.3833	0.5732
P0811	Selectivity increment	Fishery; Bin7; 2019 *	0.4408	0.2500	−0.0491	0.9307
P0812	Selectivity increment	Fishery; Bin8; 2019 *	0.2212	0.2554	−0.2793	0.7217
P0813	Selectivity increment	Fishery; Bin9; 2019 *	0.1280	0.2577	−0.3771	0.6332
P0814	Selectivity increment	Fishery; Bin10; 2019 *	0.1319	0.2624	−0.3825	0.6463
P0815	Selectivity increment	Fishery; Bin11; 2019 *	−0.0475	0.2710	−0.5786	0.4836

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0816	Selectivity increment	Fishery; Bin12; 2019 *	−0.0742	0.2705	−0.6044	0.4561
P0817	Selectivity increment	Fishery; Bin1; 2020 *	−0.0191	0.4206	−0.8434	0.8052
P0818	Selectivity increment	Fishery; Bin2; 2020 *	0.6177	0.3313	−0.0316	1.2671
P0819	Selectivity increment	Fishery; Bin3; 2020 *	0.4207	0.3273	−0.2208	1.0622
P0820	Selectivity increment	Fishery; Bin4; 2020 *	0.5413	0.3105	−0.0673	1.1498
P0821	Selectivity increment	Fishery; Bin5; 2020 *	0.1307	0.2757	−0.4096	0.6709
P0822	Selectivity increment	Fishery; Bin6; 2020 *	−0.4191	0.2437	−0.8968	0.0586
P0823	Selectivity increment	Fishery; Bin7; 2020 *	−0.5005	0.2439	−0.9786	−0.0224
P0824	Selectivity increment	Fishery; Bin8; 2020 *	−0.4159	0.2603	−0.9260	0.0942
P0825	Selectivity increment	Fishery; Bin9; 2020 *	−0.1799	0.2700	−0.7091	0.3493
P0826	Selectivity increment	Fishery; Bin10; 2020 *	−0.1332	0.2641	−0.6509	0.3844
P0827	Selectivity increment	Fishery; Bin11; 2020 *	0.0035	0.2686	−0.5231	0.5300
P0828	Selectivity increment	Fishery; Bin12; 2020 *	−0.0460	0.2683	−0.5718	0.4798
P0829	Selectivity increment	Fishery; Bin1; 2021 *	−0.0184	0.4212	−0.8439	0.8071
P0830	Selectivity increment	Fishery; Bin2; 2021 *	0.3251	0.2749	−0.2138	0.8639
P0831	Selectivity increment	Fishery; Bin3; 2021 *	0.9147	0.2699	0.3857	1.4437
P0832	Selectivity increment	Fishery; Bin4; 2021 *	−0.1532	0.2838	−0.7093	0.4030
P0833	Selectivity increment	Fishery; Bin5; 2021 *	−0.0152	0.2871	−0.5779	0.5474
P0834	Selectivity increment	Fishery; Bin6; 2021 *	−0.6384	0.2690	−1.1655	−0.1113
P0835	Selectivity increment	Fishery; Bin7; 2021 *	−0.1600	0.2500	−0.6500	0.3299
P0836	Selectivity increment	Fishery; Bin8; 2021 *	0.0909	0.2552	−0.4093	0.5910
P0837	Selectivity increment	Fishery; Bin9; 2021 *	−0.1014	0.2714	−0.6334	0.4305
P0838	Selectivity increment	Fishery; Bin10; 2021 *	−0.1057	0.2667	−0.6285	0.4170
P0839	Selectivity increment	Fishery; Bin11; 2021 *	−0.0815	0.2704	−0.6114	0.4484
P0840	Selectivity increment	Fishery; Bin12; 2021 *	−0.0567	0.2670	−0.5800	0.4666
P0841	Selectivity increment	Fishery; Bin1; 2022 *	0.0125	0.4227	−0.8160	0.8411
P0842	Selectivity increment	Fishery; Bin2; 2022 *	0.2365	0.3079	−0.3670	0.8399
P0843	Selectivity increment	Fishery; Bin3; 2022 *	−0.4506	0.2537	−0.9478	0.0466
P0844	Selectivity increment	Fishery; Bin4; 2022 *	0.5723	0.2483	0.0856	1.0591
P0845	Selectivity increment	Fishery; Bin5; 2022 *	0.0776	0.2801	−0.4714	0.6266
P0846	Selectivity increment	Fishery; Bin6; 2022 *	0.0682	0.2955	−0.5109	0.6473
P0847	Selectivity increment	Fishery; Bin7; 2022 *	−0.2542	0.2823	−0.8076	0.2991
P0848	Selectivity increment	Fishery; Bin8; 2022 *	−0.0369	0.2568	−0.5401	0.4664
P0849	Selectivity increment	Fishery; Bin9; 2022 *	−0.0857	0.2572	−0.5898	0.4184

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0850	Selectivity increment	Fishery; Bin10; 2022 *	−0.1450	0.2819	−0.6976	0.4076
P0851	Selectivity increment	Fishery; Bin11; 2022 *	−0.0111	0.2962	−0.5915	0.5694
P0852	Selectivity increment	Fishery; Bin12; 2022 *	0.0163	0.2718	−0.5165	0.5491
P0853	Selectivity increment	Fishery; Bin1; 2023 *	0.0335	0.4240	−0.7975	0.8646
P0854	Selectivity increment	Fishery; Bin2; 2023 *	−0.5536	0.3485	−1.2367	0.1295
P0855	Selectivity increment	Fishery; Bin3; 2023 *	0.1024	0.3004	−0.4864	0.6911
P0856	Selectivity increment	Fishery; Bin4; 2023 *	−0.1481	0.2631	−0.6638	0.3677
P0857	Selectivity increment	Fishery; Bin5; 2023 *	0.5445	0.2616	0.0319	1.0572
P0858	Selectivity increment	Fishery; Bin6; 2023 *	0.3275	0.2881	−0.2372	0.8923
P0859	Selectivity increment	Fishery; Bin7; 2023 *	−0.1214	0.2972	−0.7039	0.4612
P0860	Selectivity increment	Fishery; Bin8; 2023 *	−0.2840	0.3093	−0.8901	0.3222
P0861	Selectivity increment	Fishery; Bin9; 2023 *	0.0082	0.2801	−0.5408	0.5572
P0862	Selectivity increment	Fishery; Bin10; 2023 *	0.0548	0.2938	−0.5211	0.6307
P0863	Selectivity increment	Fishery; Bin11; 2023 *	0.0069	0.3066	−0.5941	0.6079
P0864	Selectivity increment	Fishery; Bin12; 2023 *	0.0292	0.2763	−0.5124	0.5707
P0865	Selectivity increment	Fishery; Bin1; 2024 *	0.0203	0.4247	−0.8122	0.8528
P0866	Selectivity increment	Fishery; Bin2; 2024 *	−0.5495	0.3664	−1.2676	0.1685
P0867	Selectivity increment	Fishery; Bin3; 2024 *	0.1606	0.3170	−0.4607	0.7819
P0868	Selectivity increment	Fishery; Bin4; 2024 *	0.1045	0.2938	−0.4713	0.6804
P0869	Selectivity increment	Fishery; Bin5; 2024 *	−0.2087	0.2602	−0.7186	0.3012
P0870	Selectivity increment	Fishery; Bin6; 2024 *	0.3659	0.2595	−0.1427	0.8745
P0871	Selectivity increment	Fishery; Bin7; 2024 *	0.2160	0.2755	−0.3239	0.7559
P0872	Selectivity increment	Fishery; Bin8; 2024 *	−0.0331	0.3171	−0.6545	0.5884
P0873	Selectivity increment	Fishery; Bin9; 2024 *	−0.0209	0.2958	−0.6007	0.5589
P0874	Selectivity increment	Fishery; Bin10; 2024 *	−0.0052	0.2983	−0.5898	0.5795
P0875	Selectivity increment	Fishery; Bin11; 2024 *	−0.0403	0.3419	−0.7104	0.6298
P0876	Selectivity increment	Fishery; Bin12; 2024 *	−0.0097	0.2891	−0.5763	0.5569
P0877	Selectivity increment	Fishery; Bin1; 2025 *	0.0465	0.4277	−0.7917	0.8847
P0878	Selectivity increment	Fishery; Bin2; 2025 *	−0.1805	0.3689	−0.9034	0.5425
P0879	Selectivity increment	Fishery; Bin3; 2025 *	−0.3735	0.3337	−1.0275	0.2805
P0880	Selectivity increment	Fishery; Bin4; 2025 *	−0.1775	0.3158	−0.7964	0.4414
P0881	Selectivity increment	Fishery; Bin5; 2025 *	−0.2383	0.2955	−0.8174	0.3409
P0882	Selectivity increment	Fishery; Bin6; 2025 *	0.1600	0.2522	−0.3343	0.6542
P0883	Selectivity increment	Fishery; Bin7; 2025 *	0.3901	0.2527	−0.1051	0.8854

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0884	Selectivity increment	Fishery; Bin8; 2025 *	0.2710	0.2754	−0.2687	0.8107
P0885	Selectivity increment	Fishery; Bin9; 2025 *	0.0724	0.2890	−0.4940	0.6388
P0886	Selectivity increment	Fishery; Bin10; 2025 *	0.0628	0.2921	−0.5098	0.6354
P0887	Selectivity increment	Fishery; Bin11; 2025 *	−0.0233	0.3339	−0.6777	0.6312
P0888	Selectivity increment	Fishery; Bin12; 2025 *	−0.0097	0.2915	−0.5811	0.5617
P0889	Selectivity increment	Fishery; Bin1; 2026 *	0.0057	0.4326	−0.8422	0.8537
P0890	Selectivity increment	Fishery; Bin2; 2026 *	−0.0131	0.4291	−0.8541	0.8278
P0891	Selectivity increment	Fishery; Bin3; 2026 *	0.0024	0.4306	−0.8416	0.8464
P0892	Selectivity increment	Fishery; Bin4; 2026 *	0.0136	0.4330	−0.8351	0.8622
P0893	Selectivity increment	Fishery; Bin5; 2026 *	0.0458	0.4381	−0.8128	0.9045
P0894	Selectivity increment	Fishery; Bin6; 2026 *	−0.1266	0.3089	−0.7320	0.4787
P0895	Selectivity increment	Fishery; Bin7; 2026 *	−0.1276	0.2822	−0.6807	0.4256
P0896	Selectivity increment	Fishery; Bin8; 2026 *	0.0580	0.2800	−0.4908	0.6068
P0897	Selectivity increment	Fishery; Bin9; 2026 *	0.0408	0.2804	−0.5087	0.5903
P0898	Selectivity increment	Fishery; Bin10; 2026 *	0.0264	0.2850	−0.5322	0.5850
P0899	Selectivity increment	Fishery; Bin11; 2026 *	0.0629	0.3040	−0.5330	0.6587
P0900	Selectivity increment	Fishery; Bin12; 2026 *	0.0118	0.2926	−0.5617	0.5853
P0901	Selectivity increment	AVO; Bin2; 1994 *	0.4268	0.3221	−0.2045	1.0581
P0902	Selectivity increment	AVO; Bin3; 1994 *	0.2014	0.3578	−0.4998	0.9027
P0903	Selectivity increment	AVO; Bin4; 1994 *	0.0307	0.3479	−0.6511	0.7125
P0904	Selectivity increment	AVO; Bin5; 1994 *	0.1390	0.3399	−0.5273	0.8052
P0905	Selectivity increment	AVO; Bin6; 1994 *	0.1843	0.3570	−0.5154	0.8839
P0906	Selectivity increment	AVO; Bin7; 1994 *	−0.2044	0.3577	−0.9055	0.4967
P0907	Selectivity increment	AVO; Bin8; 1994 *	0.0739	0.3182	−0.5498	0.6976
P0908	Selectivity increment	AVO; Bin2; 1995 *	0.2193	0.2842	−0.3376	0.7763
P0909	Selectivity increment	AVO; Bin3; 1995 *	0.2252	0.2938	−0.3507	0.8010
P0910	Selectivity increment	AVO; Bin4; 1995 *	0.1966	0.2925	−0.3766	0.7699
P0911	Selectivity increment	AVO; Bin5; 1995 *	0.1891	0.2912	−0.3817	0.7599
P0912	Selectivity increment	AVO; Bin6; 1995 *	0.2052	0.2922	−0.3676	0.7779
P0913	Selectivity increment	AVO; Bin7; 1995 *	0.2084	0.2921	−0.3640	0.7808
P0914	Selectivity increment	AVO; Bin8; 1995 *	0.2116	0.2641	−0.3061	0.7293
P0915	Selectivity increment	AVO; Bin2; 1996 *	0.2181	0.2742	−0.3193	0.7556
P0916	Selectivity increment	AVO; Bin3; 1996 *	0.2133	0.2742	−0.3240	0.7507
P0917	Selectivity increment	AVO; Bin4; 1996 *	0.2070	0.2725	−0.3271	0.7412

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0918	Selectivity increment	AVO; Bin5; 1996 *	0.1986	0.2723	−0.3351	0.7323
P0919	Selectivity increment	AVO; Bin6; 1996 *	0.2015	0.2730	−0.3335	0.7364
P0920	Selectivity increment	AVO; Bin7; 1996 *	0.2067	0.2721	−0.3266	0.7400
P0921	Selectivity increment	AVO; Bin8; 1996 *	0.2151	0.2472	−0.2694	0.6997
P0922	Selectivity increment	AVO; Bin2; 1997 *	0.2620	0.2631	−0.2535	0.7776
P0923	Selectivity increment	AVO; Bin3; 1997 *	0.2517	0.2607	−0.2594	0.7627
P0924	Selectivity increment	AVO; Bin4; 1997 *	0.2206	0.2594	−0.2877	0.7289
P0925	Selectivity increment	AVO; Bin5; 1997 *	0.2321	0.2585	−0.2745	0.7387
P0926	Selectivity increment	AVO; Bin6; 1997 *	0.2202	0.2597	−0.2888	0.7292
P0927	Selectivity increment	AVO; Bin7; 1997 *	0.2254	0.2590	−0.2822	0.7331
P0928	Selectivity increment	AVO; Bin8; 1997 *	0.2178	0.2340	−0.2408	0.6764
P0929	Selectivity increment	AVO; Bin2; 1998 *	0.2201	0.2506	−0.2710	0.7112
P0930	Selectivity increment	AVO; Bin3; 1998 *	0.2988	0.2509	−0.1930	0.7905
P0931	Selectivity increment	AVO; Bin4; 1998 *	0.2887	0.2506	−0.2025	0.7799
P0932	Selectivity increment	AVO; Bin5; 1998 *	0.2513	0.2507	−0.2401	0.7427
P0933	Selectivity increment	AVO; Bin6; 1998 *	0.2567	0.2509	−0.2351	0.7485
P0934	Selectivity increment	AVO; Bin7; 1998 *	0.2612	0.2497	−0.2283	0.7507
P0935	Selectivity increment	AVO; Bin8; 1998 *	0.2331	0.2239	−0.2057	0.6719
P0936	Selectivity increment	AVO; Bin2; 1999 *	0.2312	0.2468	−0.2526	0.7149
P0937	Selectivity increment	AVO; Bin3; 1999 *	0.3187	0.2435	−0.1584	0.7959
P0938	Selectivity increment	AVO; Bin4; 1999 *	0.3135	0.2427	−0.1622	0.7891
P0939	Selectivity increment	AVO; Bin5; 1999 *	0.2774	0.2435	−0.1999	0.7547
P0940	Selectivity increment	AVO; Bin6; 1999 *	0.2693	0.2437	−0.2083	0.7470
P0941	Selectivity increment	AVO; Bin7; 1999 *	0.2849	0.2420	−0.1894	0.7593
P0942	Selectivity increment	AVO; Bin8; 1999 *	0.2551	0.2166	−0.1694	0.6796
P0943	Selectivity increment	AVO; Bin2; 2000 *	0.2540	0.2396	−0.2156	0.7236
P0944	Selectivity increment	AVO; Bin3; 2000 *	0.2719	0.2393	−0.1972	0.7410
P0945	Selectivity increment	AVO; Bin4; 2000 *	0.3283	0.2384	−0.1390	0.7955
P0946	Selectivity increment	AVO; Bin5; 2000 *	0.2944	0.2392	−0.1744	0.7632
P0947	Selectivity increment	AVO; Bin6; 2000 *	0.2994	0.2398	−0.1707	0.7694
P0948	Selectivity increment	AVO; Bin7; 2000 *	0.2678	0.2390	−0.2007	0.7363
P0949	Selectivity increment	AVO; Bin8; 2000 *	0.2747	0.2126	−0.1419	0.6913
P0950	Selectivity increment	AVO; Bin2; 2001 *	0.3133	0.2345	−0.1463	0.7729
P0951	Selectivity increment	AVO; Bin3; 2001 *	0.3023	0.2373	−0.1628	0.7673

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0952	Selectivity increment	AVO; Bin4; 2001 *	0.2844	0.2370	−0.1800	0.7489
P0953	Selectivity increment	AVO; Bin5; 2001 *	0.3131	0.2372	−0.1517	0.7780
P0954	Selectivity increment	AVO; Bin6; 2001 *	0.3172	0.2374	−0.1480	0.7824
P0955	Selectivity increment	AVO; Bin7; 2001 *	0.2960	0.2370	−0.1685	0.7605
P0956	Selectivity increment	AVO; Bin8; 2001 *	0.2805	0.2102	−0.1314	0.6925
P0957	Selectivity increment	AVO; Bin2; 2002 *	0.3008	0.2355	−0.1608	0.7625
P0958	Selectivity increment	AVO; Bin3; 2002 *	0.3180	0.2342	−0.1411	0.7771
P0959	Selectivity increment	AVO; Bin4; 2002 *	0.3056	0.2342	−0.1533	0.7645
P0960	Selectivity increment	AVO; Bin5; 2002 *	0.3188	0.2346	−0.1410	0.7786
P0961	Selectivity increment	AVO; Bin6; 2002 *	0.3244	0.2345	−0.1353	0.7840
P0962	Selectivity increment	AVO; Bin7; 2002 *	0.3085	0.2345	−0.1510	0.7680
P0963	Selectivity increment	AVO; Bin8; 2002 *	0.2908	0.2078	−0.1165	0.6980
P0964	Selectivity increment	AVO; Bin2; 2003 *	0.2507	0.2309	−0.2019	0.7033
P0965	Selectivity increment	AVO; Bin3; 2003 *	0.2393	0.2357	−0.2227	0.7013
P0966	Selectivity increment	AVO; Bin4; 2003 *	0.3117	0.2351	−0.1490	0.7725
P0967	Selectivity increment	AVO; Bin5; 2003 *	0.3291	0.2351	−0.1317	0.7898
P0968	Selectivity increment	AVO; Bin6; 2003 *	0.3118	0.2353	−0.1493	0.7729
P0969	Selectivity increment	AVO; Bin7; 2003 *	0.3008	0.2349	−0.1597	0.7612
P0970	Selectivity increment	AVO; Bin8; 2003 *	0.3001	0.2080	−0.1075	0.7077
P0971	Selectivity increment	AVO; Bin2; 2004 *	0.2241	0.2342	−0.2349	0.6830
P0972	Selectivity increment	AVO; Bin3; 2004 *	0.2365	0.2351	−0.2244	0.6973
P0973	Selectivity increment	AVO; Bin4; 2004 *	0.3237	0.2336	−0.1342	0.7816
P0974	Selectivity increment	AVO; Bin5; 2004 *	0.3248	0.2340	−0.1338	0.7835
P0975	Selectivity increment	AVO; Bin6; 2004 *	0.3064	0.2345	−0.1533	0.7661
P0976	Selectivity increment	AVO; Bin7; 2004 *	0.2994	0.2344	−0.1600	0.7588
P0977	Selectivity increment	AVO; Bin8; 2004 *	0.2971	0.2078	−0.1101	0.7043
P0978	Selectivity increment	AVO; Bin2; 2005 *	0.3122	0.2340	−0.1465	0.7708
P0979	Selectivity increment	AVO; Bin3; 2005 *	0.2707	0.2370	−0.1939	0.7353
P0980	Selectivity increment	AVO; Bin4; 2005 *	0.2088	0.2361	−0.2539	0.6714
P0981	Selectivity increment	AVO; Bin5; 2005 *	0.2422	0.2355	−0.2193	0.7037
P0982	Selectivity increment	AVO; Bin6; 2005 *	0.2737	0.2355	−0.1878	0.7352
P0983	Selectivity increment	AVO; Bin7; 2005 *	0.2837	0.2354	−0.1776	0.7450
P0984	Selectivity increment	AVO; Bin8; 2005 *	0.2886	0.2091	−0.1212	0.6985
P0985	Selectivity increment	AVO; Bin2; 2006 *	0.2699	0.2375	−0.1956	0.7355

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P0986	Selectivity increment	AVO; Bin3; 2006 *	0.2465	0.2377	−0.2194	0.7124
P0987	Selectivity increment	AVO; Bin4; 2006 *	0.2044	0.2360	−0.2583	0.6670
P0988	Selectivity increment	AVO; Bin5; 2006 *	0.2238	0.2353	−0.2374	0.6850
P0989	Selectivity increment	AVO; Bin6; 2006 *	0.2520	0.2353	−0.2091	0.7131
P0990	Selectivity increment	AVO; Bin7; 2006 *	0.2653	0.2354	−0.1961	0.7267
P0991	Selectivity increment	AVO; Bin8; 2006 *	0.2690	0.2100	−0.1426	0.6806
P0992	Selectivity increment	AVO; Bin2; 2007 *	0.2692	0.2368	−0.1949	0.7332
P0993	Selectivity increment	AVO; Bin3; 2007 *	0.2569	0.2368	−0.2071	0.7210
P0994	Selectivity increment	AVO; Bin4; 2007 *	0.2281	0.2359	−0.2343	0.6904
P0995	Selectivity increment	AVO; Bin5; 2007 *	0.2419	0.2351	−0.2189	0.7027
P0996	Selectivity increment	AVO; Bin6; 2007 *	0.2516	0.2350	−0.2090	0.7122
P0997	Selectivity increment	AVO; Bin7; 2007 *	0.2401	0.2350	−0.2206	0.7008
P0998	Selectivity increment	AVO; Bin8; 2007 *	0.2437	0.2092	−0.1664	0.6538
P0999	Selectivity increment	AVO; Bin2; 2008 *	0.2695	0.2362	−0.1933	0.7324
P1000	Selectivity increment	AVO; Bin3; 2008 *	0.2401	0.2375	−0.2253	0.7055
P1001	Selectivity increment	AVO; Bin4; 2008 *	0.1914	0.2376	−0.2743	0.6571
P1002	Selectivity increment	AVO; Bin5; 2008 *	0.1803	0.2377	−0.2856	0.6462
P1003	Selectivity increment	AVO; Bin6; 2008 *	0.1896	0.2375	−0.2758	0.6551
P1004	Selectivity increment	AVO; Bin7; 2008 *	0.2055	0.2371	−0.2592	0.6702
P1005	Selectivity increment	AVO; Bin8; 2008 *	0.2248	0.2109	−0.1886	0.6382
P1006	Selectivity increment	AVO; Bin2; 2009 *	0.2256	0.2352	−0.2354	0.6867
P1007	Selectivity increment	AVO; Bin3; 2009 *	0.2164	0.2381	−0.2503	0.6832
P1008	Selectivity increment	AVO; Bin4; 2009 *	0.1855	0.2380	−0.2810	0.6519
P1009	Selectivity increment	AVO; Bin5; 2009 *	0.1817	0.2386	−0.2860	0.6494
P1010	Selectivity increment	AVO; Bin6; 2009 *	0.1921	0.2389	−0.2761	0.6602
P1011	Selectivity increment	AVO; Bin7; 2009 *	0.2021	0.2383	−0.2650	0.6692
P1012	Selectivity increment	AVO; Bin8; 2009 *	0.2012	0.2114	−0.2132	0.6156
P1013	Selectivity increment	AVO; Bin2; 2010 *	0.2367	0.2353	−0.2245	0.6979
P1014	Selectivity increment	AVO; Bin3; 2010 *	0.1889	0.2390	−0.2795	0.6574
P1015	Selectivity increment	AVO; Bin4; 2010 *	0.2100	0.2375	−0.2554	0.6754
P1016	Selectivity increment	AVO; Bin5; 2010 *	0.1807	0.2382	−0.2861	0.6476
P1017	Selectivity increment	AVO; Bin6; 2010 *	0.1855	0.2386	−0.2821	0.6531
P1018	Selectivity increment	AVO; Bin7; 2010 *	0.1926	0.2382	−0.2742	0.6595
P1019	Selectivity increment	AVO; Bin8; 2010 *	0.1823	0.2112	−0.2316	0.5962

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1020	Selectivity increment	AVO; Bin2; 2011 *	0.2198	0.2349	−0.2407	0.6802
P1021	Selectivity increment	AVO; Bin3; 2011 *	0.1828	0.2393	−0.2863	0.6519
P1022	Selectivity increment	AVO; Bin4; 2011 *	0.1560	0.2379	−0.3103	0.6223
P1023	Selectivity increment	AVO; Bin5; 2011 *	0.1683	0.2383	−0.2987	0.6354
P1024	Selectivity increment	AVO; Bin6; 2011 *	0.1801	0.2385	−0.2872	0.6475
P1025	Selectivity increment	AVO; Bin7; 2011 *	0.1860	0.2383	−0.2810	0.6531
P1026	Selectivity increment	AVO; Bin8; 2011 *	0.1717	0.2116	−0.2430	0.5863
P1027	Selectivity increment	AVO; Bin2; 2012 *	0.2201	0.2381	−0.2465	0.6868
P1028	Selectivity increment	AVO; Bin3; 2012 *	0.1874	0.2380	−0.2791	0.6539
P1029	Selectivity increment	AVO; Bin4; 2012 *	0.1557	0.2357	−0.3062	0.6177
P1030	Selectivity increment	AVO; Bin5; 2012 *	0.1715	0.2374	−0.2937	0.6367
P1031	Selectivity increment	AVO; Bin6; 2012 *	0.1797	0.2375	−0.2857	0.6451
P1032	Selectivity increment	AVO; Bin7; 2012 *	0.1825	0.2377	−0.2834	0.6484
P1033	Selectivity increment	AVO; Bin8; 2012 *	0.1626	0.2112	−0.2513	0.5765
P1034	Selectivity increment	AVO; Bin2; 2013 *	0.2090	0.2344	−0.2504	0.6683
P1035	Selectivity increment	AVO; Bin3; 2013 *	0.2204	0.2364	−0.2429	0.6837
P1036	Selectivity increment	AVO; Bin4; 2013 *	0.2464	0.2350	−0.2142	0.7070
P1037	Selectivity increment	AVO; Bin5; 2013 *	0.2349	0.2351	−0.2258	0.6957
P1038	Selectivity increment	AVO; Bin6; 2013 *	0.2267	0.2351	−0.2342	0.6875
P1039	Selectivity increment	AVO; Bin7; 2013 *	0.2029	0.2354	−0.2585	0.6643
P1040	Selectivity increment	AVO; Bin8; 2013 *	0.1606	0.2081	−0.2473	0.5686
P1041	Selectivity increment	AVO; Bin2; 2014 *	0.2387	0.2348	−0.2216	0.6989
P1042	Selectivity increment	AVO; Bin3; 2014 *	0.2282	0.2328	−0.2281	0.6845
P1043	Selectivity increment	AVO; Bin4; 2014 *	0.2623	0.2318	−0.1920	0.7167
P1044	Selectivity increment	AVO; Bin5; 2014 *	0.2438	0.2321	−0.2111	0.6986
P1045	Selectivity increment	AVO; Bin6; 2014 *	0.2492	0.2316	−0.2048	0.7032
P1046	Selectivity increment	AVO; Bin7; 2014 *	0.2315	0.2330	−0.2251	0.6881
P1047	Selectivity increment	AVO; Bin8; 2014 *	0.1816	0.2054	−0.2211	0.5842
P1048	Selectivity increment	AVO; Bin2; 2015 *	0.1440	0.2281	−0.3030	0.5910
P1049	Selectivity increment	AVO; Bin3; 2015 *	0.2648	0.2323	−0.1904	0.7200
P1050	Selectivity increment	AVO; Bin4; 2015 *	0.2820	0.2321	−0.1729	0.7370
P1051	Selectivity increment	AVO; Bin5; 2015 *	0.2515	0.2330	−0.2053	0.7082
P1052	Selectivity increment	AVO; Bin6; 2015 *	0.2218	0.2332	−0.2352	0.6788
P1053	Selectivity increment	AVO; Bin7; 2015 *	0.2269	0.2336	−0.2309	0.6847

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1054	Selectivity increment	AVO; Bin8; 2015 *	0.2145	0.2058	−0.1888	0.6178
P1055	Selectivity increment	AVO; Bin2; 2016 *	0.1171	0.2337	−0.3409	0.5752
P1056	Selectivity increment	AVO; Bin3; 2016 *	0.2099	0.2333	−0.2474	0.6671
P1057	Selectivity increment	AVO; Bin4; 2016 *	0.2846	0.2324	−0.1708	0.7401
P1058	Selectivity increment	AVO; Bin5; 2016 *	0.2301	0.2344	−0.2294	0.6896
P1059	Selectivity increment	AVO; Bin6; 2016 *	0.2050	0.2348	−0.2552	0.6652
P1060	Selectivity increment	AVO; Bin7; 2016 *	0.2006	0.2352	−0.2604	0.6616
P1061	Selectivity increment	AVO; Bin8; 2016 *	0.2355	0.2086	−0.1734	0.6444
P1062	Selectivity increment	AVO; Bin2; 2017 *	0.2189	0.2350	−0.2416	0.6795
P1063	Selectivity increment	AVO; Bin3; 2017 *	0.1560	0.2388	−0.3121	0.6240
P1064	Selectivity increment	AVO; Bin4; 2017 *	0.1020	0.2369	−0.3624	0.5664
P1065	Selectivity increment	AVO; Bin5; 2017 *	0.1924	0.2378	−0.2738	0.6585
P1066	Selectivity increment	AVO; Bin6; 2017 *	0.2094	0.2389	−0.2588	0.6777
P1067	Selectivity increment	AVO; Bin7; 2017 *	0.2076	0.2398	−0.2625	0.6776
P1068	Selectivity increment	AVO; Bin8; 2017 *	0.2362	0.2134	−0.1821	0.6545
P1069	Selectivity increment	AVO; Bin2; 2018 *	0.1941	0.2406	−0.2776	0.6657
P1070	Selectivity increment	AVO; Bin3; 2018 *	0.1519	0.2425	−0.3234	0.6271
P1071	Selectivity increment	AVO; Bin4; 2018 *	0.1254	0.2409	−0.3467	0.5975
P1072	Selectivity increment	AVO; Bin5; 2018 *	0.1977	0.2397	−0.2722	0.6675
P1073	Selectivity increment	AVO; Bin6; 2018 *	0.2034	0.2414	−0.2698	0.6766
P1074	Selectivity increment	AVO; Bin7; 2018 *	0.2057	0.2430	−0.2707	0.6820
P1075	Selectivity increment	AVO; Bin8; 2018 *	0.2230	0.2169	−0.2022	0.6481
P1076	Selectivity increment	AVO; Bin2; 2019 *	0.2102	0.2403	−0.2607	0.6811
P1077	Selectivity increment	AVO; Bin3; 2019 *	0.2093	0.2441	−0.2692	0.6878
P1078	Selectivity increment	AVO; Bin4; 2019 *	0.2316	0.2446	−0.2477	0.7110
P1079	Selectivity increment	AVO; Bin5; 2019 *	0.2064	0.2446	−0.2730	0.6857
P1080	Selectivity increment	AVO; Bin6; 2019 *	0.2099	0.2448	−0.2698	0.6897
P1081	Selectivity increment	AVO; Bin7; 2019 *	0.2061	0.2448	−0.2737	0.6859
P1082	Selectivity increment	AVO; Bin8; 2019 *	0.2140	0.2181	−0.2135	0.6416
P1083	Selectivity increment	AVO; Bin2; 2020 *	0.1981	0.2447	−0.2815	0.6777
P1084	Selectivity increment	AVO; Bin3; 2020 *	0.2065	0.2472	−0.2781	0.6910
P1085	Selectivity increment	AVO; Bin4; 2020 *	0.2306	0.2472	−0.2538	0.7151
P1086	Selectivity increment	AVO; Bin5; 2020 *	0.1997	0.2474	−0.2853	0.6847
P1087	Selectivity increment	AVO; Bin6; 2020 *	0.1935	0.2475	−0.2916	0.6787

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1088	Selectivity increment	AVO; Bin7; 2020 *	0.2048	0.2475	−0.2802	0.6898
P1089	Selectivity increment	AVO; Bin8; 2020 *	0.2187	0.2214	−0.2152	0.6527
P1090	Selectivity increment	AVO; Bin2; 2021 *	0.1941	0.2490	−0.2939	0.6820
P1091	Selectivity increment	AVO; Bin3; 2021 *	0.1939	0.2508	−0.2977	0.6855
P1092	Selectivity increment	AVO; Bin4; 2021 *	0.2316	0.2500	−0.2584	0.7216
P1093	Selectivity increment	AVO; Bin5; 2021 *	0.1960	0.2505	−0.2951	0.6871
P1094	Selectivity increment	AVO; Bin6; 2021 *	0.1941	0.2507	−0.2972	0.6854
P1095	Selectivity increment	AVO; Bin7; 2021 *	0.2074	0.2505	−0.2837	0.6985
P1096	Selectivity increment	AVO; Bin8; 2021 *	0.2175	0.2256	−0.2246	0.6596
P1097	Selectivity increment	AVO; Bin2; 2022 *	0.1766	0.2549	−0.3230	0.6761
P1098	Selectivity increment	AVO; Bin3; 2022 *	0.1694	0.2554	−0.3312	0.6701
P1099	Selectivity increment	AVO; Bin4; 2022 *	0.2445	0.2534	−0.2521	0.7411
P1100	Selectivity increment	AVO; Bin5; 2022 *	0.1840	0.2548	−0.3154	0.6834
P1101	Selectivity increment	AVO; Bin6; 2022 *	0.1910	0.2550	−0.3088	0.6908
P1102	Selectivity increment	AVO; Bin7; 2022 *	0.2070	0.2551	−0.2930	0.7069
P1103	Selectivity increment	AVO; Bin8; 2022 *	0.2136	0.2312	−0.2396	0.6667
P1104	Selectivity increment	AVO; Bin2; 2023 *	0.2450	0.2591	−0.2628	0.7527
P1105	Selectivity increment	AVO; Bin3; 2023 *	0.2141	0.2631	−0.3016	0.7298
P1106	Selectivity increment	AVO; Bin4; 2023 *	0.0901	0.2626	−0.4245	0.6048
P1107	Selectivity increment	AVO; Bin5; 2023 *	0.1396	0.2622	−0.3742	0.6535
P1108	Selectivity increment	AVO; Bin6; 2023 *	0.1864	0.2622	−0.3274	0.7002
P1109	Selectivity increment	AVO; Bin7; 2023 *	0.2082	0.2629	−0.3070	0.7234
P1110	Selectivity increment	AVO; Bin8; 2023 *	0.2057	0.2401	−0.2648	0.6762
P1111	Selectivity increment	AVO; Bin2; 2024 *	0.2261	0.2708	−0.3046	0.7569
P1112	Selectivity increment	AVO; Bin3; 2024 *	0.1919	0.2739	−0.3449	0.7287
P1113	Selectivity increment	AVO; Bin4; 2024 *	0.0979	0.2727	−0.4366	0.6324
P1114	Selectivity increment	AVO; Bin5; 2024 *	0.1236	0.2711	−0.4077	0.6549
P1115	Selectivity increment	AVO; Bin6; 2024 *	0.1798	0.2705	−0.3504	0.7100
P1116	Selectivity increment	AVO; Bin7; 2024 *	0.2037	0.2727	−0.3309	0.7382
P1117	Selectivity increment	AVO; Bin8; 2024 *	0.1924	0.2513	−0.3001	0.6849
P1118	Selectivity increment	AVO; Bin2; 2025 *	0.2192	0.2804	−0.3304	0.7688
P1119	Selectivity increment	AVO; Bin3; 2025 *	0.2323	0.2862	−0.3287	0.7934
P1120	Selectivity increment	AVO; Bin4; 2025 *	0.1656	0.2871	−0.3970	0.7283
P1121	Selectivity increment	AVO; Bin5; 2025 *	0.1703	0.2867	−0.3916	0.7321

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1122	Selectivity increment	AVO; Bin6; 2025 *	0.1740	0.2869	−0.3883	0.7362
P1123	Selectivity increment	AVO; Bin7; 2025 *	0.1884	0.2866	−0.3734	0.7501
P1124	Selectivity increment	AVO; Bin8; 2025 *	0.1812	0.2649	−0.3380	0.7004
P1125	Selectivity increment	AVO; Bin2; 2026 *	0.2093	0.3050	−0.3886	0.8072
P1126	Selectivity increment	AVO; Bin3; 2026 *	0.2386	0.3046	−0.3585	0.8356
P1127	Selectivity increment	AVO; Bin4; 2026 *	0.1551	0.3058	−0.4443	0.7544
P1128	Selectivity increment	AVO; Bin5; 2026 *	0.1576	0.3063	−0.4428	0.7581
P1129	Selectivity increment	AVO; Bin6; 2026 *	0.1761	0.3067	−0.4251	0.7772
P1130	Selectivity increment	AVO; Bin7; 2026 *	0.1927	0.3064	−0.4078	0.7932
P1131	Selectivity increment	AVO; Bin8; 2026 *	0.1815	0.2852	−0.3775	0.7406
P1132	Log logistic slope	Asc; BTS	−0.8376	0.1962	−1.2221	−0.4531
P1133	Logistic midpoint	Asc; BTS	6.9805	0.6409	5.7244	8.2367
P1134	BTS age-1 log selectivity	Age-1 component; BTS	−3.0479	0.3067	−3.6490	−2.4469
P1135	BTS slope deviation	Asc; BTS; 1983	0.9087	0.2274	0.4630	1.3543
P1136	BTS slope deviation	Asc; BTS; 1984	0.7382	0.2286	0.2901	1.1863
P1137	BTS slope deviation	Asc; BTS; 1985	0.2475	0.2461	−0.2348	0.7297
P1138	BTS slope deviation	Asc; BTS; 1986	0.3667	0.2364	−0.0965	0.8300
P1139	BTS slope deviation	Asc; BTS; 1987	0.3474	0.2949	−0.2306	0.9254
P1140	BTS slope deviation	Asc; BTS; 1988	0.3002	0.3348	−0.3559	0.9564
P1141	BTS slope deviation	Asc; BTS; 1989	0.5436	0.3985	−0.2374	1.3245
P1142	BTS slope deviation	Asc; BTS; 1990	1.2073	0.3076	0.6043	1.8102
P1143	BTS slope deviation	Asc; BTS; 1991	1.0413	0.2581	0.5355	1.5471
P1144	BTS slope deviation	Asc; BTS; 1992	0.9879	0.2549	0.4883	1.4875
P1145	BTS slope deviation	Asc; BTS; 1993	1.1449	0.2594	0.6365	1.6532
P1146	BTS slope deviation	Asc; BTS; 1994	1.1404	0.2336	0.6827	1.5982
P1147	BTS slope deviation	Asc; BTS; 1995	1.0736	0.2332	0.6165	1.5307
P1148	BTS slope deviation	Asc; BTS; 1996	0.6664	0.2162	0.2428	1.0901
P1149	BTS slope deviation	Asc; BTS; 1997	0.5019	0.2354	0.0406	0.9632
P1150	BTS slope deviation	Asc; BTS; 1998	0.2949	0.2285	−0.1530	0.7427
P1151	BTS slope deviation	Asc; BTS; 1999	0.2464	0.2318	−0.2079	0.7008
P1152	BTS slope deviation	Asc; BTS; 2000	0.5104	0.2470	0.0264	0.9944
P1153	BTS slope deviation	Asc; BTS; 2001	0.3417	0.2233	−0.0959	0.7793
P1154	BTS slope deviation	Asc; BTS; 2002	0.7946	0.2240	0.3556	1.2337
P1155	BTS slope deviation	Asc; BTS; 2003	1.0199	0.2354	0.5585	1.4812

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1156	BTS slope deviation	Asc; BTS; 2004	0.9485	0.2234	0.5106	1.3864
P1157	BTS slope deviation	Asc; BTS; 2005	1.0394	0.2342	0.5803	1.4985
P1158	BTS slope deviation	Asc; BTS; 2006	0.7304	0.2221	0.2950	1.1657
P1159	BTS slope deviation	Asc; BTS; 2007	1.1604	0.2459	0.6785	1.6424
P1160	BTS slope deviation	Asc; BTS; 2008	1.3498	0.2211	0.9164	1.7832
P1161	BTS slope deviation	Asc; BTS; 2009	1.0541	0.2284	0.6065	1.5016
P1162	BTS slope deviation	Asc; BTS; 2010	1.5895	0.2493	1.1008	2.0781
P1163	BTS slope deviation	Asc; BTS; 2011	1.3653	0.2144	0.9450	1.7856
P1164	BTS slope deviation	Asc; BTS; 2012	1.0054	0.2373	0.5403	1.4705
P1165	BTS slope deviation	Asc; BTS; 2013	1.1035	0.2417	0.6297	1.5773
P1166	BTS slope deviation	Asc; BTS; 2014	1.3500	0.2227	0.9136	1.7864
P1167	BTS slope deviation	Asc; BTS; 2015	1.2379	0.2284	0.7902	1.6856
P1168	BTS slope deviation	Asc; BTS; 2016	1.2450	0.2421	0.7705	1.7195
P1169	BTS slope deviation	Asc; BTS; 2017	1.0643	0.2207	0.6317	1.4968
P1170	BTS slope deviation	Asc; BTS; 2018	0.5311	0.2347	0.0711	0.9911
P1171	BTS slope deviation	Asc; BTS; 2019	0.8316	0.2327	0.3755	1.2876
P1172	BTS slope deviation	Asc; BTS; 2020	0.7195	0.2985	0.1345	1.3046
P1173	BTS slope deviation	Asc; BTS; 2021	0.6483	0.2169	0.2232	1.0734
P1174	BTS slope deviation	Asc; BTS; 2022	0.8388	0.2317	0.3848	1.2929
P1175	BTS slope deviation	Asc; BTS; 2023	0.5333	0.2269	0.0887	0.9779
P1176	BTS slope deviation	Asc; BTS; 2024	0.5612	0.2527	0.0660	1.0564
P1177	BTS slope deviation	Asc; BTS; 2025	0.6074	0.2290	0.1586	1.0562
P1178	BTS slope deviation	Asc; BTS; 2026	0.8008	0.2609	0.2895	1.3121
P1179	BTS midpoint deviation	Asc; BTS; 1983	−0.3111	0.0866	−0.4808	−0.1414
P1180	BTS midpoint deviation	Asc; BTS; 1984	−0.1267	0.0895	−0.3021	0.0488
P1181	BTS midpoint deviation	Asc; BTS; 1985	0.0383	0.0942	−0.1463	0.2229
P1182	BTS midpoint deviation	Asc; BTS; 1986	0.0274	0.0975	−0.1637	0.2186
P1183	BTS midpoint deviation	Asc; BTS; 1987	−0.0750	0.1102	−0.2909	0.1409
P1184	BTS midpoint deviation	Asc; BTS; 1988	−0.2158	0.1232	−0.4573	0.0258
P1185	BTS midpoint deviation	Asc; BTS; 1989	−0.2963	0.1204	−0.5324	−0.0603
P1186	BTS midpoint deviation	Asc; BTS; 1990	−0.3506	0.1113	−0.5688	−0.1323
P1187	BTS midpoint deviation	Asc; BTS; 1991	−0.2509	0.1111	−0.4688	−0.0331
P1188	BTS midpoint deviation	Asc; BTS; 1992	−0.2448	0.1085	−0.4575	−0.0322
P1189	BTS midpoint deviation	Asc; BTS; 1993	−0.2921	0.1020	−0.4920	−0.0922

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1190	BTS midpoint deviation	Asc; BTS; 1994	−0.2291	0.0979	−0.4210	−0.0373
P1191	BTS midpoint deviation	Asc; BTS; 1995	−0.1343	0.1012	−0.3326	0.0640
P1192	BTS midpoint deviation	Asc; BTS; 1996	0.0653	0.0961	−0.1230	0.2536
P1193	BTS midpoint deviation	Asc; BTS; 1997	0.1110	0.1035	−0.0917	0.3138
P1194	BTS midpoint deviation	Asc; BTS; 1998	0.1916	0.1005	−0.0054	0.3885
P1195	BTS midpoint deviation	Asc; BTS; 1999	0.1329	0.1076	−0.0780	0.3439
P1196	BTS midpoint deviation	Asc; BTS; 2000	−0.0032	0.1111	−0.2210	0.2145
P1197	BTS midpoint deviation	Asc; BTS; 2001	0.0394	0.1009	−0.1583	0.2372
P1198	BTS midpoint deviation	Asc; BTS; 2002	−0.0939	0.1011	−0.2920	0.1042
P1199	BTS midpoint deviation	Asc; BTS; 2003	−0.1374	0.1034	−0.3402	0.0653
P1200	BTS midpoint deviation	Asc; BTS; 2004	−0.1130	0.0971	−0.3033	0.0773
P1201	BTS midpoint deviation	Asc; BTS; 2005	−0.1589	0.0977	−0.3504	0.0326
P1202	BTS midpoint deviation	Asc; BTS; 2006	−0.0355	0.0960	−0.2237	0.1527
P1203	BTS midpoint deviation	Asc; BTS; 2007	−0.2089	0.0993	−0.4035	−0.0143
P1204	BTS midpoint deviation	Asc; BTS; 2008	−0.2335	0.0961	−0.4219	−0.0451
P1205	BTS midpoint deviation	Asc; BTS; 2009	−0.1517	0.0988	−0.3454	0.0420
P1206	BTS midpoint deviation	Asc; BTS; 2010	−0.3339	0.0978	−0.5256	−0.1421
P1207	BTS midpoint deviation	Asc; BTS; 2011	−0.2094	0.0940	−0.3936	−0.0251
P1208	BTS midpoint deviation	Asc; BTS; 2012	−0.1593	0.0986	−0.3526	0.0339
P1209	BTS midpoint deviation	Asc; BTS; 2013	−0.1905	0.0946	−0.3760	−0.0050
P1210	BTS midpoint deviation	Asc; BTS; 2014	−0.2790	0.0959	−0.4670	−0.0911
P1211	BTS midpoint deviation	Asc; BTS; 2015	−0.3231	0.0976	−0.5144	−0.1318
P1212	BTS midpoint deviation	Asc; BTS; 2016	−0.2689	0.0984	−0.4617	−0.0762
P1213	BTS midpoint deviation	Asc; BTS; 2017	−0.1865	0.0941	−0.3709	−0.0020
P1214	BTS midpoint deviation	Asc; BTS; 2018	−0.0140	0.0995	−0.2090	0.1811
P1215	BTS midpoint deviation	Asc; BTS; 2019	−0.1490	0.0969	−0.3390	0.0410
P1216	BTS midpoint deviation	Asc; BTS; 2020	−0.0624	0.1177	−0.2930	0.1683
P1217	BTS midpoint deviation	Asc; BTS; 2021	0.0125	0.0977	−0.1790	0.2041
P1218	BTS midpoint deviation	Asc; BTS; 2022	−0.0671	0.0992	−0.2617	0.1274
P1219	BTS midpoint deviation	Asc; BTS; 2023	0.0978	0.0985	−0.0952	0.2909
P1220	BTS midpoint deviation	Asc; BTS; 2024	0.0068	0.1052	−0.1993	0.2130
P1221	BTS midpoint deviation	Asc; BTS; 2025	0.0317	0.1021	−0.1684	0.2319
P1222	BTS midpoint deviation	Asc; BTS; 2026	−0.1099	0.1145	−0.3344	0.1146
P1223	BTS age-1 deviation	Age-1 component; BTS; 1983	−0.0145	0.1177	−0.2452	0.2162

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1224	BTS age-1 deviation	Age-1 component; BTS; 1984	0.0371	0.1648	−0.2858	0.3600
P1225	BTS age-1 deviation	Age-1 component; BTS; 1985	−0.1923	0.1416	−0.4698	0.0851
P1226	BTS age-1 deviation	Age-1 component; BTS; 1986	−0.4013	0.1448	−0.6851	−0.1174
P1227	BTS age-1 deviation	Age-1 component; BTS; 1987	−0.2452	0.1911	−0.6198	0.1294
P1228	BTS age-1 deviation	Age-1 component; BTS; 1988	−0.2987	0.1852	−0.6616	0.0643
P1229	BTS age-1 deviation	Age-1 component; BTS; 1989	−0.0598	0.1804	−0.4134	0.2939
P1230	BTS age-1 deviation	Age-1 component; BTS; 1990	0.1030	0.1485	−0.1881	0.3941
P1231	BTS age-1 deviation	Age-1 component; BTS; 1991	−0.1431	0.1368	−0.4113	0.1250
P1232	BTS age-1 deviation	Age-1 component; BTS; 1992	−0.0165	0.1370	−0.2849	0.2520
P1233	BTS age-1 deviation	Age-1 component; BTS; 1993	0.0411	0.1297	−0.2131	0.2953
P1234	BTS age-1 deviation	Age-1 component; BTS; 1994	−0.1102	0.1451	−0.3946	0.1741
P1235	BTS age-1 deviation	Age-1 component; BTS; 1995	−0.1370	0.1474	−0.4259	0.1519
P1236	BTS age-1 deviation	Age-1 component; BTS; 1996	−0.1046	0.1174	−0.3346	0.1254
P1237	BTS age-1 deviation	Age-1 component; BTS; 1997	0.0048	0.1318	−0.2534	0.2631
P1238	BTS age-1 deviation	Age-1 component; BTS; 1998	0.0560	0.1447	−0.2275	0.3395
P1239	BTS age-1 deviation	Age-1 component; BTS; 1999	0.0971	0.1390	−0.1753	0.3695
P1240	BTS age-1 deviation	Age-1 component; BTS; 2000	0.1927	0.1380	−0.0778	0.4631
P1241	BTS age-1 deviation	Age-1 component; BTS; 2001	0.0776	0.1172	−0.1521	0.3073
P1242	BTS age-1 deviation	Age-1 component; BTS; 2002	0.0837	0.1210	−0.1534	0.3209
P1243	BTS age-1 deviation	Age-1 component; BTS; 2003	0.2335	0.1390	−0.0389	0.5059
P1244	BTS age-1 deviation	Age-1 component; BTS; 2004	0.0473	0.1386	−0.2242	0.3189
P1245	BTS age-1 deviation	Age-1 component; BTS; 2005	−0.0147	0.1483	−0.3054	0.2759
P1246	BTS age-1 deviation	Age-1 component; BTS; 2006	−0.0132	0.1166	−0.2417	0.2153
P1247	BTS age-1 deviation	Age-1 component; BTS; 2007	−0.0358	0.1179	−0.2668	0.1952
P1248	BTS age-1 deviation	Age-1 component; BTS; 2008	0.2024	0.1181	−0.0291	0.4340
P1249	BTS age-1 deviation	Age-1 component; BTS; 2009	0.3586	0.1115	0.1400	0.5773
P1250	BTS age-1 deviation	Age-1 component; BTS; 2010	0.3397	0.1320	0.0809	0.5984
P1251	BTS age-1 deviation	Age-1 component; BTS; 2011	−0.0718	0.1226	−0.3121	0.1685
P1252	BTS age-1 deviation	Age-1 component; BTS; 2012	−0.0599	0.1366	−0.3276	0.2078
P1253	BTS age-1 deviation	Age-1 component; BTS; 2013	0.2090	0.1341	−0.0539	0.4719
P1254	BTS age-1 deviation	Age-1 component; BTS; 2014	0.1700	0.1236	−0.0723	0.4123
P1255	BTS age-1 deviation	Age-1 component; BTS; 2015	0.0604	0.1400	−0.2139	0.3347
P1256	BTS age-1 deviation	Age-1 component; BTS; 2016	−0.0342	0.1521	−0.3322	0.2639
P1257	BTS age-1 deviation	Age-1 component; BTS; 2017	−0.0710	0.1368	−0.3392	0.1972

Complete parameter estimates and uncertainty						
ID	Group	Parameter context	Estimate	SE	95% lower	95% upper
P1258	BTS age-1 deviation	Age-1 component; BTS; 2018	−0.0545	0.1279	−0.3053	0.1962
P1259	BTS age-1 deviation	Age-1 component; BTS; 2019	0.2778	0.1171	0.0483	0.5074
P1260	BTS age-1 deviation	Age-1 component; BTS; 2020	0.1488	0.2105	−0.2638	0.5614
P1261	BTS age-1 deviation	Age-1 component; BTS; 2021	0.0198	0.1341	−0.2431	0.2826
P1262	BTS age-1 deviation	Age-1 component; BTS; 2022	0.0699	0.1373	−0.1992	0.3390
P1263	BTS age-1 deviation	Age-1 component; BTS; 2023	−0.0696	0.1428	−0.3494	0.2102
P1264	BTS age-1 deviation	Age-1 component; BTS; 2024	0.1744	0.1502	−0.1201	0.4689
P1265	BTS age-1 deviation	Age-1 component; BTS; 2025	0.0580	0.1478	−0.2317	0.3477
P1266	BTS age-1 deviation	Age-1 component; BTS; 2026	−0.0438	0.2085	−0.4524	0.3649
* A parameter is shared across additional positions. The full shared-position mappings are supplied in the numerical parameter table.						
The BTS age-1 component reuses native coordinates named Descending; it is a separate first-age component rather than a descending limb. Its baseline is log selectivity, and its annual deviations multiply that baseline log value. Midpoint deviations likewise multiply the baseline midpoint. The selectivity table combines these parameters and their uncertainty.						

Note. All estimated parameters and approximate 95% confidence intervals in Model 26.0, on their estimation scales. The parameter ID links to the complete numerical data and shared-parameter mappings. Recruitment and selectivity deviations are expressed on their fitted deviation or increment scales; the biological and selectivity schedules show their realized effects. The full precision data also supply natural-scale transformations where applicable.

Table D2 *Largest Pairwise Correlations Among Model 26.0 Parameters*

Strong parameter correlations		
Parameter 1	Parameter 2	Correlation
P0001: Recruitment level: EBS Pollock; R0	P0039: Recruitment deviation: EBS Pollock; 2001	−0.96375
P0001: Recruitment level: EBS Pollock; R0	P0051: Recruitment deviation: EBS Pollock; 2013	−0.96370
P0001: Recruitment level: EBS Pollock; R0	P0047: Recruitment deviation: EBS Pollock; 2009	−0.96198
P0001: Recruitment level: EBS Pollock; R0	P0052: Recruitment deviation: EBS Pollock; 2014	−0.96036
P0001: Recruitment level: EBS Pollock; R0	P0028: Recruitment deviation: EBS Pollock; 1990	−0.95676
P1133: Logistic midpoint: Ascending; BTS; Sex combined	P1207: BTS midpoint deviation: Ascending; BTS; Sex combined; 2011	−0.95670
P1133: Logistic midpoint: Ascending; BTS; Sex combined	P1213: BTS midpoint deviation: Ascending; BTS; Sex combined; 2017	−0.95529
P0001: Recruitment level: EBS Pollock; R0	P0040: Recruitment deviation: EBS Pollock; 2002	−0.95229
P0001: Recruitment level: EBS Pollock; R0	P0035: Recruitment deviation: EBS Pollock; 1997	−0.95002
P0001: Recruitment level: EBS Pollock; R0	P0045: Recruitment deviation: EBS Pollock; 2007	−0.94802

Note. Ten largest absolute pairwise correlations among Model 26.0 estimated parameters. IDs link to the complete parameter table; correlations close to plus or minus one identify parameter combinations that change together. These are parameter correlations rather than correlations between survey observations.

Appendix E
Proposed Projection Input Curves

The proposed three-year fishery-selectivity average is compared with the window including the inferred 2026 curve (Figure E1). The technical appendix records their conditional projection effects.



Note. Proposed three-year fishery selectivity and the alternative window check. The two curves are distinguished by colour, line type and point shape.

Figure E1 *Proposed Fishery Selectivity and Alternative Averaging Window*

The projected fishery weights retain their distinction from population and spawning schedules (Figure E2). The forecast year and the schedules held constant are specified in Section 7.



Note. Weight schedules used in the proposed projection. The 2027 fishery forecast is distinguished from 2026 fishery weights and from the carried population and spawning schedules.

Figure E2 *Weight Schedules for the Proposed Projection*

Appendix F
Model Equations

Scope. This appendix carries forward the structure and model components of the 2024 *Appendix EBS Pollock Model Description* (SAFE pp. 172–182). The equations describe the Model 23.0 ADMB comparison. The proposed Model 26.0 configuration and its weight schedules are described in Section 5 and Section 5.2. The historical Tier 1 material documents an available calculation; the December 2024 SSC adopted Tier 3a.

Dynamics

The statistical age-structured model combines the catch equation with annual survival and an accumulating terminal age group (Fournier and Archibald 1982; Hilborn and Walters 1992; Schnute and Richards 1995; McAllister and Ianelli 1997). Let t index years, $a=1,\ldots,A$ index ages, and $A=15$ denote ages 15 and older. $N_{t,a}$ is abundance at the start of the year, $F_{t,a}$ and $M_{t,a}$ are annual instantaneous mortality rates, and $Z_{t,a}$ is their sum. Catch at age and catch biomass are

$$\begin{aligned} Z_{t,a} &= F_{t,a} + M_{t,a}, \\ C_{t,a} &= N_{t,a}\frac{F_{t,a}}{Z_{t,a}}\left(1-e^{-Z_{t,a}}\right), \\ Y_t &= \sum_{a=1}^{A} C_{t,a} w^{f}_{t,a},\; p^{f}_{t,a} = \frac{C_{t,a}}{\sum_{b} C_{t,b}}. \end{aligned} \tag{2}$$

Here $w^{f}_{t,a}$ is fishery weight at age. Abundance in millions of fish multiplied by kg per fish gives biomass in kt. The population advances as

$$\begin{aligned} N_{t+1,1} &= R_{t+1}, \\ N_{t+1,a+1} &= N_{t,a}e^{-Z_{t,a}}, 1 \le a < A, \\ N_{t+1,A} &= N_{t,A-1}e^{-Z_{t,A-1}} + N_{t,A}e^{-Z_{t,A}}. \end{aligned} \tag{3}$$

The middle line applies to entry from each younger age; the final line adds surviving members of the plus group. This corrects the age subscript in 2024 Eq. 14.2, which printed $a-1$ on the right side while advancing a to $a+1$ on the left.

Fishing mortality separates annual intensity from selectivity at age:

$$F_{t,a} = s^{f}_{t,a}\mu_F e^{\epsilon_t}. \tag{4}$$

The assessment estimates selectivity coefficients with penalties on changes over time and on curvature across ages (Butterworth et al. 2003). In a general penalty notation, $\Delta_t \log s_{t,a}$ controls annual changes and $\Delta_a^2 \log s_{t,a}$ controls smoothness across ages. Their weights and allowed change years come from the run controls.

The bottom-trawl survey (BTS) uses logistic selectivity above age one and a separate age-one term. The implemented form is

$$\begin{aligned} s^{b}_{t,a} &= \left\{1+\exp\left[-k_t\left(a-a_{50,t}\right)\right]\right\}^{-1}, a>1, \\ \log s^{b}_{t,1} &= b_1\exp\left(d_t\right), \\ k_t &= \acute{k}\exp\left(u_t\right), a_{50,t} = \acute{a}_{50}\exp\left(v_t\right). \end{aligned} \tag{5}$$

This follows compute_selectivity() and the age-one assignment in Get_Selectivity(). It replaces the ambiguous slope/intercept and age-one expressions printed in 2024. The 2024 description reported a process variance of 0.04 for an earlier random-walk formulation. The retained controls instead select the penalty on changes in log-selectivity at the

specified ages, plus the separate age-one deviation penalty. Its coefficient is 8 for the squared first differences of d_t. The actual controls and source identify which parameter or log-selectivity penalty is applied.

Acoustic-trawl survey (ATS) selectivity is flexible across ages; the optional separate age-one index leaves ages 2–15+ in its composition likelihood. Constant survey catchability and time-varying selectivity represent separate model components.

Recruitment

Recruitment is abundance at age one. Female spawning biomass accounts for survival from the start of the year to spawning:

$$B_t = \sum_{a=1}^{A} N_{t,a} \exp\left(-\tau Z_{t,a}\right) \phi_a w_{t,a}^{s}, \tau = 0.25. \quad (6)$$

Here $w_{t,a}^{s}$ is spawning weight and ϕ_a includes both maturity and the female fraction. The assessment reads a maturity schedule and applies the female fraction internally; the one-sex SPM handoff reads maturity and halves total abundance. Applying the female fraction once preserves the same biomass definition. The spawning time is April 1 (month 4).

Annual recruitment deviations accompany a stock–recruit function, $R_t = f\left(B_{t-1}\right) \exp\left(\varepsilon_t\right)$, with log-scale deviation penalties and the implementation's bias adjustment. The Ricker option selected by SrType=1 is

$$f(B) = \frac{B}{\psi_0} \exp\left[\alpha_R \left(1 - \frac{B}{B_0}\right)\right], B_0 = R_0 \psi_0, \alpha_R = \log\left(\frac{4h_{ADMB}}{1 - h_{ADMB}}\right). \quad (7)$$

ψ_0 is unfished spawning biomass per recruit. The final expression is the source code's transformation of its parameter named steepness; it is retained as h_{ADMB} to identify that convention. The 2024 printed Ricker equation misplaced R_0 in the biomass ratio. The expression here follows SRecruit() and Get_Bzero() in the preserved template.

The alternative Beverton–Holt relationship (Francis 1992) is

$$f_{BH}(B) = \frac{B}{\alpha_{BH} + \beta_{BH} B}, \alpha_{BH} = \psi_0 \frac{1 - h}{4h}, \beta_{BH} = \frac{5h - 1}{4h R_0}. \quad (8)$$

For this alternative, h is the fraction of unfished recruitment at 20% of unfished spawning biomass. The 2024 controls fix $\sigma_R = 1$ and specify a symmetric beta prior with shape parameters 14.93209877 for the selected stock–recruit parameter. The earlier Beverton–Holt example in the 2024 appendix used a different prior; it remains an alternative rather than the selected 2024 setting.

Diagnostics

The historical replay diagnostic retains fitted recruitment anomalies while adjusting recruitment for the change in spawning biomass under a hypothetical history with fishing set to zero:

$$R_t^{'} = \hat{R}_t \frac{f\left(B_{t-1}^{'}\right)}{f\left(B_{t-1}\right)}. \quad (9)$$

Retrospective analyses repeat estimation after removing recent years and compare estimates for common years (Parma 1993; Ianelli and Fournier 1998). Their fit controls and terminal years accompany each saved run.

The effective number of ages contributing to spawning biomass is the exponential of Shannon entropy (Jost 2006):

$$D_t = \exp\left(-\sum_a p_{t,a}^{B} \log p_{t,a}^{B}\right), p_{t,a}^{B} = \frac{N_{t,a} e^{-\tau Z_{t,a}} \phi_a w_{t,a}^{s}}{B_t}. \quad (10)$$

The mean-age diagnostic uses observed and predicted composition means and the predicted variance at age (Francis 2011):

$$\acute{a}_t=\sum_a a\,p_{t,a},\hat{\acute{a}}_t=\sum_a a\,\hat{p}_{t,a},$$
$$V_t=\sum_a (a-\hat{\acute{a}}_t)^2\hat{p}_{t,a}, \qquad (11)$$
$$z_t=\frac{\acute{a}_t-\hat{\acute{a}}_t}{\sqrt{V_t/n_t}},v=\{Var_t(z_t)\}^{-1}.$$

The implied adjusted sample size is $v\,n_t$. This diagnostic supports evaluation of weighting; changing assessment inputs requires a recorded decision. The expression for V_t replaces the malformed κ_t formula in the printed 2024 appendix.

Parameter estimation

The objective adds negative log likelihoods, prior densities and selectivity penalties. Phase-based estimation introduces parameters in stages. For age compositions, the multinomial contribution, up to a data-only constant, is

$$\ell_{age}=-\sum_t n_t\sum_a p_{t,a}\log(\hat{p}_{t,a}+c),\hat{p}_t^{read}=\hat{p}_t^{true}E. \qquad (12)$$

c is the small numerical constant in the implementation. When enabled, $E_{a,b}$ is the probability that true age a is read as age b. The selected historical controls have use_age_err=0. The original appendix also described a robust composition option (Fournier et al. 1990), with variance inflation and a small outlier component. In the retained 2024 controls its activation phase is 1350; the six-phase completed comparison uses the multinomial form. This distinction prevents the general software description from implying that the robust option was active.

For survey s at within-year time τ_s, expected abundance and biomass at age are

$$\hat{N}^s_{t,a}=q_s s^s_{t,a}N_{t,a}e^{-\tau_s Z_{t,a}},\hat{I}^s_t=\sum_a \hat{N}^s_{t,a}w^s_{t,a}. \qquad (13)$$

An abundance index omits the weight multiplier. The original description used midyear survey timing; actual run inputs specify the timing used in each comparison. The AVO index shares ATS selectivity and its documented weight basis, while retaining a separate catchability coefficient.

The independent lognormal-index and multivariate BTS contributions have the forms

$$\ell_s=\frac{1}{2}\sum_t\frac{[\log I^{obs}_{s,t}-\log\hat{I}_{s,t}]^2}{\sigma^2_{s,t}},\ell_{BTS}=\frac{1}{2}r^T\Sigma^{-1}r,r_t=\log I^{obs}_t-\log\hat{I}_t. \qquad (14)$$

Fixed-variance normalization terms are constant for these comparisons. The normal-index option described in 2024 instead uses residuals on the original scale. The BTS covariance provides Σ for the multivariate lognormal fit. Catch biomass enters the implemented objective as $\lambda_C\sum_t[\log(Y^{obs}_t+c)-\log(\hat{Y}_t+c)]^2$, with $\lambda_C=200$ in the retained controls. This equals $1/(2\sigma_C^2)$ for $\sigma_C=0.05$ and removes the extra factor of two in the printed 2024 likelihood expression. Priors and penalties retain their actual source-code weights.

Uncertainty in mean body mass

The 2024 appendix described estimating future weights within the assessment to propagate uncertainty into derived quantities (Ianelli and Williamson 2001). Its earlier option used $w_a^{future}\sim N(\acute{w}_a,\sigma^2_{w,a})$. The retained implementation fits a mean growth curve plus cohort and year effects. For starting modeled age a_0, its mean curve is

$$\acute{w}_a = \alpha_w \left\{ L_1 + (L_2 - L_1) \frac{1 - K^{a-a_0}}{1 - K^{A-1}} \right\}^3. \quad (15)$$

At the final estimation phase, the recurrence in Est_Fixed_Effects_wts() is

$$\begin{aligned} \hat{w}_{t,a_0} &= \acute{w}_{a_0} \exp(\sigma_c^2/2 + \sigma_c c_t), \\ \hat{w}_{t,a} &= \hat{w}_{t-1,a-1} + (\acute{w}_a - \acute{w}_{a-1}) \exp(\sigma_y^2/2 + \sigma_y y_t), a > a_0. \end{aligned} \quad (16)$$

The initial year starts at $\acute{w}_a$; dataset-specific scale vectors map the latent weights to observation series. This notation corrects the increment index in 2024 and explicitly retains the implementation's sigma scaling and final-phase offsets. The cohort and year deviations receive standard-normal penalties. The four-term weight objective is

$$\ell_w = \frac{1}{2} \sum_{h,t,a} \frac{(w_{h,t,a}^{obs} - \acute{w}_a)^2}{\sigma_{h,t,a}^2} + \frac{1}{2} \sum_{h,t,a} \frac{(w_{h,t,a}^{obs} - \hat{w}_{h,t,a})^2}{\sigma_{h,t,a}^2} + \frac{1}{2} \sum_t c_t^2 + \frac{1}{2} \sum_t y_t^2. \quad (17)$$

The 2026 numerical baseline allocates all four terms and retains their sum.

Tier 1 projections: historical calculation

The 2024 description retained a Tier 1 calculation based on fishable biomass, the harmonic-mean harvest rate at MSY, and a spawning-biomass adjustment:

$$\begin{aligned} ABC_t &= B_{GM,t}^f \hat{u}_{HM} \zeta_t, \\ B_{GM,t}^f &= \exp\{\log \hat{B}_t^f - \sigma_{\log B^f}^2/2\}, \\ \hat{u}_{HM} &= \exp\{\log \hat{u}_{MSY} - \sigma_{\log u}^2/2\}, \\ \zeta_t &= \begin{cases} (B_t/B_{MSY} - 0.05)/0.95, & B_t < B_{MSY}, \\ 1, & B_t \geq B_{MSY}. \end{cases} \end{aligned} \quad (18)$$

The geometric and harmonic adjustments assume the documented lognormal approximations. This historical formula describes the earlier calculation over its applicable biomass range; final advice uses the adopted tier and the complete corresponding harvest-control rule.

Tier 3 projections

For fixed mortality, maturity, weights and selectivity, equilibrium spawning biomass per recruit at fishing intensity F is

$$\begin{aligned} l_1(F) = 1, l_a(F) &= l_{a-1}(F) e^{-M_{a-1} - F s_{a-1}}, 1 < a < A, \\ l_A(F) &= \frac{l_{A-1}(F) e^{-M_{A-1} - F s_{A-1}}}{1 - e^{-M_A - F s_A}}, \\ \psi(F) &= \sum_a l_a(F) \phi_a w_a^s e^{-\tau(M_a + F s_a)}, \\ B_{100\%} = \acute{R} \psi(0), B_{x\%} &= \frac{x}{100} B_{100\%}, \frac{\psi(F_{x\%})}{\psi(0)} = \frac{x}{100}. \end{aligned} \quad (19)$$

Tier 3a uses $F_{40\%}$ for maximum ABC and $F_{35\%}$ for OFL, with the adopted biomass-dependent rule applied when needed. Projection catch follows Equation 2. The 2024 SPM calculation used one sex category split equally into females and males, fixed biological schedules and selectivity, and stochastic recruitment generated from the historical arithmetic and harmonic means with an inverse-Gaussian distribution. Its 1,000 simulations retained recruitment process variability; uncertainty in fitted abundance, mean recruitment, selectivity and weights was omitted. The saved seed is 123 in the historical SPM source.

The recovered 2024 handoff uses five preceding selectivity years (2019–2023), April spawning, the terminal spawning weights and fitted future fishery weights. SPM averages age-one recruitment over 1978–2023; the assessment's $B_{100\%}$ calculation includes 2024. These different recruitment means produce different reference-point values. The selected scenario fixes catches at 1,350 kt in both 2025 and 2026. Its archived simulation means provide the historical forecast comparison.

Appendix G
FT-NIRS Age Prediction for EBS Pollock

Authored by: Derek Chamberlin, Esther Goldstein, Beth Matta, and Jason Conner

FT-NIRS Team: John Brogan, Jason Conner, Derek Chamberlin, Esther Goldstein, Brenna Hsieh, Beth Matta, Jon Short

FT-NIRS predicts age from otolith spectra and associated specimen information, using traditional microscopic ages for calibration and validation. The workflow proceeds from age prediction through tester-age conversion to the assessment likelihood (Table G1). The 2024 and 2025 fishery age compositions use their respective independent tester-age keys applied to released FT-NIRS predictions. The 2025 source is model a00-d3. The year-specific keys also convert biological weight inputs (Table G2). Both composition series use the assessment's identity ageing-error assumption after their selected age treatment.

Year-specific conditional keys redistribute source abundance and biomass among tester ages. Mean weights are then calculated as converted biomass divided by converted abundance, with terminal ages pooled using those totals. The partial 2024 fishery diagnostic retains unsupported source-age-13 mass (Table G3). Selected assessment weights fill that row using pooled age offsets and assign a 100% CV to the target-age-15+ weight observation. The 2025 conversion uses 668 independent tester pairs and fills the unsupported source-age-13 row with the midpoint of the age-12 and age-14 conditional probabilities. The same key supplies the 2025 age composition and weight observations (Table G4).

BTS 2025 and 2026 use separate tester-age keys. The 2026 key retains 307 pairs after excluding 22 flagged pairs, with exact TMA ages 1–18 converted before ages 15–18 are pooled into 15+. Total abundance and biomass are conserved. Converted weight SEs propagate the saved joint abundance–weight covariance (Table G5). The fitted weight model retains its existing marginal observation likelihood, and empirical-key uncertainty is excluded. Numerical checks distinguish fitted use from the retained full-pair conversion diagnostics (Table G6).

Provisional 2026 ATS ages use the BTS age–length key (ALK) applied to ATS population length-frequency estimates, followed by conversion to the TMA basis. FT-NIRS ages from the 2026 ATS samples will be available in 2027. Unsupported lengths remain unallocated. The resulting weights also supply the 2026 AVO predictions.

Fourier transform near-infrared spectroscopy (FT-NIRS) Overview

Fourier transform near-infrared spectroscopy (FT-NIRS) has emerged as an effective secondary age prediction tool and has successfully been implemented in production ageing settings (Matta and Helser, 2024), with FT-NIRS predicted ages recently being used to produce stock assessment age composition data (Dick et al., 2025; Ianelli, 2025). FT-NIRS is a non-destructive technique that interrogates a sample using light in the near-infrared region of the light spectrum (typically 4,000 to 12,000 cm^{-1}).

Scanning with FT-NIRS produces a spectral signature of the sample based on the absorption of near infrared light by different chemical bonds, such as C-H (aliphatic), C-H (aromatic), C-O (carboxyl), O-H (hydroxyl), and N-H (amine and amide) (Murray and Williams, 1987). For otoliths, these spectra capture changes in the organic protein matrix (otolin) relative to the inorganic calcium carbonate structure. This relationship correlates strongly with fish growth and age (Passerotti et al., 2022) and has been used to develop models, calibrated with traditional microscopic ages (TMA) and FT-NIR spectra, to predict the age of unknown samples based on their spectral signatures with an estimated time-cost saving of 23-40% (Adamack et al., 2025). FT-NIRS has been applied to estimate age for a multitude of fish species from Australia (Wedding et al., 2014), the Mediterranean Sea (Basilone et al., 2024), the eastern Bering Sea (Benson et al., 2023), and the Atlantic Ocean (Adamack et al., 2025).

Traditional Microscopic Ageing (TMA) QA/QC

The FT-NIRS models rely on traditional microscopic ageing (TMA) to calibrate the model, thus the accuracy of the age predictions is directly impacted by the accuracy of the calibration ages (Chamberlin et al., 2026). The AFSC Age and Growth Program (AGP) has a long history of ageing walleye pollock by microscopic examination of otolith growth patterns (Matta et al., 2026) and has employed a rigorous quality control program since 1983 (Kimura and Anderl, 2005). In TMA of pollock, presumed annual marks ("annuli") are enumerated from the otolith surface in younger fish or from burned or toasted otolith cross sections in older fish (Matta and Kimura, 2012; Matta et al., 2026). The accuracy of the

AGP's TMA for pollock has been independently corroborated using a variety of methods, including radiometric ageing, edge and marginal increment analysis, year-class tracking, and captive-rearing studies (Kastelle and Kimura, 2006; Kimura et al., 2006; Kimura et al., 2007; Lai, 1985).

To test FT-NIRS model prediction skill, random 20% subsamples of pollock otoliths from each collection source (fishery, bottom-trawl survey, and hydroacoustic midwater trawl) were subjected to TMA. Spectral outliers (i.e., those that predicted fish lengths outside an established threshold) were also selected for TMA; see FT-NIRS QA/QC below for more details. Each otolith selected for TMA was independently aged by two experienced readers. The TMA subsamples were evaluated for inter-reader precision and bias. All discrepancies between the two readers were resolved. Any TMA age-length outliers were reexamined, and re-aged or recorded as collection errors when appropriate (Kimura and Anderl, 2005).

FT-NIRS QA/QC

In addition to the TMA QA/QC, the FT-NIR spectra underwent a three-step QA/QC process to identify poor quality spectra due to data collection errors, instrument issues, or otolith irregularities such as crystallization, contamination, or fragmentation (Goldstein et al., 2024). Spectrometer operators conducted qualitative QA/QC concurrent with data collection to identify missing data and unusual spectra using length predictions from a species- and region-specific partial least-squares regression (PLSR) model in comparison to observed lengths. Real-time length predictions were used to identify transcription errors and provide assurance of proper instrument function.

Following data collection, rigorous QA/QC was conducted to identify outliers using principal component analysis (PCA) and PLSR models developed from a multi-year collection of spectral data from each data source (AFSC groundfish survey, AFSC hydroacoustic survey, fishery data) and projections (PCA) and predictions (PLSR) of new data. PCA scores by scan date were used to visually identify instrument drift. Gross spectral data outliers were identified from PCA using Hotelling's T^2 and Q residuals (Pomerantsev and Rodionova, 2014) and PLSR outlier analyses identified discrepancies between spectra and specimen length by combining Hotelling's T^2 and Q residuals (full X-distance) and Y-distance (observed vs. predicted; Rodionova and Pomerantsev, 2020). Analyses used a threshold of 1% to identify outliers. Outliers were then aged using traditional microscopic methods. Once aged, outlier spectra were excluded from the final dataset if age predictions from the MMCNN models were also outside the MMCNN 95% Monte Carlo dropout prediction intervals discussed below.

FT-NIRS Model

Age was predicted from FT-NIR spectra using a multimodal convolutional neural network model (MMCNN). MMCNNs identify non-linear relationships in spectral data, incorporate additional inputs such as fish length or capture location, and learn relevant spectral features during training. Classical chemometric approaches such as PLS require the user to select key wavenumbers (Chen and Wang, 2019). The MMCNN used here is an updated version of the MMCNN presented in Benson et al. (2023) and Chamberlin et al. (2026). The MMCNN includes one convolutional layer and six dense layers to extract information from spectral and associated fish length data to predict age.

The first layer of the MMCNN is the spectral input data which is fed to a 1D convolutional layer, which acts as an automated feature extractor to identify patterns in the spectra. This output is then processed with a Rectified Linear Unit (ReLU) activation function to introduce nonlinearity into the network necessary for approximating continuous functions (Brownlee, 2019; Géron, 2019). The data are then flattened into a one-dimensional array and passed through a dense layer (all dense layers use Exponential Linear Unit (ELU) activation). This reduces the dimensionality of the spectral input and balances its contribution relative to the metadata. The metadata input is fed to a dense layer, then concatenated with the spectral input layer. The output is passed through a fully connected dense layer with ELU activation to model the interactions between spectral data and metadata. To mitigate overfitting, 5 dropout layers follow each dense layer to randomly ignore a subset of neurons and prevent codependency between nodes (Srivastava et al., 2014). Crucially, these dropout layers remain active during the inference phase, a technique known as Monte Carlo (MC) dropout, allowing the model to perform multiple forward passes per sample to generate a distribution of age estimates and calculate Monte Carlo prediction intervals. Finally, a fully connected dense layer with a single node and a batch normalization layer are used to produce a continuous age estimate, allowing for loss estimation via a Mean Absolute Error (MAE) function. The model's hyperparameters (the number of filters in the convolutional layer, the number of nodes in the concatenated layer,

learning rate, batch size and the dropout rate) were optimized using hyperband optimization (Li et al., 2018) with early stopping.

The walleye pollock model is trained on spectra and TMA ages from 2014-2025 using an "n−1" annual model updating approach. Under this scheme, spectral and age data from all fleets up to the year prior to the target prediction year are included (for example, predicting 2025 fishery ages uses data from 2024 and earlier). For MMCNN training, input data are first split into development (80%) and testing (20%) pools using a randomly shuffled split that is stratified on fleet and year. Samples receive a normalized weighting score based on 2 coefficients: *decay* is an integer [1, 2, 3] that defines the rate of exponential decay for samples based on year, and *alpha* [0, 0.1, 0.2, …, 1.0] defines whether sample year or sample age frequency has more influence in weighting the sample, with *alpha* = 1 turning decay off. This allows for testing of whether model performance improves when more recent calibration data or rare age classes are weighted more heavily. For hyperparameter optimization, the development data are further split into training (90%) and validation (10%) datasets to inform early stopping, using the previously defined sample weights. For each scenario a series of 10 trainings, with random seeds, are run to estimate the median number of epochs needed to train the model but avoid overfitting, prior to a final training run based on the median number of epochs.

In total, 31 training models were produced with a grid search for the 2 sample weighting coefficients, and age predictions were generated for bottom trawl survey, hydroacoustic survey and fishery fleets. The final candidate model for each fleet was selected using four statistical metrics: Lin's concordance correlation coefficient (Lin's CCC), the mean Winkler score, R^2, and the slope of the regression between FT-NIRS predicted ages and corresponding TMA ages. Lin's CCC measures how closely the observed data aligns with a 45° line of perfect agreement, while the mean Winkler score evaluates average performance based on the width of the Monte Carlo prediction intervals and the frequency of samples falling outside them. The four metrics capture different aspects of model accuracy, and visual assessments complemented them. Ultimately, the best-performing model was identified by synthesizing these visual and quantitative criteria. Age predictions were then finalized by removing any samples that were both flagged as a spectral outlier and an outlier from the Monte Carlo prediction intervals. This finalized dataset was then transmitted to end users and coded as FT-NIRS predicted ages.

Age conversion and assessment use

The assessment uses year-specific independent tester TMA conversions for the 2024 and 2025 fishery compositions, matching annual weight conversions and separate survey-specific age conversions (Table G1). These treatments serve different purposes: predicting age from spectra, allocating abundance and biomass among ages, and specifying the assessment likelihood are separate steps. Current input and fit records establish their use; the retained partial-conversion diagnostics describe their original support and uncertainty limits.

The 2026 BTS conditional tester-age key excludes calibration pairs marked nir_outlier = Y, removing 22 of 329 complete pairs and retaining 307. It targets independent test_age conditional on FT-NIRS final_nir_age; each FT-NIRS row is normalized by its retained paired count. Conversion uses exact TMA ages 1–18 before pooling ages 15–18 into 15+. Screening applies to calibration pairs, preserving the released abundance vector. All unflagged complete pairs are eligible, including checks outside the systematic 20% sample.

Screening changed the exact mean converted age from 5.318978 to 5.316448 years, with a largest age-proportion change of approximately 0.124 percentage points and unchanged total abundance. The same 307-pair key supplies BTS weight means and their joint-covariance delta-method SEs, provisional ATS age allocations and biomass-derived weights, and the 2026 AVO prediction weights. The standalone RE-WAA refit and assessment fits using these inputs are complete S10. The 2025 BTS key is unchanged. The original 329-pair products remain historical evidence. Empirical-key uncertainty is excluded; the fitted weight model retains its existing marginal observation likelihood rather than using the full converted covariance as a new likelihood.

The contribution describes **31 MMCNN training candidates**, distinguished by sample-weight settings and fleet-specific selection. PCA and PLSR provide spectral QA/QC; PLSR also provides length predictions. The a00-d3 and a03-d1 labels on the saved 600-fish comparison identify prediction sets, separately from the assessment's cumulative 0.x input sequence.

Table G1 *FT-NIRS Use and Transformations by Data Source and Year*

Gear or survey	Collection year	Input or model version	Age basis and year-specific key	Application and verified status
Commercial fishery, pelagic trawl (NORPAC gear 2)	2024	Saved sampler_2024_2025_20261016T160737Z; current fishery CAA	Released FT-NIRS ages; distinct 2024 key recovered from the earlier release: 2,985 released records and 693 independent tester pairs. Specimen/age-field hashes match the original-release verification record.	The ordinary assessment uses the independent tester TMA composition key, with the source-age-13 row completed from pooled 2024 source-age offsets at ages 9–12 (34 pairs) and likelihood N=578. The selected weight input uses the same key and a 100% CV for the target-age-15+ observation. The retained partial weight diagnostic describes its historical supported-only scope. The accepted marginal Model 26.1.2 fit uses the same year-specific composition and weight inputs. S2, S3, S6, S10.
Commercial fishery, pelagic trawl (NORPAC gear 2)	2025	FT-NIRS model a00-d3; annual catch-at-age expansion	Independent tester TMA ages; year-specific conditional key from 668 complete pairs. The unsupported source-age-13 row is the midpoint of the age-12 and age-14 rows.	Converted proportions use ages 1–14 and 15+ with selected likelihood N = 530 and identity ageing error. Total numbers are preserved. The matching weight observations use the same key. Key and interpolation uncertainty are additional to sampling uncertainty. S1, S2, S3, S10.
Commercial fishery, pelagic trawl	2025	Matching annual weight observations	Same independent tester-age key and source-age-13 midpoint. Source age 1 has zero abundance.	The key redistributes abundance and biomass in each of 1,000 saved bootstrap draws; weight ratios and SDs are calculated from the converted quantities. These observations supply the selected fitted weights. Target ages 13 and 14 have no converted abundance. Empirical-key and interpolation uncertainty are excluded. S3, S10.
Bottom-trawl survey (BTS), EBS/NBS	2025	Revised tinyVAST_abundance.csv / tinyVAST_props.csv; saved bts_2026_candidate lineage	FT-NIRS age estimates; bts_2025 independent tester key, dated 2025-09-24, 390 pairs.	Converted abundance and age proportions are used in the working assessment. Exact TMA ages 1–18 are converted before ages 15–18 are pooled into 15+. This key transfer supplies no additional abundance SE or empirical-key uncertainty. The earlier candidate exclusion is retained only in the historical staging record. S1, S4, S6, S10.

Gear or survey	Collection year	Input or model version	Age basis and year-specific key	Application and verified status
Bottom-trawl survey (BTS), EBS/NBS	2026	Revised spatial estimates; screened 29 October rerun input	Independent test_age conditional key from the 2026-09-24 release: 329 complete pairs before screening, 307 after excluding 22 nir_outlier = Y pairs.	Screened key applied to working CAA and WAA inputs. Exact TMA ages 1–18 are converted before 15+ pooling; source abundance, sample-size choices and 2025 key are retained. Original full-pair products are retained unchanged. S1, S4, S10.
Bottom-trawl survey, Indivero/VAST weight products	2025 and 2026, converted separately	Saved VAST joint N/W covariance; 29 October screened-key preparation and standalone RE-WAA refit	The 2025 full tester key is retained; 2026 uses the screened 307-pair key with the matching VAST abundance and weight estimates.	Ratios and delta-method SEs use abundance–weight and cross-age/year covariance under the same method. Nine 2026 observation means and nine SEs change; 2025 observations remain unchanged. The standalone RE-WAA numerical checks pass. Empirical-key uncertainty is excluded. S3, S5, S10.
BTS historical age-comparison material	2014–2018	FTNIR Compare.qmd uses historical fishery/survey paired releases	Historical comparison data span 2014–2018. The new contribution describes training data spanning 2014–2025 with annual n−1 updating.	These are distinct vintages. The historical comparison files alone do not establish the precise training set/model ID used for each 2025/2026 release; do not transfer old keys to new years. S7, supplied contribution above.
Acoustic-trawl survey (ATS), acoustic expansion with trawl biological samples	2024	ATS_2024_age comp ALK.xlsx, length totals × age-length proportions	The preparation record quotes the supplied 2025-09-08 email as TMA ages sent to MACE. The workbook has a 2024 age-length pivot and matrix multiplication of expanded numbers/biomass.	This local workbook verifies a length-to-age step, not an automatic FT-NIRS designation. A separate ATS age-age conversion and exact predictive model/key lineage are unverified here. S3, S8.
Acoustic-trawl survey (ATS)	2026	ATS/ stock_assessment_output/ *202604.csv; 29 October provisional rerun products	Released BTS FT-NIRS length key followed by the screened 307-pair BTS conditional tester-age key; source FT-NIRS allocations and unsupported lengths are preserved.	Abundance and biomass are redistributed with the same key; pooled ratios supply ATS and 2026 AVO prediction weights. Provisional estimates are used in the completed working fits and await direct ATS age-composition data in 2027. Transfer assumptions and uncertainty scope remain unchanged. S9, S10.

Note. FT-NIRS use and transformations by collection and year. Current fitted applications are distinguished from retained historical preparation diagnostics. Unfiltered counts describe the original complete released tester pairs; the 2026 screened input excludes the 22 flagged complete pairs.

The original keys contain 693 and 668 pairs for fishery 2024 and 2025, and 390 and 329 for BTS 2025 and 2026 before screening. The current 2026 BTS key retains 307 pairs. The fishery gear label follows the saved NORPAC records: gear 2 is PELAGIC in both 2024 and 2025. The contribution's fishery, bottom-trawl survey and hydroacoustic categories retain their own age basis and sampling conditions.

Predictive model and assessment input differences

The MMCNN combines spectral and fish-length information, MC dropout prediction intervals, hyperband optimization and 31 sample-weight settings. Its annual n−1 updating rule uses information available before the prediction year; for

example, a 2025 prediction uses data through 2024. The selected fleet-by-year model checksum, coefficient setting and exact mapping to the saved 600-fish prediction files remain to be supplied.

The full tester-key release uses final_nir_age as predictor and independent test_age as target, excluding resolved final_age as the calibration target. It contains 3,000 fishery-2025 records, 1,638 BTS-2025 records and 1,586 BTS-2026 records; 2,992, 1,619 and 1,582 have released FT-NIRS ages. Their complete-pair counts are 668, 390 and 329, respectively; the screened 2026 BTS application retains 307 S1.

For the 2025 fishery, model a00-d3 FT-NIRS ages are converted to independent tester ages using the 668 complete paired observations. The conditional probability of tester age j given FT-NIRS age i is $K_{ij}=n_{ij}/\sum_{j} n_{ij}$. Converted numbers are $N_j^{TMA}=\sum_{i} N_i^{FTNIR}K_{ij}$. Empirical rows retain the released paired observations, including flagged pairs. The unsupported source-age-13 row is $K_{13,j}=(K_{12,j}+K_{14,j})/2$, giving probabilities of one third at tester age 10, one sixth at age 11 and one half at age 15. The age-12 row has six pairs and the age-14 row one pair, so this interpolation has limited direct support. Source and tester ages in the release are exact ages 2–15; the assessment terminal group contains ages 15 and older. The selected likelihood sample size is 530. The key is treated as fixed; empirical-key and interpolation uncertainty are excluded.

The 2024 fishery composition uses its 693-pair independent tester key. The unsupported source-age-13 row uses pooled TMA-minus-FT-NIRS offsets from the same year's source ages 9–12 (34 pairs). Empirical rows remain unchanged; the selected likelihood input is 578. Both recent years are converted before fitting and use a 15-by-15 identity ageing-error matrix, equivalent to the ADMB setting with ageing error switched off. Key and tail-allocation uncertainty are additional to the sampling precision represented by the selected likelihood weights.

Weight conversion applies the same conditional key to source numbers and biomass, then divides converted biomass by converted numbers. Terminal-age weights use pooled totals. The partial empirical 2024 diagnostic leaves unsupported source-age-13 mass unallocated. The selected weight treatment fills that row using pooled offsets from source ages 9–12 and assigns a 100% CV to the target-age-15+ observation. Those fixed approximations add no empirical calibration pairs; their uncertainty differs from a sampling SE. The 2025 weight observations use the same source-age-13 midpoint as the age composition. Their zero-abundance source-age-1 padding contributes no converted numbers or biomass. The midpoint is specific to 2025; source age 14 also lacks pairs in the 2024 key.

The older staging record S6 describes a 2024 endpoint in pm_26_candidate.dat: fishery CAA at 0.4, provisional BTS WAA/biomass at 0.5/0.6, ATS at 0.7, and separate AVO-only 0.8/0.9 components. These are historical input-sequence labels from the incremental-model plan. Current fitted use follows the documented working input and run records, not those earlier staging exclusions.

Source and validation limits

The partial 2024 empirical diagnostic retains its unsupported source-age-13 mass. Selected assessment weights use the pooled-offset rule and 100% CV described above. The 2024 and 2025 compositions use their respective tester-age keys and selected support rules. Weight SEs propagate the available joint abundance–weight covariance, while the fitted model retains its marginal observation likelihood. Empirical-key uncertainty remains excluded.

The assessment author identified a00-d3 as the selected 2025 fishery model. The exact training-run checksum, coefficient setting and complete collection-by-year release record remain to be documented. The 31 predictive candidates and the assessment's cumulative input stages identify different choices. The earlier mixed-basis BTS workbook remains unavailable; the separately identified current workbook is excluded from assessment assembly. Its source basis must be established before using it in a comparison.

Provisional ATS products retain their transferred BTS age–length and tester-age keys, unsupported-length limits and separate sampling conditions. Direct ATS age and weight validation remains open, with age-composition data expected in 2027. Archived full-pair calibration products remain available alongside the screened-key input records. Prior fit, projection and posterior records retain their original dates and input identities. Final model and recommendation selection remain separate review decisions.

Sources and reproducibility

The paired-age releases, preparation products and current run records retain their source identities. Detailed file locations, checksums and the original preparation narrative are preserved in the technical source record.

S1 — Paired tester-age release: the full year-specific tester-key workbook and tidy release, with independent tester ages as the calibration target.

S2 — Fishery observations: catch-expanded annual numbers based on released FT-NIRS ages, with year-specific tester-age conversions: 693 pairs and N=578 for 2024; 668 pairs and N=530 for 2025.

S3 — Weight conversion: year-specific fishery conversion keys, partial 2024 diagnostics and independent fishery arithmetic checks.

S4 — Survey age compositions: revised BTS abundance and proportion estimates, with their source age assignments.

S5 — Survey weights and covariance: matching model-based abundance and weight estimates, joint covariance and delta-method validation.

S6 — Historical staging: the earlier preparation sequence, retained separately from current selected input and fit records.

S7 — Earlier age comparison: fishery and survey comparisons for 2014–2018, distinct from later training and prediction releases.

S8 — ATS length-to-age calculation: the 2024 age–length workbook and expanded number/biomass calculations.

S9 — ATS survey products: regional survey length and biomass products used for the provisional BTS-key transfer.

S10 — Current screened-key inputs and fits: the 307-pair calibration, completed standalone RE-WAA refit and current assessment run records. The original 329-pair products retain their historical identity.

Year-specific weight-conversion diagnostics

The retained full-pair conversion record separates three completed year-specific conversions from the partial 2024 fishery diagnostic (conversion-status table). Fishery 2025 and BTS 2025–2026 use independent tester-age keys; positive unsupported source-age-13 mass remains unallocated in fishery 2024.

Mean weights generally increase with age, with the largest source-versus- converted differences concentrated at older ages (2024 fishery weight table). The partial fishery 2024 values at TMA ages 13 and 14 are both 1.2878 kg, while its target 15+ weight remains missing. Its prepared older-age SEs are large relative to the means; the partial converted SDs condition on the fixed key and supported source ages.

The 2025 fishery conversion raises the 15+ mean from 0.8102 to 1.2676 kg; target ages 13 and 14 retain missing weights because converted abundance is zero (fishery weight comparison). Thus the older-age differences include changes in support as well as in ratios.

BTS 2025 converted weights are generally below the source weights from age 5 upward; in 2026, weights at ages 3–8 remain close to or above their source values, while older-age weights are lower (BTS weight comparison; Figure G1). The 15+ mean changes from 1.7646 to 1.5474 kg in 2025 and from 1.8390 to 1.3665 kg in 2026. These are conversion contrasts that vary by age and year. The smaller conditional SEs in several converted cells retain the stated fixed-key uncertainty scope.

Independent fishery arithmetic, preservation of earlier fishery values and survey covariance/derivative checks establish computational consistency (conversion checks). The partial 2024 diagnostic leaves unsupported source-age-13 mass unallocated. Current selected weights use the separate pooled-offset tail treatment and 100% CV, and the screened-key standalone RE-WAA and assessment fits are complete. These distinctions preserve the historical diagnostic's support limits and the current fit's fixed-key uncertainty scope.

Adjustments to FT-NIRs for assessment application

The following tables retain the full-pair calibration used to evaluate conversion arithmetic, including the partial 2024 fishery results. Current assessment inputs use the separately documented selected tail treatment and screened 2026 key. The table values preserve their original calibration basis.

Complete conversion report, including Table 3 for fishery 2024–2025

Diagnostic basis and current use

The retained diagnostics convert fishery **2025** and BTS **2025–2026** abundance and biomass with their year-specific independent tester-age keys. Fishery **2024** has a distinct 693-pair key, but its empirical partial conversion leaves positive source-age-13 mass unallocated. The selected assessment weight treatment fills that row using pooled offsets from source ages 9–12 and assigns a 100% CV to the target-age-15+ weight observation. The current standalone RE-WAA refit and assessment fits are complete; the historical partial diagnostic remains separate. The BTS tables below retain the original full-pair calibration, while the current 2026 fit uses the screened 307-pair key.

Table G2 *Conversion Status by Data Source*

Stream	Conversion key	Current action
Fishery 2024	Distinct 2024 key: 693 pairs	Partial diagnostic; source age 13 unallocated
Fishery 2025	2025 fishery tester-age key: 668 pairs	Converted with two approved probability rules
BTS 2025	2025 BTS tester-age key: 390 pairs	Converted with joint model uncertainty
BTS 2026	2026 BTS tester-age key: 329 pairs	Converted with joint model uncertainty

Note. Conversion status by new data source.

Conversion and approved approximations

Each key gives the distribution of TMA tester age conditional on FTNIR age. Abundance and biomass are redistributed together, using the same conditional probabilities. Converted mean weight is converted biomass divided by converted abundance. Within a source FTNIR bin, this calculation assigns the same source mean weight to every target age.

For fishery 2025, Jim approved two rules for unsupported source ages:

> **Age 1:** a dummy identity row assigns FTNIR age 1 to TMA age 1. All saved source age-1 catch estimates are zero, so this row retains zero catch.
>
> **Age 13:** each target-age probability is the arithmetic average of its probabilities in the source age-12 and age-14 rows. The resulting row sums to one. This approximation uses the existing age-12 and age-14 pairs.

Converted target-age 13 and 14 abundances are zero in the 2025 fishery key. Their weights and SEs remain paired missing observations. The historical data continue to inform these ages in the future wtAgeRE fit. All pre-2025 fishery mean/SE decimal values are preserved exactly.

Exact BTS target ages 1–18 are retained during conversion. Target ages 15–18 are then pooled into 15+, conserving total abundance and biomass. The 2024 fishery key also retains exact target ages through 18; the 2025 fishery key has target ages through 15. The same terminal pooling convention applies. Age selection for wtAgeRE occurs after conversion; its input bins are 3–15+.

Fishery 2024 results and remaining gap

The recovered 2024 release contains **2,985 released FTNIR ages and 693 complete independent tester pairs**. The target is test_age; the resolved final_age is a separate field. Canonical specimen and age-field hashes match the independent original-workbook verification record. The original workbook itself remains outside this local snapshot. The empirical key uses all complete released pairs, including additional QA/QC selections; their representativeness remains an assumption.

Source ages **1–12** have empirical probability rows. Source ages **13–15** lack pairs; only age 13 has positive mass in the saved sampler draws. Its unallocated observed-point mass is **0.0323% of abundance** and **0.0817% of biomass**.

Allocated plus unallocated totals balance across all 1,001 saved rows to relative errors below 6.7e-16. The diagnostic retains that mass separately and preserves the original totals.

The table and figure report the **partial, fixed-key bootstrap mean and SD** from the supported source rows. Every converted 2024 bin remains conditional on the unresolved source-age-13 allocation. Target TMA age 15+ receives zero allocated abundance, so its mean and SD remain missing. Source FTNIR age 13 and target TMA age 13 are distinct: supported source ages can populate target ages 13 and 14.

The 2025 midpoint rule stays specific to its approved key. In 2024, source age 14 also lacks calibration pairs; applying the age-12/14 midpoint rule would require additional empirical support. The 2024 key supports source age 1 directly; its positive source catch is retained.

Table G3 *Fishery 2024 Weights and Partial Supported-Only Conversion*

Age	Prepared mean	Prepared SE	Partial TMA mean	Partial TMA SD
3	0.3832	0.0093	0.4353	0.0053
4	0.4841	0.0067	0.4807	0.0033
5	0.5317	0.0035	0.5596	0.0021
6	0.6628	0.0040	0.6175	0.0023
7	0.8195	0.0166	0.6965	0.0060
8	0.9733	0.0439	0.8252	0.0138
9	0.9850	0.0469	1.0162	0.0182
10	1.1167	0.0401	1.0163	0.0227
11	1.3455	0.0774	1.0774	0.0227
12	1.2878	0.1082	1.2464	0.0455
13	0.7485	0.7800	1.2878	0.1082
14	1.1658	0.6356	1.2878	0.1082
15+	0.9705	0.9011	Missing	Missing

Note. Fishery 2024 weights in kilograms: preserved prepared values and partial supported-only conversion. SDs are conditional fixed-key diagnostics; all converted bins remain incomplete.

Weight and uncertainty comparisons



Note. Mean weights before and after conversion, with source SEs and converted bootstrap SDs or survey SEs. Fishery 2024 shows a partial supported-only diagnostic; source-age-13 mass remains unallocated.

Figure G1 *Mean Weights Before and After Conversion*

Converted fishery weights and bootstrap SDs are recalculated from the first 1,000 paired sampler bootstrap draws after transforming each draw's abundance and biomass. The final observed-data point row is excluded from those summaries. The previously selected legacy convention is retained: a target age absent in a bootstrap draw has zero weight for that draw. The SD uses the sample denominator. Known limitations of the legacy sampler's resampling selectors remain in the source record.

BTS uncertainty uses the saved joint VAST parameter covariance and conditional random-effect covariance. It includes dependence between source abundance and weight, among ages through shared parameters, and between 2025 and 2026. The ratio's numerator and denominator both contribute to the delta-method SE. The calculation evaluates the saved VAST model at its recorded parameter estimates.

All conversion keys are treated as fixed. The reported SEs exclude uncertainty in the empirical keys, the dummy row, and the interpolated row.

Table G4 *Fishery 2024–2025 Weights and Uncertainty*

Year	Age	Original mean	Original SE	TMA mean	TMA SE / SD
2024	3	0.3832	0.0093	0.4353	0.0053
2024	4	0.4841	0.0067	0.4807	0.0033
2024	5	0.5317	0.0035	0.5596	0.0021
2024	6	0.6628	0.0040	0.6175	0.0023
2024	7	0.8195	0.0166	0.6965	0.0060
2024	8	0.9733	0.0439	0.8252	0.0138
2024	9	0.9850	0.0469	1.0162	0.0182
2024	10	1.1167	0.0401	1.0163	0.0227
2024	11	1.3455	0.0774	1.0774	0.0227
2024	12	1.2878	0.1082	1.2464	0.0455
2024	13	0.7485	0.7800	1.2878	0.1082
2024	14	1.1658	0.6356	1.2878	0.1082
2024	15+	0.9705	0.9011	Missing	Missing
2025	3	0.4354	0.0217	0.3143	0.0115
2025	4	0.5753	0.0102	0.5559	0.0085
2025	5	0.6356	0.0063	0.6214	0.0066
2025	6	0.6891	0.0050	0.6929	0.0027
2025	7	0.7913	0.0086	0.7295	0.0029
2025	8	0.8824	0.0175	0.8555	0.0073
2025	9	1.0084	0.0296	0.9657	0.0115
2025	10	1.0939	0.0411	1.1395	0.0292
2025	11	1.1054	0.0557	1.1943	0.0406
2025	12	1.4336	0.1344	1.0815	0.0241
2025	13	1.3513	0.6052	Missing	Missing
2025	14	1.1133	0.6965	Missing	Missing
2025	15+	0.8102	0.8284	1.2676	0.0950

Note. Fishery 2024–2025 weights and uncertainty in kilograms. The 2024 TMA results are a partial supported-only conversion; source-age-13 mass remains unallocated. Missing target ages have no converted abundance.

2024: partial bootstrap mean and fixed-key SD, conditional on the supported source ages; target age 15+ remains missing. 2025: converted mean and bootstrap SE.

Table G5 *BTS Mean Weights and Standard Errors After Joint Covariance Propagation*

Year	Age	Original mean	Original SE	TMA mean	TMA SE
2025	3	0.3655	0.0357	0.2255	0.0161
2025	4	0.4928	0.0352	0.5260	0.0166
2025	5	0.5951	0.0186	0.5871	0.0134
2025	6	0.6964	0.0114	0.6758	0.0098
2025	7	0.8329	0.0206	0.7340	0.0100
2025	8	0.9690	0.0331	0.8125	0.0125
2025	9	1.0288	0.0325	0.9026	0.0131
2025	10	1.1360	0.0383	1.0872	0.0197
2025	11	1.2675	0.0653	1.0659	0.0200
2025	12	1.4825	0.0943	1.1496	0.0257
2025	13	1.4562	0.1232	1.2577	0.0358
2025	14	1.7453	0.2714	1.3987	0.0558
2025	15+	1.7646	0.3427	1.5474	0.0834
2026	3	0.3850	0.0291	0.3971	0.0190
2026	4	0.5211	0.0184	0.5440	0.0155
2026	5	0.6188	0.0218	0.6400	0.0165
2026	6	0.7139	0.0237	0.7254	0.0109
2026	7	0.7551	0.0162	0.8076	0.0101
2026	8	0.8707	0.0144	0.8755	0.0101
2026	9	1.0041	0.0203	0.9476	0.0125
2026	10	1.1782	0.0322	1.0064	0.0132
2026	11	1.3129	0.0472	1.0415	0.0156
2026	12	1.4490	0.0762	1.2159	0.0209
2026	13	1.9177	0.1171	1.3719	0.0298
2026	14	1.6711	0.1496	1.5171	0.0648
2026	15+	1.8390	0.1928	1.3665	0.0336

Note. BTS mean weights and SEs in kilograms after propagating the joint model covariance.

Validation and source record

Table G6 *Validation of the Transformations*

Check	Result
Fishery independent recalculation	PASS
Pre-2025 fishery decimal preservation	Pass
Survey source SE reconstruction: largest relative difference	1.22e-15
Ratio derivative check: largest scaled difference	1.68e-11
Analytic versus numerical converted SE: largest relative difference	3.15e-11
Survey random-effect gradient: maximum	4.36e-12
Saved survey fixed Hessian positive definite	Yes
2024 specimen/age-field identity and partial arithmetic	Pass: 22 checks; positive source-age-13 support unresolved
Input adapter	Selected inputs used in completed working fits; historical partial-conversion support limits retained
wtAgeRE optimization	Screened-key standalone RE-WAA fit complete; earlier partial-conversion branch remains a separate diagnostic

Note. Validation of the transformations. Numerical checks retain the full-pair conversion basis; current fitted inputs use the selected 2024 tail treatment and screened 2026 key.

The source records retain matching abundance and weight estimates from the same saved survey fit, along with input/output hashes and fixed-key approximation rules. The empirical key is treated as fixed. Current fitting records and the historical conversion diagnostics are identified separately in the technical source record.

The historical 2024 partial-conversion diagnostics preserve their original inputs and supported-age limits. The current selected tail allocation, screened BTS key and completed fits have separate source and run identities. Earlier preparation states and the original verification wording remain archived in the technical source record.

This is an internal methods record. It uses a compact Quarto report and TAF-compatible source, transformation, output, and provenance stages. Assessment adoption is documented in the separate input and run records.

Historical partial-diagnostic fishery input: converted 2025, preserved 2024

Converted BTS input

Fishery approximation key

Conversion manifest

BTS uncertainty validation

2024 empirical key

2024 partial bootstrap summaries

2024 exact and pooled point candidates

2024 key support and unallocated source mass

2024 validation

Complete numbered conversion report · Editable analytical Quarto source · Subsection source manifest

Provisional ATS age compositions

Assessment preparation note supplied by Jim Ianelli. Provisional ATS age compositions were estimated by applying the BTS age–length key to the ATS abundance-at-length data. These estimates will be updated with ATS age-composition data in 2027.

References

Adamack, A.T., Antaya, K., Dhaliwal, S., Dwyer, K., Bourne, C.M., Forbes, G., 2025. Exploring the use of Fourier transform near infrared spectroscopy for aging Newfoundland and Labrador Atlantic cod (Gadus morhua). Fish. Res. 285, 107331. https://doi.org/10.1016/j.fishres.2025.107331

Basilone, G., Lo Cicero, G., Fortuna, M., Luviner, A., Ferreri, R., Aronica, S., Genovese, S., Giacalone, G., Fontana, I., Bonanno, A., 2024. Fourier-transform near-infrared spectroscopy first application to age determination in European fish species: the case of the Atlantic horse mackerel from the central Mediterranean Sea. Front. Mar. Sci. 11, 1429150. https://doi.org/10.3389/fmars.2024.1429150

Benson, I.M., Helser, T.E., Marchetti, G., Barnett, B.K., 2023. The future of fish age estimation: deep machine learning coupled with Fourier transform near-infrared spectroscopy of otoliths. Can. J. Fish. Aquat. Sci. cjfas-2023-0045. https://doi.org/10.1139/cjfas-2023-0045

Brownlee, J., 2019. Deep learning for computer vision: image classification, object detection, and face recognition in python.

Chamberlin D.W., Helser T.E., Brogan J.D., Benson I.M., Conner J., Chin A.T., Gburski C.K., Matta M.E., Pearce J.A., Stone K.R., TenBrink T.T., Arrington M.B. 2026. " A Framework to Investigate the Effects of Observation Error on Neural Network Predictions of Fish Age." Fish and Fisheries 27, no. 4: 974–986. https://doi.org/10.1111/faf.70094.

Chen, Y., Wang, Z., 2019. End-to-end quantitative analysis modeling of near-infrared spectroscopy based on convolutional neural network. J. Chemom. 33, e3122. https://doi.org/10.1002/cem.3122

Dick, E.J., Field, J.C., Grunloh, N., Rogers, T., 2025. The Status of Chilipepper Rockfish (Sebastes goodei) in U.S. Waters off California, Oregon, and Washington in 2025. Pac. Fish. Manag. Counc. Portland OR.

Géron, A., 2019. Hands-on machine learning with Scikit-Learn, Keras, and TensorFlow: concepts, tools, and techniques to build intelligent systems. O'Reilly Media, Inc., Sebastopol, CA.

Goldstein, E. D., Hsieh, B. C., Arrington, M. B., Helser, T. E. 2024. Quality assurance and quality control of Fourier transform near-infrared spectroscopy data for age prediction. In: Matta, M. E., and T. E. Helser (Eds). Proceedings of the Fourth Research Workshop on the Rapid Estimation of Fish Age Using Fourier Transform Near Infrared Spectroscopy. Alaska Fisheries Science Center, Resource Ecology and Fisheries Management Division, AFSC Processed Report 2024-01.

Ianelli, J.N., 2025. Eastern Bering Sea walleye pollock stock assessment. https://noaa-afsc.github.io/EBS_pollock/

Kastelle, C.R., and D.K. Kimura. 2006. Age validation of walleye pollock (Theragra chalcogramma) from the Gulf of Alaska using the disequilibrium of Pb-210 and Ra-226. ICES J. Mar. Sci. 63(8):1520-1529.

Kimura, D.K., and D.M. Anderl. 2005. Quality control of age data at the Alaska Fisheries Science Center. Mar. Freshwat. Res. 56(5):783-789.

Kimura, D.K., C.R. Kastelle, B.J. Goetz, C.M. Gburski, and A.V. Buslov. 2006. Corroborating the ages of walleye pollock (Theragra chalcogramma). Mar. Freshwat. Res. 57(3):323-332.

Kimura, D.K., D.M. Anderl, and B.J. Goetz. 2007. Seasonal marginal growth on otoliths of seven Alaska groundfish species support the existence of annual patterns. Alaska Fish. Res. Bull. 12(2):243-251.

Lai, H.L. 1985. Evaluation and validation of age determination for sablefish, pollock, Pacific cod and yellowfin sole; optimum sampling design using age-length key; and implication of aging variability in pollock. Doctoral dissertation, University of Washington, Seattle, WA.

Li, L., Jamieson, K., DeSalvo, G., Rostamizadeh, A., Talwalkar, A., 2018. Hyperband: A Novel Bandit-Based Approach to Hyperparameter Optimization. Hyperband Nov. Bandit-Based Approach Hyperparameter Optim. 18, 1–52. https://doi.org/10.48550/arXiv.1603.06560

Matta, M.E., Helser, T.E. (Eds.), 2024. Proceedings of the Fourth Research Workshop on the Rapid Estimation of Fish Age Using Fourier Transform Near Infrared Spectroscopy. https://doi.org/10.25923/g30w-8268

Matta, Mary Elizabeth, and Daniel K. Kimura (editors). 2012. Age determination manual of the Alaska Fisheries Science Center Age and Growth Program. NOAA Professional Paper NMFS 13, 97 p.

Matta, M. E., Short, J. A., Brogan, J. D., and Chamberlin, D. W. 2026. Characteristics and reliability of groundfish age data produced by the Alaska Fisheries Science Center's Age and Growth Program: reflecting on 50 years of age determination. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-AFSC-520, 222 p.

Murray, I., Williams, P., 1987. Chemical principles of near-infrared technology, in: Williams, P., Norris, K. (Eds.), Near Infrared Technology in the Agricultural and Food Industries. American Association of Cereal Chemists, St. Paul, MN, pp. 29–31.

Passerotti, M.S., Reichert, M.J.M., Robertory, B.A., Marsh, Z., Stefik, M., Quattro, J.M., 2022. Physicochemical mechanisms of FT-NIRS age prediction in fish otoliths. Mar. Freshw. Res. 73, 846–865. https://doi.org/10.1071/MF21341

Pomerantsev, A. L., and O. Y. Rodionova. 2014. Concept and role of extreme objects in PCA/SIMCA. J. Chemom. 28(5):429-438. https://doi.org/10.1002/cem.2506

Rodionova, O. Y., and A. L. Pomerantsev. 2020. Detection of outliers in projection-based modeling. Anal. Chem. 92:2656-2664. https://doi.org/10.1021/acs.analchem.9b04611

Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., Salakhutdinov, R., 2014. Dropout: A Simple Way to Prevent Neural Networks from Overfitting. J. Mach. Learn. Res. 15, 1929–1958.

Wedding, B.B., Forrest, A.J., Wright, C., Grauf, S., Exley, P., Poole, S.E., 2014. A novel method for the age estimation of Saddletail snapper (Lutjanus malabaricus) using Fourier Transform-near infrared (FT-NIR) spectroscopy. Mar. Freshw. Res. 65, 894. https://doi.org/10.1071/MF13244

Appendix H
BTS Spatiotemporal Biomass Index and Abundance at Age, Including the Northern Bering Sea

This appendix describes the spatiotemporal modeling approach used to estimate the bottom-trawl survey (BTS) biomass index and abundance at age for the combined eastern and northern Bering Sea (EBS and NBS). This approach was adopted in the 2024 assessment for both abundance at age and survey indices and follows the approach used by the AFSC Groundfish Assessment Program (GAP) in previous assessments. The sections below describe its application to the 2026 survey data, including model specifications and uncertainty estimates.

Scope and relationship to the earlier VAST appendix

The earlier VAST appendix describes a different software generation and mixed 1982–2023/2024 coverage. It is retained as a historical source. The description uses the settings recorded in the saved production fit objects. The saved production fit objects identify **sdmTMB 1.0.0** for the biomass index and **tinyVAST 1.6.3** for abundance at age. Model-based integration across EBS and NBS extends the approach used to address pollock distribution shifts (O'Leary et al. 2020). The sdmTMB framework is described by Anderson et al. (2022).

Both output series contain **44 years, 1982–2019 and 2021–2026**. The survey was absent in 2020. The index fit retains 2020 as an additional latent time step representing a model prediction. The age fit retains annual time levels across 1982–2026, while the abundance and composition data files omit 2020. NBS sampling coverage varies among years within the prediction domain.

Biomass index

The pollock index code reads density-dependence-corrected station biomass density from the VAST-named input file. The file name is historical; the saved model class is sdmTMB. Input density is converted from kg/ha to kg/km² by multiplying by 100. The saved fit uses a Poisson-link delta-gamma observation model with separate year effects and spatial and spatiotemporal fields. Both components have spatial fields and annual AR(1) spatiotemporal structure. The retained call enables geometric anisotropy and includes a spatially varying coefficient for the environmental covariate.

The companion pollock code defines that covariate as standardized cold-pool area, the area of bottom water at or below 2°C (Rohan et al. 2022). The saved object contains its numerical covariate and the formula; the complete annual cold-pool release manifest remains unavailable. The exact environmental release and production script revision require confirmation.

Predicted density is integrated over a fixed regional grid. For grid-cell area ($A_{r,j}$) in km² and predicted biomass density ($d_{t,j}$) in kg/km², the regional biomass index in tons is

$$\hat{I}_{r,t} = \frac{1}{1000} \sum_{j \in r} A_{r,j} \hat{d}_{t,j}. \quad (20)$$

The biomass-index estimates include separate EBS, NBS and combined **Both** products. The combined 2026 point estimate is **9,034,017.809 t**; the separate EBS and NBS point estimates sum to that value within numerical precision. These survey estimates remain subject to assessment input selection and acceptance.

Abundance at age

The saved tinyVAST fit uses numerical density at station and age, with age-by-year fixed effects in both delta components. All 15 labelled age classes have age-specific spatial variance terms and a common estimated lag-1 temporal coefficient in the spatiotemporal specification. The saved control has anisotropy disabled. This model uses age-by-year effects with the environmental terms excluded.

The companion pollock code multiplies CPUE_num by 100, converting fish/ha to fish/km². Its comment describing this as a conversion of kg is inconsistent with the numerical input; the calculation is an abundance calculation. Numerical density is integrated over the combined EBS–NBS grid and scaled to billions of fish:

$$\widehat{N}_{r,t,a}[billion fish]=\frac{1}{10^9}\sum_{j\in r}A_{r,j}\hat{n}_{t,a,j},\hat{p}_{r,t,a}=\frac{\widehat{N}_{r,t,a}}{\sum_b \widehat{N}_{r,t,b}}.\quad(21)$$

The abundance-at-age matrix contains 15 rows and 44 year columns. Its row order agrees with the companion age_1 through age_15 composition columns: normalizing every matrix column reproduces the composition file to a maximum absolute difference of (8.4^{-16}). The 2026 combined total is **16.064690 billion fish**. The companion composition file supplies the age labels for the matrix and must remain attached to it. Displayed ages are 1–14 and 15+; the terminal category represents fish aged 15 and older. The original source column names are retained in the data. The age export supplies only the combined EBS–NBS estimates (terminal-age interpretation).

These expanded compositions use the age assignments and age–length-key preparation in their station-level inputs. Age assignment and FT-NIRS-to-traditional-age conversion precede the tinyVAST expansion. The exact release, assignment rules for 2025–2026, and any age-age correction used in this production input remain part of the source reconciliation.

Verified settings and prediction-code evidence

The retained index fit uses 250 interior mesh knots, anisotropy and a spatially varying environmental coefficient; the age fit uses 50 interior knots with anisotropy disabled and a separate spatial variance structure by age (Table H1). Prediction-grid and integration settings are identified separately as companion-code evidence. The local code checkout is 2a9e6f26062d34ee0375f8eac9462e51273e65a3. Its default species selections are Pacific cod for the index and yellowfin sole for age compositions. The saved pollock fit objects supply the production evidence for this appendix; the generic default selections remain code context.

Table H1 *Delivered 2026 Pollock Fit and Companion Prediction Settings*

Setting	Biomass-index fit	Abundance-at-age fit
Platform and retained version	sdmTMB 1.0.0	tinyVAST 1.6.3
Response	Biomass density	Numerical density by age
Observation family	Poisson-link delta-gamma	Poisson-link delta-gamma
Fixed effects	Year in both components	Age × year in both components
Spatial mesh	250 interior knots; 307 total vertices	50 interior knots; 106 total vertices
Spatiotemporal structure	AR(1) in both components	Shared lag-1 coefficient; age-specific variance structure
Environmental coefficient	Spatially varying coefficient for env	None retained
Anisotropy	Enabled in saved call	Disabled in saved control
Observation years	1982–2019, 2021–2026	1982–2019, 2021–2026
Saved input rows	17,236	265,350 station–age rows
Reported output regions	EBS, NBS, Both	Both
Prediction grid in companion code	5 nmi; 1,578 EBS + 672 NBS cells	Same combined 2,250-cell grid
Integration bias/SE settings in companion code	get_index: bias_correct=TRUE	apply.epsilon=TRUE; bias.correct=FALSE; getsd=FALSE

Note. Retained settings in the 2026 pollock fit objects, with prediction-grid and integration settings identified separately as companion-code evidence. The complete production prediction record remains unavailable.

The code grid has 492,989.921 km² in the EBS and 198,866.805 km² in the NBS, summing to 691,856.726 km². Those grid files are hashed in the source record. The complete production prediction grid and expansion invocation remain unavailable. This draft therefore distinguishes companion-code settings from verified saved-fit settings. The historical 2-nmi grid and 750-knot index describe the earlier VAST workflow.

Uncertainty, diagnostics and outstanding review

The index code requests bias-corrected integration. It also contains a 500-simulation covariance workflow. The assessment selects its covariance file and uncertainty treatment separately from those integration settings. The age

expansion code sets bias.correct=FALSE and getsd=FALSE despite apply.epsilon=TRUE. Its age output therefore excludes epsilon bias correction and a full estimated age-composition covariance. Assessment weighting and the selected design-based covariance remain separate decisions.

The saved index object reports a positive definite Hessian and maximum absolute retained gradient **0.0138148**. The saved age object reports optimizer return code 0, a positive definite Hessian and maximum retained gradient **0.000205084**. These recorded diagnostics require provider review for numerical acceptance. Both gradients exceed the workspace's (10^{-4}) assessment-model convergence guardrail; their suitability for these survey fits requires provider review. This appendix uses the saved survey fits by the provider.

The saved index data have **17,236 rows**, while the adjacent raw index export has **17,237**; the counts differ in 2026 (350 versus 351). The saved age fit has **265,350 rows**, versus **258,555** in the adjacent age-density export. Historical differences occur in 1982, 1985, 1988, 1991, 1994, 2001, 2005, 2006 and 2018; 2026 has 5,250 fitted rows versus 5,265 exported rows. These differences prevent a claim that the saved objects exactly reproduce the neighboring raw files. Their causes, the intended final input versions and the fit-to-export chain require confirmation from the survey analysts.

Before finalizing this appendix, confirm the production script revision, prediction grid, annual environmental release, full R/TMB/dependency versions, age-assignment lineage, treatment of the input mismatches, and the accepted covariance and diagnostic assessment. The latest linked general writeup and terms-of-reference table remain unavailable through the connected source. Their agreement with this draft requires author review. Historical VAST dependency versions retain their original dates.

Source and reproducibility record

The local provenance bundle retains saved-object metadata and checks, source file hashes, output consistency checks, and the read-only extraction scripts. It also preserves the original historical VAST text and historical PDF. The original fit objects and station-level inputs remain in their owning data directory. The report consumes saved metadata and aggregate outputs.

Current-method source code: index workflow and age-composition workflow. Author-documents awaiting access: general methods writeup and terms-of-reference settings.

Appendix I
Assessment Development and Supporting Record

The September 2026 assessment development document (Ianelli et al. 2026), published 3 September, presents the starting model comparisons and exploratory appendices using data through 2024. Corrected through-2024 results are archived with that document; the main assessment uses observations through 2026.

The Plan Team excerpt provides the seven recommendation summaries (Table A1). Its final report citation and any subsequent SSC direction require verification before release.

The source guidelines are *Alaska Groundfish Stock Assessment Guidelines*, July 2024, in a file named *Stock Assessment Guidelines Tier 3.docx*. Their publication procedures inform eventual release preparation. The development copy remains local under the user's release timing.

Early–late fishery sensitivity summaries

Model 26.1.2 evaluates sensitivity to separating the early and late fisheries. Model 26.0 supplies the assessment recommendations.

Model 26.1.2 Early-late fishery split comparison				
Quantity	As estimated or specified in 2024 for:		Model 26.1.2 sensitivity for:	
	2025	2026	2027	2028
Natural mortality (annual rate, ages 3+)	0.300	0.300	0.300	0.300
Tier and sub-tier	3a	3a	3a	3a
Projected total biomass (ages 3+, t)	8,526,000	8,075,000	11,392,000	12,652,000
Projected female spawning biomass (t)	3,118,000	3,342,000	3,354,000	3,716,000
B100% (t)	5,902,000	5,902,000	6,296,000	6,296,000
B40% (t)	2,361,000	2,361,000	2,518,000	2,518,000
B35% (t)	2,066,000	2,066,000	2,204,000	2,204,000
F_OFL (annual rate)	0.513	0.513	0.696	0.696
Maximum F_ABC (annual rate)	0.394	0.394	0.544	0.544
Recommended F_ABC (annual rate)	0.394	0.394	Not selected	Not selected
OFL (t)	2,957,000	2,496,000	2,692,000	2,853,000
Maximum ABC (t)	2,417,000	2,036,000	2,187,000	2,316,000
Recommended ABC (t)	2,417,000	2,036,000	Not selected	Not selected
Historical source: December 2024 SAFE, Executive Summary, Tier 3 version, p. 4; B40% is printed in section 7.2, p. 35.				
2027–2028 projections assume catch of 1,375,000 t in each year, mean 2023–2025 selectivity and fixed biological inputs. Total biomass covers ages 3+ at the start of the year; spawning biomass is female biomass at spawning. F_OFL and maximum F_ABC are the fully selected F35% and F40% rates. Biomass and catch are rounded to the nearest 1,000 t.				
Model 26.1.2 is a sensitivity analysis. Recruitment is fixed at the fitted 1978–2023 mean; these projections show point estimates. Model 26.0 supplies the assessment recommendation.				

Model 26.1.2 Early-late fishery split stock-status summary. The 2025–2026 columns report status for 2025 and 2026; historical determinations are from the 2024 assessment.				
Status	As determined in 2024 for:		As determined this year for:	
	2023	2024	2025	2026
Overfishing	No	n/a	No	n/a
Overfished	n/a	No	n/a	No
Approaching overfished	n/a	No	n/a	No
Historical determinations are from the December 2024 EBS pollock SAFE, Executive Summary, p. 4.				
The 2025–2026 entries are assessment conclusions, separate from final NMFS status determinations and SSC/Council review. Overfishing for the most recent complete year compares the updated 2025 catch of 1,326,215.61 t with the specified 2025 Bering Sea subarea OFL of 2,957,000 t. The 2026 overfished determination uses each model's estimated 2026 female spawning biomass; approaching status uses its Scenario 7 projections through 2028 (Section 9). Model 26.1.2 is the Early-late fishery split comparison. n/a denotes a year/status combination outside this reporting layout.				

Assessment archive record

The assessment archive links model source and executable versions, prepared inputs, run controls, saved fits, diagnostics, retrospectives, projections and report products. Source manifests document provenance and access restrictions. Scientific acceptance is recorded separately from numerical diagnostics and report preparation.

Annual data-update register

The annual register identifies the input streams, observation years, revisions and weight-at-age treatments used in the assessment (Table I1).

Table I1 *Annual Data Additions, Revisions and Weight-at-Age Assumptions*

Annual data and weight-at-age register		
Data source	Year	Update or weight-at-age treatment
Fishery catch	2024	Historical catch revision check
Fishery catch	2025	Observed annual update
Fishery catch	2026	Year-to-date plus annual completion forecast
BTS index	2025	Observed annual update
BTS index	2026	Observed annual update
ATS index	2026	Observed annual update
AVO index	2025	Observed annual update
AVO index	2026	Observed annual update
BTS ages	2025	Observed annual update
BTS ages	2026	Observed annual update
ATS ages	2026	Observed annual update
Fishery ages	2024	Observed annual update
Fishery ages	2025	Observed annual update
Fishery weight at age	2024	Observed inputs plus forecast 2026
Fishery weight at age	2025	Observed inputs plus forecast 2026
Fishery weight at age	2026	Forecast schedule
Spawning weight at age	2024	A-season schedule correction
Spawning weight at age	2025	A-season schedule correction
Spawning weight at age	2026	A-season schedule correction
BTS weight at age	2025	Survey product with recorded fills
BTS weight at age	2026	Survey product with recorded fills
ATS weight at age	2025	Assumed carry of 2024 WAA
ATS weight at age	2026	Carry or provisional survey product

Note. Annual data additions, revisions and weight-at-age assumptions relative to the 2024 assessment. Listed updates are incorporated in the Model 23.x and 26.x fits. Cumulative input-update fits retain their stated stepwise history.

Appendix J
Contents

Table of Contents

List of Tables

List of Figures