Appendix 3K: Results of the Spatially Explicit Stock Assessment Models for Alaska Sablefish with Data Through 2026

Matthew LH. Cheng, Daniel R. Goethel

Introduction

Alaska sablefish (*Anoplopoma fimbria*) are currently managed as a single panmictic stock across the Gulf of Alaska (GOA), Bering Sea (BS), and Aleutian Islands (AI), informed by an age-and sex-structured statistical catch-at-age model. While the operational assessment assumes that the stock is spatially homogenous, research over the past several decades suggest that spatial structure likely exists to some degree. For instance, individuals have been shown to demonstrate age-specific (ontogenetic) movement patterns, wherein younger individuals reside in the BS and AI, and move towards the GOA as they mature (Cheng et al., 2025; Heifetz and Fujioka, 1991). In recent years, large recruitment events of age-2 individuals have also generally occurred in western regions (BS and AI). In general, these events reflect greater overall recruitment to the population, rather than a redistribution of recruits among regions, especially considering the BS historically serving as a stronghold for recruits. Moreover, fishery effort is primarily concentrated in the GOA regions, likely resulting in differential harvest impacts across space. Taken together, these patterns suggest that spatial heterogeneity in recruitment, movement, and harvest dynamics likely play a role in shaping Alaska sablefish population dynamics.

In the following, we present a spatially-explicit stock assessment model for Alaska sablefish, which attempts to explicitly model spatial heterogeneity in population dynamics. Spatially-explicit stock assessment models generally resemble single-region stock assessment models, but data are typically disaggregated across spatial regions, and movement dynamics are often incorporated into population processes. However, it is important to note that spatial models often involve an increase in the number of parameters estimated, coupled with a decrease in the amount of data informing parameter estimates through regional partitioning of data. Thus, spatial stock assessment models are often subject to greater tensions between model complexity and data availability, raising the practical question of whether the gains in structural realism directly translate to improved management performance relative to simpler non-spatial models. Nonetheless, several benefits can be envisioned from the implementation of a spatial model. Firstly, spatial models can provide a more mechanistic understanding of regional population dynamics and allow monitoring of localized depletion, which is not feasible under a non-spatial stock assessment model. Second, spatial models can serve as conditioning frameworks for operating models used in management strategy evaluations (MSEs). Lastly, spatial models potentially provide a means to address spatial changes in survey design, allowing the accommodation of missing data, and circumventing the need to interpolate missing data, as would be the case in a single-region model.

All model files for running the spatial assessment as well as model outputs and diagnostics can be found on the 2026 sablefish SAFE GitHub repo (here).

- Model *26.3_Spat_5Reg*: here
- Model *26.4_Spat_3Reg*: here

Methods

General Model Structure

Spatial stock assessment models were developed using the SPoRC (Stochastic Population over Regional Components) modelling framework (Cheng et al., 2026). SPoRC is a generalized statistical catch-at-age model written in RTMB (Kristensen et al., 2016) with the ability to support multi-region, multi-fleet, multi-sex configurations, while also integrating tagging data. Two configurations of spatial models were developed: a three-region model and a five-region model. The three-region model partitions the population across the BS, AI, and GOA, while the five-region model further disaggregates the GOA into the western GOA (WGOA), the central GOA (CGOA), and the eastern GOA (EGOA). Both configurations share the same population dynamics and model specifications, differing only in their spatial resolution and the number of estimated parameters. The spatial assessment models were generally based on the dynamics of the non-spatial models, the conceptual model shown in Figure 3D.1, and the parameterization of pervious work (Cheng et al., 2025). Specifically, spatial models represented 30 ages (2 – 31+), sex-specific dynamics (females and males), 67 years (1960 – 2026), two fishery fleets (longline and trawl), and two survey fleets (domestic and Japanese longline surveys). Within each annual time step, the population was subject to the following processes in sequence: recruitment of age-2 individuals, movement among regions, followed by mortality from fishing and natural mortality. Further details on model dynamics, observation equations, and likelihoods are described in https://chengmatt.github.io/SPoRC/articles/c_model_equations.html.

Recruitment

Recruitment of age-2 individuals in both spatial models were assumed to follow a mean recruitment relationship, where global mean recruitment is apportioned across spatial regions. Lognormal recruitment deviations were specified to vary across space and time, with the standard deviation of recruitment deviations fixed at 1.2 across all regions, following the convention of the non-spatial models. Similar to many spatial models, we assume that recruits are unable to move, in order to mitigate potential confounding with movement and recruitment dynamics.

Natural Mortality

Natural mortality is sex-specific and assumed to be constant across sexes and ages, consistent with the non-spatial models presented. Both values are fixed according to those estimated in the fleets as areas model (*26.2b_FAA_3Flt_LLS_Only*).

Growth and Maturity

Weight-at-age and maturity-at-age are treated as fixed quantities and are assumed to be constant across space, where values are taken from the non-spatial models. Weight-at-age is assumed to be sex-specific and follows a von Bertalanffy growth function (females asymptote at a larger size than males), while maturity-at-age is only modelled for females and is represented as a logistic function.

Movement

Movement among regions was modelled as a discrete time Markovian process, assuming constant rates across time, but age-specific movement. Age-specific movement was represented as three age-blocks representing young individuals (ages 2 – 7), intermediate aged individuals (8 – 15),

and old individuals (16 – 31) to capture ontogenetic shifts in spatial distribution (i.e., movement from western to eastern regions as individuals matured).

Selectivity and Catchability

Across all fishery and survey fleets, selectivity was parameterized as sex-specific logistic functions. The longline fishery fleet estimated time-varying selectivity as two time-blocks to capture a change in fleet structure (i.e., shifting from hook-and-line gear to pot gear), while the trawl fishery fleet estimated time-invariant selectivity. Both survey fleets similarly estimated time-invariant selectivity. Selectivity was assumed to be constant across regions. Catchability for both survey fleets were freely estimated and assumed to remain constant across space and time.

Tag Reporting

Tag reporting rates were also estimated to characterize tag recapture rates. Tag reporting was assumed to be constant across regions, but was allowed to vary across time as time blocks. In particular, three time-blocks were estimated, coinciding with large changes in fishery dynamics (inception of an individual fishing quota fishery and a change in fixed-gear fleet structure).

Observation Models and Likelihoods

Catch

Region-specific expected removals (weight) by the longline and trawl fishery are predicted using Baranov's catch equation, and are fit to assuming a lognormal likelihood with negligible error (standard deviation of catch is assumed to be fixed at 0.05).

Survey Indices

Region-specific survey indices (numbers) are fit to for both the domestic and Japanese longline survey fleets assuming a lognormal likelihood and design-based estimates of uncertainty. Unlike the single-region models, regional indices are fit directly without the need for interpolation or carryover values, and is able to adequately accommodate missing data.

Compositions

Both age and length-compositions are fit to assuming a Multinomial likelihood, where input sample sizes are generally based on *a priori* knowledge of sampling practices (i.e., regions and fleets with higher sampling coverage have larger sample sizes). Sex-specific compositions are available for all fishery and survey fleets and are fit to jointly (i.e., considering sex-specific compositions as a single vector; Cheng et al., 2025). Unlike the single-region counterparts, no data-weighting is performed to further adjust sample sizes, given complexities with respect to data-weighting in spatial models. Data-weighting assumptions for composition data will be explored in further iterations of the model.

Tagging Data

Since the inception of the longline survey, approximately 400,000 individuals have been tagged, with about 35,000 individuals recaptured. Thus, tagging data are expected to be highly informative of movement dynamics. Expected tag recaptures are predicted using a Brownie tag attrition submodel, and are fit using a multinomial release-conditioned likelihood, where the probability of recapture in each region is derived from the model's predicted movement and survival processes.

Moreover, the maximum tag liberty was restricted to 15 years (i.e., tags were not tracked or fit to beyond 15 years at liberty) to reduce computation time. To prevent tagging data from having due influence on overall model estimation, multinomial weights are scaled by a factor of 0.25. Similar to composition data, data-weighting assumptions for tagging data will continue to be explored in the future.

Model Estimation and Diagnostics

All parameters were estimated using penalized maximum likelihood using RTMB, where the objective function was the sum of negative log-likelihoods for catch data, survey indices, compositions, tagging, penalties (i.e., recruitment), and priors placed on parameters. The three-region model estimates a total of 732 parameters, while the five-region model estimates a total of 1223 parameters. For both models, the largest share of parameters primarily results from fishing mortality deviations (~50%), followed by recruitment deviations (~27%), initial age deviations (~12%), and movement parameters (~5%).

Several diagnostic tools were used to assess model adequacy, stability, and performance. In particular, all models were inspected to ensure that maximum gradient components for all parameters were < 1e-4, as well as a positive definite Hessian matrix. Model fits to catch data, survey indices, both age- and length-compositions, and tagging data were also evaluated (full diagnostics can be found on the GitHub repo). Lastly, retrospective analyses (7-year peels) were conducted to understand model stability under sequential data removal and to evaluate retrospective consistency in recruitment and spawning stock biomass (*SSB*).

Results

Both spatial models converged with maximum gradient components for all parameters < 1e-4, along with an invertible Hessian matrix. In general, most data sources were adequately fit and did not demonstrate systematic misfits. For instance, both spatial models provided good fits to regional catch data (Fig. 3D.2 and 3D.3). Likewise, abundance indices from both survey fleets were generally fit well, although indices in the most recent years for the GOA regions appeared to be overpredicted (Fig. 3D.4 and 3D.5), potentially reflecting the need to model time-varying movement dynamics. With respect to composition data, both models appeared to capture general patterns, wherein western regions often had higher proportions of younger fish, and eastern regions demonstrated higher proportions of older fish. While fits to composition data were generally adequate, there were some complex patterns that neither spatial model was able to reconcile. In particular, exceptionally large plus groups as observed by the longline fishery in the AI region were underpredicted in some years (see the GitHub repo for composition data fits). Relative to other data sources, tagging data were moderately well fit, although recaptures in recent periods appeared more difficult to reconcile (Fig. 3D.6 and 3D.7). Similar to the fits described for survey abundance indices, this may either reflect the need to explicitly represent time-varying movement dynamics, and/or changes in tag reporting dynamics.

Retrospective analyses for both three-region and five-region models generally demonstrated similar patterns, where all regions exhibited positive retrospective patterns in *SSB* that coincided with overestimated recruitment values (relative to the terminal year) from several years prior (Fig. 3D.8 and 3D.9). Interestingly, recruitment estimates in the AI from the three-region model

demonstrated some instability in certain retrospective peels (i.e., recruits in the Al are estimated at values close to 0), likely reflecting confounding (and revision of estimates) between movement and recruitment as age data were sequentially removed from the analysis (Fig. 3D.8). Thus, results from retrospective analyses highlight that spatially-explicit stock assessment models for Alaska sablefish will likely require careful annual tuning and evaluation of model structure, parameterization, and data weighting to ensure stable and biologically realistic dynamics through time.

Model estimates of recruitment and *SSB* were largely concordant between spatial models (Fig. 3D.10). Both the pattern and scale of both recruitment and *SSB* were similar, with large recruitment events primarily occurring in western regions (BS and AI) and sustained recruitment events in the eastern regions (WGOA, CGOA, and EGOA) (Fig. 3D.10 and 3D.11). Moreover, regions exhibiting high recruitment did not necessarily coincide with regions of high SSB, presumably due to the estimated ontogenetic movement dynamics (i.e., younger individuals primarily residing in western regions and moving eastward as they mature), as well as potential unmodelled larval dispersal processes (Cheng et al., 2025) (Fig. 3D.10 and 3D.11).

Overall, results from the spatial model highlight that spatial heterogeneity in population dynamics likely exists, driven in part by movement dynamics, differential spatial harvest, and underlying oceanographic processes. Despite the presence of such heterogeneity, aggregated time-series estimates from spatial models generally align with simpler alternatives in the case of Alaska sablefish (e.g., the single-region model; Cheng et al., 2025), suggesting that non-spatial models can provide similar advice for management purposes in the interim.

References

Cheng, M.LH., Goethel, D.R., Cunningham, C.J., Hulson, P.F., Ianelli, J.N., Omori, K.L., 2026. The SPoRC Stock Assessment Package: A Generalized Next-Generation Platform to Assess Spatial, Age and Sex-Structured Populations. Fish and Fisheries faf.70082. https://doi.org/10.1111/faf.70082

Cheng, M.LH., Hulson, P.-J.F., Goethel, D.R., Cunningham, C.J., 2025. A mathematical proof comparing the statistical properties between two common approaches for parameterizing sex-composition likelihoods in fishery stock assessments. Fisheries Research 281, 107231. https://doi.org/10.1016/j.fishres.2024.107231

Cheng, M.L.H., Marsh, C.A., Goethel, D.R., Hulson, P.F., Echave, K., Williams, B.C., Berger, A.M., Cunningham, C.J., 2025. Panmictic Panacea? Demonstrating Good Practices for Developing Spatial Stock Assessments Through Application to Alaska Sablefish (*Anoplopoma fimbria*). Fish and Fisheries faf.70002. https://doi.org/10.1111/faf.70002

Heifetz, J., Fujioka, J.T., 1991. Movement dynamics of tagged sablefish in the northeastern Pacific. Fisheries Research 11, 355–374. https://doi.org/10.1016/0165-7836(91)90009-5

Kristensen, K., Nielsen, A., Berg, C.W., Skaug, H., Bell, B., 2016. TMB: Automatic Differentiation and Laplace Approximation. J. Stat. Soft. 70. https://doi.org/10.18637/jss.v070.i05

Figures




Figure 3D.1. Conceptual model of spatial sablefish dynamics. Black solid arrows represent hypotheses on larval dispersal, blue dashed arrows are hypotheses regarding the movement dynamics of younger individuals, and orange dashed arrows are hypotheses on general movement dynamics of individuals as they mature.




Figure 3D.2. Fits to regional catch data (kt) by fishery fleet for the three-region model. Fleet 1 represents the longline fishery while fleet 2 represents the trawl fishery. The redline denotes the model expectation and point ranges represents observed data and their associated 95% confidence intervals.




Figure 3D.3. Fits to regional catch data (kt) by fishery fleet for the five-region model. Fleet 1 represents the longline fishery while fleet 2 represents the trawl fishery. The redline denotes the model expectation and point ranges represents observed data and their associated 95% confidence intervals.




Figure 3D.4. Fits to regional survey indices (numbers) by survey fleet for the three-region model. The redline denotes the model expectation and point ranges represents observed data and their associated 95% confidence intervals.




Figure 3D.5. Fits to regional survey indices (numbers) by survey fleet for the three-region model. The redline denotes the model expectation and point ranges represents observed data and their associated 95% confidence intervals.




Figure 3D.6. Fits to tag recaptures across regions for the three-region model. Colored lines represent the model expectation while grey points denote observed recaptures.




Figure 3D.7. Fits to tag recaptures across regions for the five-region model. Colored lines represent the model expectation while grey points denote observed recaptures.




Figure 3D.8. Retrospective analysis from the three-region model. Column panels represent time series of recruitment and spawning stock biomass (*SSB*) as data are sequentially removed. Row panels denote the modeled regions. Mohn's rho values are shown in the upper left of each panel.




Figure 3D.9. Retrospective analysis from the five-region model. Column panels represent time series of recruitment and spawning stock biomass (*SSB*) as data are sequentially removed. Row panels denote the modeled regions. Mohn's rho values are shown in the upper left of each panel.




Figure 3D.10. Comparison of time-series of recruitment (age-2; millions of individuals) and spawning stock biomass (*SSB*; kt) between the three-region and five-region spatial models. Time-series estimates for the Gulf of Alaska from the five-region model are summed up across the western, central, and eastern Gulf of Alaska for comparison with the three-region model.




Figure 3D.11. Time-series of recruitment (age-2; millions of individuals) and spawning stock biomass (*SSB*; kt) for the five-region spatial model.