

Math 3B: Lecture 4

Noah White

September 30, 2016

Problem set 2 and homework

- Problem set 2 is online now.

Problem set 2 and homework

- Problem set 2 is online now.
- Homework 1 is due next Friday 7 October at the start of the lecture

Problem set 2 and homework

- Problem set 2 is online now.
- Homework 1 is due next Friday 7 October at the start of the lecture
- The homework will be problems 2(a), 3, and 8.

Last time

Last time, we spoke about

- Slanted asymptotes

Last time

Last time, we spoke about

- Slanted asymptotes
- What are maxima and minima

Last time

Last time, we spoke about

- Slanted asymptotes
- What are maxima and minima
- Critical points

Last time

Last time, we spoke about

- Slanted asymptotes
- What are maxima and minima
- Critical points
- Finding maxima and minima

Last time

Last time, we spoke about

- Slanted asymptotes
- What are maxima and minima
- Critical points
- Finding maxima and minima
- First derivative test

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)


- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .



First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)


- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .



Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Note: If $f''(a) = 0$ then we cannot conclude anything! E.g x^3 or x^4 .

Example 1

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

Example 1

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

Example 1

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

a	$f''(a)$
0	6
2	-6

Example 1

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

a	$f''(a)$
0	6
2	-6

- Local minimum at $x = 0$

Example 1

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

a	$f''(a)$
0	6
2	-6

- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example 1

- Local minimum at $x = 0$

Example 1


- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example 1

- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example 1

- Local minimum at $x = 0$
- Local maximum at $x = 2$



Example 2

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

Example 2

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

Example 2

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

The critical points are

a	$f''(a)$
0	-16

Example 2

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

The critical points are

a	$f''(a)$
0	-16

- Local maximum at $x = 0$

Example 2


- Local and global maximum at $x = 0$

Example 2

- Local and global maximum at $x = 0$

Example 2

- Local and global maximum at $x = 0$



Example 3

$$f(x) = \begin{cases} -x & \text{if } x < -1 \\ 3x + 4 & \text{if } x \geq -1 \end{cases} \quad (\text{domain: } [-5, 2])$$

Example 3

$$f(x) = \begin{cases} -x & \text{if } x < -1 \\ 3x + 4 & \text{if } x \geq -1 \end{cases} \quad (\text{domain: } [-5, 2])$$

$$f'(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3 & \text{if } x > -1 \\ \text{undef.} & \text{if } x = -5, -1, 2 \end{cases}$$

The critical points are

a	$f(a)$
-5	5
-1	1
2	10

Example 3

$$f(x) = \begin{cases} -x & \text{if } x < -1 \\ 3x + 4 & \text{if } x \geq -1 \end{cases} \quad (\text{domain: } [-5, 2])$$

$$f'(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3 & \text{if } x > -1 \\ \text{undef.} & \text{if } x = -5, -1, 2 \end{cases}$$

The critical points are

a	$f(a)$
-5	5
-1	1
2	10

- Local minimum at $x = -1$

Example 3

$$f(x) = \begin{cases} -x & \text{if } x < -1 \\ 3x + 4 & \text{if } x \geq -1 \end{cases} \quad (\text{domain: } [-5, 2])$$

$$f'(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3 & \text{if } x > -1 \\ \text{undef.} & \text{if } x = -5, -1, 2 \end{cases}$$

The critical points are

a	$f(a)$
-5	5
-1	1
2	10

- Local minimum at $x = -1$
- Local maximums at $x = -5, 2$

Example 3

- Local minimum at $x = -1$

Example 3


- Local minimum at $x = -1$
- Local maximums at $x = -5, 2$

Example 3

- Local minimum at $x = -1$
- Local maximums at $x = -5, 2$

Example 3

- Local minimum at $x = -1$
- Local maximums at $x = -5, 2$



Example 4

Question

Suppose you have M feet of fencing, what is the largest rectangular area you can enclose?

Example 4

Question

Suppose you have M feet of fencing, what is the largest rectangular area you can enclose?

Solution

Suppose we create a rectangle with length x feet and width y feet. They must satisfy

$$2x + 2y = M \quad \text{i.e.} \quad y = \frac{1}{2}(M - 2x)$$

and the area is given by $A(x, y) = xy$.

Example 4

Question

Suppose you have M feet of fencing, what is the largest rectangular area you can enclose?

Solution

Suppose we create a rectangle with length x feet and width y feet. They must satisfy

$$2x + 2y = M \quad \text{i.e.} \quad y = \frac{1}{2}(M - 2x)$$

and the area is given by $A(x, y) = xy$.

Example 4

Question

Suppose you have M feet of fencing, what is the largest rectangular area you can enclose?

Solution

Suppose we create a rectangle with length x feet and width y feet. They must satisfy

$$2x + 2y = M \quad \text{i.e.} \quad y = \frac{1}{2}(M - 2x)$$

and the area is given by $A(x, y) = xy$. So

$$A(x) = x \frac{1}{2}(M - 2x) = \frac{M}{2}x - x^2$$

Example 4

Question

Suppose you have M feet of fencing, what is the largest rectangular area you can enclose?

Solution

Suppose we create a rectangle with length x feet and width y feet. They must satisfy

$$2x + 2y = M \quad \text{i.e.} \quad y = \frac{1}{2}(M - 2x)$$

and the area is given by $A(x, y) = xy$. So

$$A(x) = x \frac{1}{2}(M - 2x) = \frac{M}{2}x - x^2$$

We would like to find the value of x which maximises this function!

Example 4

The derivative:

$$A'(x) = \frac{M}{2} - 2x$$

Example 4

The derivative:

$$A'(x) = \frac{M}{2} - 2x$$

A critical point at $x = \frac{M}{4}$.

Example 4

The derivative:

$$A'(x) = \frac{M}{2} - 2x$$

A critical point at $x = \frac{M}{4}$.

The second derivative $A''(x) = -2$ is always negative so this must be a maximum! Thus the dimensions for the rectangle with the largest area are

$$x = y = \frac{M}{4}$$

Catalina island

Question

You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/*km* on land, and you can charter a boat for \$100/*km*. You are very thrifty and want to save money, how far from the LA River should you come ashore?

Catalina island


Question

You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/*km* on land, and you can charter a boat for \$100/*km*. You are very thrifty and want to save money, how far from the LA River should you come ashore?

Catalina island

Question


You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/ km on land, and you can charter a boat for \$100/ km . You are very thrifty and want to save money, how far from the LA River should you come ashore?



Catalina island

Question


You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/ km on land, and you can charter a boat for \$100/ km . You are very thrifty and want to save money, how far from the LA River should you come ashore?



Catalina island

Question


You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/ km on land, and you can charter a boat for \$100/ km . You are very thrifty and want to save money, how far from the LA River should you come ashore?



Catalina island

Question

You want to get from Catalina Island to Long Beach/LA River. An UberX costs \$2/*km* on land, and you can charter a boat for \$100/*km*. You are very thrifty and want to save money, how far from the LA River should you come ashore?



Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Setting the derivative equal to zero we obtain

$$2 = -\frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Setting the derivative equal to zero we obtain

$$2 = -\frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

$$2\sqrt{x^2 - 46x + 2378} = 460 - 20x$$

Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Setting the derivative equal to zero we obtain

$$2 = -\frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

$$2\sqrt{x^2 - 46x + 2378} = 460 - 20x$$

$$4x^2 - 184x + 9512 = 211600 - 18400x + 400x^2$$

Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Setting the derivative equal to zero we obtain

$$2 = -\frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

$$2\sqrt{x^2 - 46x + 2378} = 460 - 20x$$

$$4x^2 - 184x + 9512 = 211600 - 18400x + 400x^2$$

$$0 = 396x^2 - 18216x + 202088$$

Catalina island

$$C(x) = 2x + 100\sqrt{49^2 + (23 - x)^2}$$

$$C'(x) = 2 + \frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

Setting the derivative equal to zero we obtain

$$2 = -\frac{20(x - 23)}{\sqrt{x^2 - 46x + 2378}}$$

$$2\sqrt{x^2 - 46x + 2378} = 460 - 20x$$

$$4x^2 - 184x + 9512 = 211600 - 18400x + 400x^2$$

$$0 = 396x^2 - 18216x + 202088$$

So the solutions are

$$x = 23 - \frac{43}{3\sqrt{11}} \quad \text{and} \quad 23 + \frac{43}{3\sqrt{11}}$$