

Math 3B: Lecture 9

Noah White

October 16, 2018

Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

The **challenge** is to find all the functions $y = f(x)$ (or even just one) that satisfy a given equation.

Newton's second law (motivation)

The original differential equation!

$$F = ma$$

Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If $h(t)$ measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If $h(t)$ measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

The force due to gravity is roughly $-10m$ Newtons, so

$$-10m = mh''(t)$$

Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

Assumption

The rate of growth of a population is proportional to its current size

Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

Assumption

The rate of growth of a population is proportional to its current size

Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

Assumption

The rate of growth of a population is proportional to its current size

If $P(t)$ is the population at time t :

$$\frac{dP}{dt} = rP(t)$$

Some more examples of differential equations

$$\frac{dy}{dx} = f(x)$$

$$\frac{dy}{dx} = y(1 - y)$$

$$y'' = \sqrt{a^2 - (y')^2}$$

$$\frac{dy}{dt} = k(A - y)^2$$

Antiderivatives

We have been solving differential equations of the form

$$\frac{dy}{dx} = f(x).$$


Any antiderivative $y = F(x)$ of $f(x)$ is a solutions to this differential equation!

Slope fields

In some cases it is impossible to find the antiderivative (without special functions). E.g.

$$f(x) = e^{-x^2}$$

But we can still (approximately) graph the antiderivative! First we draw the slope field



Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the xy -plane.

Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the xy -plane.
2. At every point (x, y) what would the slope of $y = F(x)$ be if it passed through that point?

Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the xy -plane.
2. At every point (x, y) what would the slope of $y = F(x)$ be if it passed through that point?
3. Answer given by differential equation above, slope is $f(x)$

Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the xy -plane.
2. At every point (x, y) what would the slope of $y = F(x)$ be if it passed through that point?
3. Answer given by differential equation above, slope is $f(x)$
4. Draw a small arrow with slope $f(x)$ and the point (x, y)

Slope fields (how to draw)


How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the xy -plane.
2. At every point (x, y) what would the slope of $y = F(x)$ be if it passed through that point?
3. Answer given by differential equation above, slope is $f(x)$
4. Draw a small arrow with slope $f(x)$ and the point (x, y)
5. Do this for a grid of points on the xy -plane.


Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



Accumulated change

Often we encounter problems involving **accumulated change**.

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

These problems involve finding the area under some curve.

Example 1


If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

Example 1

If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

Solution

We model the car's speed using the function $s(t) = 30$. So we can see that the area under this curve



is the distance travelled (75 miles)

Example 2


If a car accelerates for 20 seconds at a rate of $2m/s^2$ and then decelerates for 30 seconds at a rate of $1m/s^2$, how far has it travelled?

Example 2

If a car accelerates for 20 seconds at a rate of 2m/s^2 and then decelerates for 30 seconds at a rate of 1m/s^2 , how far has it travelled?


Solution

The car's speed is given by $s(t) = 2t$ when $0 \leq t \leq 20$ and $s(t) = 60 - t$ when $20 \leq t \leq 50$. So the graph looks like




More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).

Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between $t = a$ and $t = b$?

Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between $t = a$ and $t = b$?
- Answer: area under $f(t)$ between a and b .

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

A first approach to calculating the area under a curve is to approximate using rectangles:

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

A first approach to calculating the area under a curve is to approximate using rectangles:

(Too hard to draw, lets look at an animation)

The definite integral

Defintion

The definite integral of a function $f(x)$ is defined to be

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$

where $\Delta x = \frac{b-a}{n}$.


The definite integral

Defintion

The definite integral of a function $f(x)$ is defined to be

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$


where $\Delta x = \frac{b-a}{n}$.



Properties of definite integrals

Zero area


$$\int_a^a f(x) \, dx = 0$$



Properties of definite integrals

Adding areas

$$\int_a^c f(x) \, dx = \int_a^b f(x) \, dx + \int_b^c f(x) \, dx$$



More properties of definite integrals

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

More properties of definite integrals

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

More properties of definite integrals

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

Linearity (scalars factor out)

$$\int_a^b \alpha f(x) \, dx = \alpha \int_a^b f(x) \, dx$$

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .
- every input x produces a number as an output.

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Why?

Well $F(x) = \int_a^x f(t) \, dt + C$ for some a and C . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Solution

An antiderivative of $x^2 - 4$ is $\frac{1}{3}x^3 - 4x$ so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Solution

An antiderivative of $\sin x$ is $-\cos x$ so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\frac{d}{dx} F(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \end{aligned}$$

Why is the FTC true?


$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt \end{aligned}$$


Why is the FTC true?

When h is very small $\int_x^{x+h} f(t) \approx$



Why is the FTC true?

When h is very small $\int_x^{x+h} f(t) \approx hf(x)$



Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$

Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} hf(x)\end{aligned}$$

Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x) \\&= f(x)\end{aligned}$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Why?

Well $F(x) = \int_a^x f(t) \, dt + C$ for some a and C . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

The indefinite integral

We also use the following notation for the general antiderivative:

Definition

$$\int f(x) \, dx := F(x) + C$$

The indefinite integral

We also use the following notation for the general antiderivative:

Definition

$$\int f(x) \, dx := F(x) + C$$

Example

$$\int \sin(x) - x \, dx = -\cos(x) - \frac{1}{2}x^2 + C$$

Example

Question

Find an antiderivative of $f(x) = |x|$?

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of a .

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) dt$$

is an antiderivative for any choice of a .

- Lets use $a = 0$.

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) dt$$

is an antiderivative for any choice of a .


- Lets use $a = 0$.
- How should we calculate $F(x)$?

Example

Use the definition!

$$F(x) = \int_0^x |t| \, dt$$

is the area under $|t|$!




$$F(x) = \frac{1}{2}x^2 \quad \text{if } x \geq 0$$

Example

If $x \leq 0$ then

$$F(x) = \int_0^x |t| \, dt = - \int_x^0 |t| \, dt$$

is the negative of the area under $|t|$!



$$F(x) = -\frac{1}{2}x^2 \quad \text{if } x \leq 0$$

Example

In summary

$$F(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } x \geq 0 \\ -\frac{1}{2}x^2 & \text{if } x \leq 0 \end{cases}$$

or

$$F(x) = \frac{1}{2}x|x|$$