

This week on the problem set you will get practice at calculating integrals using substitution and integration by parts.

\*Numbers in parentheses indicate the question has been taken from the textbook:

S. J. Schreiber, *Calculus for the Life Sciences*, Wiley,

and refer to the section and question number in the textbook.

**Homework:** The first homework will be due on Friday 12 October, at 8am, the *start* of the lecture. It will consist of questions:

5 and 8

1. (5.3) Express the limits as definite integrals of the form  $\int_0^1 f(x) dx$ .

- (a) (5.3.1)  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{i}{n^2}$   
 (b) (5.3.5)  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left(1 - \frac{i^2}{n^2}\right) \frac{1}{n}$   
 (c) (5.3.6)  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \sin\left(\frac{\pi i}{n} - \pi\right) \frac{\pi}{n}$

**Solution:** We know that if the integral ranges from  $x = 0$  to  $x = 1$  then  $\Delta x = \frac{1}{n}$  and  $x_i = \frac{i}{n}$ . Using the definition of the definite integral

$$\begin{aligned} \int_0^1 f(x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{i}{n}\right) \frac{1}{n}. \end{aligned}$$

So we need that  $f\left(\frac{k}{n}\right) \frac{1}{n} = \sin\left(\frac{\pi i}{n} - \pi\right) \frac{\pi}{n}$  i.e. that

$$f\left(\frac{k}{n}\right) = \sin\left(\frac{\pi i}{n} - \pi\right) \pi$$

which is obviously achieved if  $f(x) = \pi \sin(\pi x - \pi)$ . That means the definite integral corresponding to the Riemann sum is

$$\int_0^1 \pi \sin(\pi x - \pi) dx.$$

2. (5.3) Express the definite integrals as limits of Riemann sums.

- (a) (5.3.8)  $\int_{-1}^1 (x^2 - x) dx$

**Solution:** We know that if the integral ranges from  $x = -1$  to  $x = 1$  then  $\Delta x = \frac{2}{n}$  and  $x_i = \frac{2i}{n} - 1$ . Using the definition of the definite integral

$$\begin{aligned} \int_{-1}^1 f(x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{2i}{n} - 1\right) \frac{2}{n}. \end{aligned}$$

Thus, since  $f(x) = x^2 - x = x(x - 1)$  we get that

$$\int_{-1}^1 x^2 - x \, dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left( \frac{2i}{n} - 1 \right) \left( \frac{2i}{n} - 2 \right) \frac{2}{n}.$$

- (b) (5.3.9)  $\int_0^1 e^x \, dx$   
 (c) (5.3.11)  $\int_{-1}^1 |x| \, dx$
3. (5.5) Calculate the following integrals using substitution.
- (a) (5.5.12)  $\int \frac{x}{\sqrt{x^2+1}} \, dx$   
 (b) (5.5.14)  $\int \sin^3 t \cos t \, dt$   
 (c) (5.5.16)  $\int \frac{z^3}{\sqrt{z^4+12}} \, dz$   
 (d) (5.5.19)  $\int_1^2 \frac{e^{1/x}}{x^2} \, dx$   
 (e) (5.5.23)  $\int_1^2 x\sqrt{x-1} \, dx$   
 (f) (5.5.24)  $\int_0^2 (e^x - e^{-x})^2 dx$
4. (5.5-30) Suppose an environmental study indicates that the ozone level,  $L$ , in the air above a major metropolitan center is changing at a rate modeled by the function

$$L'(t) = \frac{0.24 - 0.03t}{\sqrt{36 + 16t - t^2}}$$

parts per million per hour (ppm/h)  $t$  hours after 7:00 A.M.

- (a) Express the ozone level  $L(t)$  as a function of  $t$  if  $L$  is 4 ppm at 7:00 A.M.

**Solution:** The function  $L(t)$  expressing the ozone level at time  $t$  will be an antiderivative of  $L'(t)$ . That is

$$L(t) = \int \frac{0.24 - 0.03t}{\sqrt{36 + 16t - t^2}} \, dt.$$

We will use the substitution  $u = 36 + 16t - t^2$ . Thus  $u' = 2(8 - t)$ . Note that  $0.24 - 0.03t = 0.03(8 - t)$ . Thus

$$\begin{aligned} L(t) &= \int \frac{0.03}{\sqrt{36 + 16t - t^2}} \cdot \frac{1}{2} 2(8 - t) \, dt \\ &= 0.03 \int \frac{1}{2\sqrt{u}} \, du \\ &= 0.03\sqrt{u} + C \\ &= 0.03\sqrt{36 + 16t - t^2} + C \end{aligned}$$

To find the constant  $C$  we simply solve the equation  $L(0) = 4$ , that is,

$$\begin{aligned} 0.03\sqrt{36 + 16 \cdot 0 - 0^2} + C &= 4 \\ 0.03\sqrt{36} + C &= \\ 0.18 + C &= \\ C &= 4 - 0.18 = 3.82. \end{aligned}$$

Thus

$$L(t) = 0.03\sqrt{36 + 16t - t^2} + 3.82.$$

- (b) Find the time between 7:00 A.M. and 7:00 P.M. when the highest level of ozone occurs. What is the highest level? (*Note: part b has been changed slightly from what is written in the textbook.*)

**Solution:** First we find the critical points by setting  $L'(t) = 0$ . This happens when  $t = 8$ , i.e. at 3pm. Using the first derivative test we know this is a maximum. Thus the highest level of ozone is


$$L(8) = 0.03\sqrt{26 + 16 \cdot 8 - 64} - 3.82 = 0.09\sqrt{10} - 3.82 = 4.10\text{ppm}$$

5. The circle  $x^2 + (y + 1)^2 = 4$  has area  $4\pi$ . What is the area of the portion of the circle lying above the  $x$  axis?

You may use the fact that

$$\int \sqrt{1 - t^2} dt = \frac{1}{2} \left( t\sqrt{1 - t^2} + \sin^{-1} t \right) + C.$$

**Solution:** We first draw a picture so that we can visualise the area we would like to find.



We want to find the shaded area. The circle is given by the equation  $x^2 + (y + 1)^2 = 4$ , which means the function that describes the top half semicircle is

$$y = \sqrt{4 - x^2} - 1$$

and the area is given by the integral

$$A = \int_a^b \sqrt{4 - x^2} - 1 \, dx.$$

Here  $a$  and  $b$  are the  $x$ -intercepts of the semicircle. We can find these by setting  $y = 0$  and solving for  $x$ :

$$\begin{aligned} 0 &= \sqrt{4 - x^2} - 1 \\ 1 &= \sqrt{4 - x^2} \\ 1 &= 4 - x^2 \\ x^2 &= 4 - 1 = 3 \\ x &= \pm\sqrt{3}. \end{aligned}$$

Thus  $a = -\sqrt{3}$  and  $b = \sqrt{3}$ . Thus

$$A = \int_{-\sqrt{3}}^{\sqrt{3}} \sqrt{4-x^2} - 1 \, dx$$

which we can separate,

$$= \int_{-\sqrt{3}}^{\sqrt{3}} \sqrt{4-x^2} \, dx - \int_{-\sqrt{3}}^{\sqrt{3}} dx$$

and factor out the 4,

$$= \int_{-\sqrt{3}}^{\sqrt{3}} 2\sqrt{1-\left(\frac{x}{2}\right)^2} \, dx - \int_{-\sqrt{3}}^{\sqrt{3}} dx.$$

Note that

$$\int_{-\sqrt{3}}^{\sqrt{3}} dx = 2\sqrt{3}. \quad (1)$$

We can solve the first part of  $A$  by using the substitution  $u = \frac{x}{2}$ , so  $u' = \frac{1}{2}$ . Note that when  $x = \pm\sqrt{3}$  then  $u = \pm\frac{\sqrt{3}}{2}$ . This means

$$2 \int_{-\sqrt{3}}^{\sqrt{3}} \sqrt{1-\left(\frac{x}{2}\right)^2} \, dx = 4 \int_{-\sqrt{3}/2}^{\sqrt{3}/2} \sqrt{1-u^2} \, dx$$

now we can apply the antiderivative given in the question,

$$= 4 \left[ \frac{1}{2} u \sqrt{1-u^2} + \frac{1}{2} \sin^{-1} u \right]_{-\sqrt{3}/2}^{\sqrt{3}/2}$$

noting that  $\sin^{-1}\left(\pm\frac{\sqrt{3}}{2}\right) = \pm\frac{\pi}{3}$ , we get

$$\begin{aligned} &= \sqrt{3} \cdot \frac{1}{2} + 2 \cdot \frac{\pi}{3} - \left(-\sqrt{3}\right) \cdot \frac{1}{2} - 2 \cdot \left(-\frac{\pi}{3}\right) \\ &= \sqrt{3} + \frac{4\pi}{3}. \end{aligned} \quad (2)$$

Since  $A = (2) - (1)$  we have  $A = \frac{4\pi}{3} - \sqrt{3}$ .

6. (5.6) Calculate the following integrals using integration by parts.

(a) (2)  $\int e^t \sin t \, dt$

(b) (6)  $\int x^2 \ln x \, dx$

(c) (9)  $\int \sin x \cos x \, dx$

(d) (14)  $\int_0^\pi x \sin x \, dx$

(e) (16)  $\int_1^e x^3 \ln x \, dx$


7. Use the fundamental theorem of calculus and the interpretation of the definite integral as an area to find a formula for the general antiderivative of the function  $f(x) = |x|$ .

**Solution:** The fundamental theorem of calculus says that for any constant  $a$ , the function

$$F(x) = \int_a^x |t| \, dt$$

will be an antiderivative of  $f(x) = |x|$ .

Let us choose  $a = 0$  for simplicity. To evaluate this integral we consider two cases. First when  $x \geq 0$ . In this case we can interpret the integral as an area.




This is a triangle with height  $x$  and base length  $x$  so the area is  $\frac{1}{2}x^2$ . I.e.  $F(x) = \frac{1}{2}x^2$  when  $x \geq 0$ . Now consider the case when  $x < 0$ . We cannot use the area interpretation of the integral

$$F(x) = \int_0^x |t| \, dt$$

since we are going from right to left, however we can use the property of definite integrals that says we can swap the limits at the expense of a minus sign:

$$F(x) = \int_0^x |t| \, dt = - \int_x^0 |t| \, dt.$$

The integral on the right is now one we can evaluate using the same area interpretation:



The area again is a triangle with area  $\frac{1}{2}x^2$ , thus  $F(x) = -\frac{1}{2}x^2$  when  $x < 0$ . Summarising, the general antiderivative is thus and shift by a constant of what we have found above:

$$F(x) = \begin{cases} \frac{1}{2}x^2 + C & \text{if } x \geq 0 \\ -\frac{1}{2}x^2 + C & \text{if } x < 0 \end{cases}$$

$$= \frac{1}{2}x|x|.$$

8. Use the fundamental theorem of calculus and the interpretation of the definite integral as an area to find a formula for the general antiderivative of the function  $f(x) = \frac{1}{x}$ .

**Solution:** The fundamental theorem of calculus says that for any constant  $a$ , the function


$$F(x) = \int_a^x \frac{1}{t} dt$$

will be an antiderivative of  $f(x) = \frac{1}{x}$ . We know what happens when  $x > 0$ , in this case  $F(x) = \ln x + C$ . So we concentrate on  $x < 0$ .

Let us choose  $a = -1$  for simplicity. We are interested in the area

$$F(x) = \int_{-1}^x \frac{1}{t} dt$$

This is the green shaded area below



By symmetry, this is exactly the negative of the red area! So

$$F(x) = \int_{-1}^x \frac{1}{t} dt = - \int_{-x}^1 \frac{1}{t} dt = [-\ln t]_{-x}^1 = \ln(-x).$$

So, summarising,

$$F(x) = \begin{cases} \ln x + C & \text{if } x > 0 \\ \ln -x + C & \text{if } x < 0 \end{cases}$$

$$= \ln |x|.$$