

# Math 3B: Lecture 18

Noah White

February 22, 2019

# Linear models

## Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants  $a$  and  $b$ .

# Linear models

## Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants  $a$  and  $b$ .

# Linear models

## Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants  $a$  and  $b$ . In other words, the right hand side is a linear function of  $y$ .

# Linear models

## Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants  $a$  and  $b$ . In other words, the right hand side is a linear function of  $y$ .

## Examples

$$\frac{dy}{dt} = ay, \quad \frac{dy}{dt} = -\lambda y.$$

# Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant

so

# Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

# Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so



# Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

# Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

# Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

## Note

**Something** could mean (for example)

- concentration of a drug in bloodstream

# Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

## Note

**Something** could mean (for example)

- concentration of a drug in bloodstream
- pollutant in water supply

## General solution

Using separation of variables, we can show that the general solution to

$$\frac{dy}{dt} = a - by$$

is

$$y(t) = \frac{a}{b} - Ce^{-bt}$$

where  $C$  is an arbitrary constant.

## Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration  $y(t)$  at time  $t$ .

Solution

## Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration  $y(t)$  at time  $t$ .

### Solution

- Ignoring infusion, every 2 hours the amount of drug halves.

## Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration  $y(t)$  at time  $t$ .

### Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with  $M$  mg, after  $t$  hours there will be

$$M \left( \frac{1}{2} \right)^{t/2} = M e^{-0.5t \ln(2)} \text{ mg left}$$



## Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration  $y(t)$  at time  $t$ .

### Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with  $M$  mg, after  $t$  hours there will be

$$M \left( \frac{1}{2} \right)^{t/2} = M e^{-0.5t \ln(2)} \text{ mg left}$$

- Thus the rate at which the drug is leaving (at time  $t$ ) is given by

$$0.5 \ln(2) M e^{-0.5t \ln(2)} = 0.5 \ln(2) (\text{current concentration}) \text{ mg/h.}$$

## Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

## Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

## Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream,  $y(0) = 0$ ,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

## Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream,  $y(0) = 0$ ,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

- Thus at time  $t$  the concentration is

$$y(t) = 28.9 - 28.9e^{-0.3t} = 28.9(1 - e^{-0.3t})$$