### Math 3B: Lecture 3

Noah White

October 4, 2017

# AAP tutoring

• AAP Peer Learning Facilitators available for Math 3B.

## AAP tutoring

- AAP Peer Learning Facilitators available for Math 3B.
- Sign-ups start Wednesday (10/4/17) and the first session begins the following week.

## AAP tutoring

- AAP Peer Learning Facilitators available for Math 3B.
- Sign-ups start Wednesday (10/4/17) and the first session begins the following week.
- If you are not a part of AAP, applications for this program are found in Campbell Hall!

Last time, we spoke about

• Graphing using calculus

- Graphing using calculus
- Horizontal asymptotes

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative


Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

Note: Problem set 2 and homework will be posted today.

# Example time

... On the board.



 $\bullet$  An asymptote is a straight line which the function approaches as  $x\to\pm\infty$ 

- $\bullet$  An asymptote is a straight line which the function approaches as  $x\to\pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

• We then know the function has a slanted asymptote y = mx + b.

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

- We then know the function has a slanted asymptote y = mx + b.
- To find *b*:

$$b = \lim_{x \to \pm \infty} (f(x) - mx)$$

# Example time

... On the board.