

Math 3B: Lecture 17

Noah White

November 8, 2017

Modelling using differential equations

Question

Why are differential equations important?

Modelling using differential equations

Question

Why are differential equations important?

Answer

They describe nature!

Modelling using differential equations

Question

Why are differential equations important?

Answer

They describe nature!

Modelling using differential equations

Question

Why are differential equations important?

Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function $y(t)$ that describes something we are interested in (e.g. population/mass/etc)

Modelling using differential equations

Question

Why are differential equations important?

Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function $y(t)$ that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)

Modelling using differential equations

Question

Why are differential equations important?

Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function $y(t)$ that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)
- We can't do this directly, but we can write down an ODE that y satisfies instead.

Example 1

The senario

Want to find a function $N(t)$, descibing the sie of a population at time t (in years). Assume

Example 1

The senario

Want to find a function $N(t)$, descibing the sie of a population at time t (in years). Assume

- The population is 100 people when $t = 0$

Example 1

The senario

Want to find a function $N(t)$, descibing the sie of a population at time t (in years). Assume

- The population is 100 people when $t = 0$
- No immigration or emmigration

Example 1

The senario

Want to find a function $N(t)$, descibing the sie of a population at time t (in years). Assume

- The population is 100 people when $t = 0$
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

$bN(t)$ births per year, for some b

Example 1

The senario

Want to find a function $N(t)$, descibing the sie of a population at time t (in years). Assume

- The population is 100 people when $t = 0$
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

$bN(t)$ births per year, for some b

- Number of deaths is proportional to the total number of people. So

$dN(t)$ deaths per year, for some d

Example 1

The total change in population at time t is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$

Example 1

The total change in population at time t is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$

In real life we would determine b and d experimentally.

Example 1

The total change in population at time t is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$

In real life we would determine b and d experimentally. Let $r = b - d$. the **instinsic growth rate**. So our model is

$$\frac{dN}{dt} = rN.$$

and we know $N(0) = 100$.

Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so $N(t) = 100$ for all t).

Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so $N(t) = 100$ for all t).

Case 2: $r > 0$

The population is increasing indefinitely.

Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so $N(t) = 100$ for all t).

Case 2: $r > 0$

The population is increasing indefinitely.

Case 3: $r < 0$

The population is decreasing indefinitely.

Solution to a simple ODE

Theorem

For any constant a , if y is a solution to the ODE

$$\frac{dy}{dx} = ay$$

then y is given by

$$y = Ce^{ax}$$

for some constant C .

Solution to a simple ODE

Theorem

For any constant a , if y is a solution to the ODE

$$\frac{dy}{dx} = ay$$

then y is given by

$$y = Ce^{ax}$$

for some constant C .

Next time

We will see why, but for now we can verify it is actually a solution:

$$\frac{dy}{dx} = \frac{d}{dx} Ce^{ax} = C \frac{d}{dx} e^{ax} = Cae^{ax} = ay.$$

Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C .

Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C . To determine C , we need one extra piece of information, $N(0) = 100$.

Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C . To determine C , we need one extra piece of information, $N(0) = 100$.

$$100 = Ce^{(b-d)0} \quad \text{so} \quad C = 100e^{(d-b)}.$$

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d \propto N(t)).$$

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d + kN(t)) N(t).$$

- This means

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d + kN(t)) N(t).$$

- This means

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d + kN(t)) N(t).$$

- This means

$$\frac{dN}{dt} = bN - (d + kN)N$$

Where $K = r/k$.

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d + kN(t)) N(t).$$

- This means

$$\begin{aligned}\frac{dN}{dt} &= bN - (d + kN)N \\ &= (b - d - kN)N = (r - kN)N\end{aligned}$$

Where $K = r/k$.

Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time t

$$(d + kN(t)) N(t).$$

- This means

$$\begin{aligned}\frac{dN}{dt} &= bN - (d + kN)N \\ &= (b - d - kN)N = (r - kN)N \\ &= r \left(1 - \frac{kN}{r}\right) N = r \left(1 - \frac{N}{K}\right) N\end{aligned}$$

Where $K = r/k$.

Logistic growth

The equation

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

is called the **Logistic equation** and K is the **carrying capacity**.

Behaviour of logistic growth

Assume that $r > 0$ and $K > 0$.

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

Case 1. $N(0) = 0$

In this case the growth rate is 0 initially, so $N(t)$ does not increase or decrease, so remains 0.

Behaviour of logistic growth

Assume that $r > 0$ and $K > 0$.

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

Case 1. $N(0) = 0$

In this case the growth rate is 0 initially, so $N(t)$ does not increase or decrease, so remains 0.

Case 2. $N(0) = K$

In this case the growth rate is 0 initially, so $N(t)$ does not increase or decrease, so remains K .

Behaviour of logistic growth

Assume that $r > 0$ and $K > 0$.

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

Case 1. $N(0) = 0$

In this case the growth rate is 0 initially, so $N(t)$ does not increase or decrease, so remains 0.

Case 2. $N(0) = K$

In this case the growth rate is 0 initially, so $N(t)$ does not increase or decrease, so remains K .

Key takeaway

Both $N(t) = 0$ and $N(t) = K$ are solutions to the ODE. They are called **equilibrium solutions**.

Behaviour of logistic growth

Assume that $r > 0$ and $K > 0$.

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

Case 3. $0 \leq N(0) \leq K$

In this case, N is initially increasing and so becomes more positive, slowing down as it gets close to K .

Behaviour of logistic growth

Assume that $r > 0$ and $K > 0$.

$$\frac{dN}{dt} = r \left(1 - \frac{N}{K} \right) N$$

Case 3. $0 \leq N(0) \leq K$

In this case, N is initially increasing and so becomes more positive, slowing down as it gets close to K .

Case 4. $N(0) \geq K$

In this case N is initially decreasing but decreases slower and slower as it gets close to K .

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

$$\frac{dy}{dx} = e^{\sin x} \cos x$$

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

$$\begin{aligned}\frac{dy}{dx} &= e^{\sin x} \cos x \\ y \cos x &= e^{\sin x} \cos x\end{aligned}$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

4. solve for y !

Examples

On the board...

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Examples

$$\frac{dy}{dt} = ay, \quad \frac{dy}{dt} = -\lambda y.$$

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream

Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream
- pollutant in water supply

General solution

Using separation of variables, we can show that the general solution to

$$\frac{dy}{dt} = a - by$$

is

$$y(t) = \frac{a}{b} - Ce^{-bt}$$

where C is an arbitrary constant.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = M e^{-0.5t \ln(2)} \text{ mg left}$$

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = Me^{-0.5t \ln(2)} \text{ mg left}$$

- Thus the rate at which the drug is leaving (at time t) is given by

$$0.5 \ln(2) Me^{-0.5t \ln(2)} = 0.5 \ln(2)(\text{current concentration}) \text{ mg/h.}$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

- Thus at time t the concentration is

$$y(t) = 28.9 - 28.9e^{-0.3t} = 28.9(1 - e^{-0.3t})$$

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

$$\frac{dT}{dt} = k(A - T)$$

General solution

$$T(t) = A - Ce^{-kt}.$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.
- Thus

$$90 = 70 - C \quad \text{so} \quad C = -20.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

- Rearranging we get $20e^{-0.01t} = 5$ i.e.

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

- So we get

$$t = -100 \ln\left(\frac{1}{4}\right) \approx 138 = 2 \text{ hours } 18 \text{ minutes.}$$