

Midterm 1 practice 2

UCLA: Math 115A, Spring 2019

Instructor: Noah White

Date:

Version: 1

- This exam has 4 questions, for a total of 20 points.
- Please print your working and answers neatly.
- Write your solutions in the space provided showing working.
- All final answers should be exact values. Decimal approximations will not be given credit.
- Indicate your final answer clearly.
- Full points will only be awarded for solutions with correct working.
- You may write on the reverse of a page or on the blank pages found at the back of the booklet however these will not be graded unless very clearly indicated.
- Non programmable and non graphing calculators are allowed.

Name: _____

ID number: _____

| Question | Points | Score |
|----------|--------|-------|
| 1 | 5 | |
| 2 | 5 | |
| 3 | 5 | |
| 4 | 5 | |
| Total: | 20 | |

Question 1 is multiple choice. Once you are satisfied with your solutions, indicate your answers by marking the corresponding box in the table below.

Please note! The following three pages will not be graded. You must indicate your answers here for them to be graded!

Question 1.

| <i>Part</i> | A | B | C | D |
|-------------|---|---|---|---|
| (a) | | | | |
| (b) | | | | |
| (c) | | | | |
| (d) | | | | |
| (e) | | | | |

1. Each of the following questions has exactly one correct answer. Choose from the four options presented in each case. No partial points will be given.

- (a) (1 point) In the vector space $\mathbb{C}[x]$, the set $\{x^2 - x, x^2 + 1\}$ is
- A. linearly dependent
 - B. linearly independent
 - C. a spanning set
 - D. none of the above

The following two questions concern the subsets

$$U = \left\{ \begin{pmatrix} a \\ b \\ c \end{pmatrix} \mid a^2 + b^2 = c \right\} \subseteq \mathbb{R}^3$$
$$V = \{ p(x) \in \mathbb{R}_2[x] \mid p(1) = \lambda \} \subseteq \mathbb{R}_2[x]$$

for some choice of $\lambda \in \mathbb{R}$. Recall that $\mathbb{R}_2[x]$ is the space of polynomials of degree at most 2.

- (b) (1 point) Which of the following is a true statement?
- A. Both U and V are subspaces regardless of the value of $\lambda \in \mathbb{R}$.
 - B. Only U is a subspace.
 - C. V is a subspace for any λ .
 - D. Only V is a subspace when $\lambda = 0$.

- (c) (1 point) When $\lambda = 0$, the subspace V has dimension
- A. 1
 - B. 2
 - C. 3
 - D. 4

(d) (1 point) Let U and W be two, finite dimensional subspaces of a vector space V . Which of the following statements is true?

- A. We must have $U \cap W = \{0\}$.
- B. If $U \cap W = \{0\}$ then $U + W = V$.
- C. $\dim U + \dim W \geq \dim(U + W)$.
- D. The dimension of $U + W$ is unrelated to $\dim U$ and $\dim W$.

(e) (1 point) Which of the following definitions, makes $T : \mathbb{R}_2[x] \longrightarrow \mathbb{R}^2$ into a surjective linear map?

- A. $T(p) = \begin{pmatrix} 0 \\ p(1) \end{pmatrix}$
- B. $T(p) = \begin{pmatrix} p(1) \\ p(1) \end{pmatrix}$
- C. $T(p) = \begin{pmatrix} p'(1) \\ p(1) \end{pmatrix}$
- D. $T(p) = \begin{pmatrix} p - 1 \\ p + 1 \end{pmatrix}$

2. Give (simple) examples of all of the following situations.

(a) (2 points) Two subspaces U, V of $\text{Mat}_{2 \times 2}(\mathbb{R})$ such that $U+V = \text{Mat}_{2 \times 2}(\mathbb{R})$ but $\text{Mat}_{2 \times 2}(\mathbb{R}) \neq U \oplus V$.

(b) (2 points) A basis for each of your subspaces U and V above.

(c) (1 point) A basis for $\text{Mat}_{2 \times 2}(\mathbb{R})$ that does not contain either of the bases from the previous part.

3. Consider the following maps. Prove or disprove that they are linear and if linear, find the dimension of the kernel (nullspace).

(a) (1 point) $T : \mathbb{R}^2 \longrightarrow \mathbb{R}$ given by $T(a, b) = a^2 - b^2$.

(b) (4 points) $R : \text{Mat}_{2 \times 2}(\mathbb{C}) \longrightarrow \mathbb{C}^2$ given by

$$R(M) = M \cdot \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

4. (a) (2 points) Let V and W be vector spaces and let $T : V \longrightarrow W$ be a linear map. Prove that the kernel (nullspace), $\ker T \subset V$ is a subspace.

- (b) (3 points) Suppose W_1 and W_2 are two subspaces of a vector space V such that $V = W_1 \oplus W_2$. If B_1 and B_2 are bases of W_1 and W_2 respectively, show that $B_1 \cup B_2$ is a basis for V .

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.