Math 3B: Lecture 22

Noah White

November 28, 2018

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t} = f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

We want points (t, y) such that f(y) = 0.

• Suppose f(a) = 0.

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

We want points (t, y) such that f(y) = 0.

- Suppose f(a) = 0.
- Then (t, a) is on the nullcline, for any t.

Definition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t} = f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property


The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

We want points (t, y) such that f(y) = 0.


- Suppose f(a) = 0.
- Then (t, a) is on the nullcline, for any t.
- So the line y = a is part of the nullcline, whenever f(a) = 0.

Slope fields and nullclines for autonomous systems

Phase lines/diagram



Phase lines/diagram



Phase lines/diagram



Recipe to draw phase lines

1. Draw a vertical corresponding to y axis

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- 3. Draw up arrows on intervals between dots where the derivative is positive

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- 3. Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition

 An equalibrium is stable if the two arrows are pointing towards it.

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition


- An equalibrium is stable if the two arrows are pointing towards it.
- It is unstable if the two arrows are pointing away from it.

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative


Definition


- An equalibrium is stable if the two arrows are pointing towards it.
- It is unstable if the two arrows are pointing away from it.
- It is semistable if the arrows point in the same direction.





$$\frac{\mathrm{d}y}{\mathrm{d}t} = y(y-10)(25-y)$$


$$\frac{\mathrm{d}y}{\mathrm{d}t} = y(y-10)(25-y)$$









































Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

- (i.e. f(a) = 0) then a is
 - stable if f'(a) < 0

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

- stable if f'(a) < 0
- unstable if f'(a) > 0

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

- stable if f'(a) < 0
- unstable if f'(a) > 0
- indeterminate if f'(a) = 0

