

# Math 3B: Lecture 6

Noah White

January 22, 2017

# Introduction

Last time

- Examples of maxima/minima problems

Quiz 3

# Introduction

Last time

- Examples of maxima/minima problems
- Maximising functions with constraints

Quiz 3

# Introduction

Last time

- Examples of maxima/minima problems
- Maximising functions with constraints
- Competing economies

Quiz 3

# Introduction

Last time

- Examples of maxima/minima problems
- Maximising functions with constraints
- Competing economies

Quiz 3

- Arrive on time

# Introduction

Last time

- Examples of maxima/minima problems
- Maximising functions with constraints
- Competing economies

Quiz 3

- Arrive on time
- You will have 10 mins

# Introduction

Last time

- Examples of maxima/minima problems
- Maximising functions with constraints
- Competing economies

Quiz 3

- Arrive on time
- You will have 10 mins
- Look at questions 1-8 on PS3

# Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$



# Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

The **challenge** is to find all the functions  $y = f(x)$  (or even just one) that satisfy a given equation.

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If  $h(t)$  measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If  $h(t)$  measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

The force due to gravity is roughly  $-10m$  Newtons, so

$$-10m = mh''(t)$$

## Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*

# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*

# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*

If  $P(t)$  is the population at time  $t$ :

$$\frac{dP}{dt} = rP(t)$$



## Some more examples of differential equations

$$\frac{dy}{dx} = f(x)$$

$$\frac{dy}{dx} = y(1 - y)$$

$$y'' = \sqrt{a^2 - (y')^2}$$

$$\frac{dy}{dt} = k(A - y)^2$$

# Antiderivatives

We will be concentrating on solving differential equations of the form

$$\frac{dy}{dx} = f(x)$$

# Antiderivatives

We will be concentrating on solving differential equations of the form

$$\frac{dy}{dx} = f(x)$$

A solution  $y = F(x)$  is called an **antiderivative** of  $f(x)$ .

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + 4$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + 8$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + C$$



## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2 - x$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2 - x + C$$

## Example 3

### Question

What is the antiderivative of  $f(x) = e^{2x}$ ?

## Example 3

### Question

What is the antiderivative of  $f(x) = e^{2x}$ ?

### Solution

$$F(x) = \frac{1}{2}e^{2x}$$

## Example 4

### Question

What is the antiderivative of  $f(x) = \frac{1}{x}$  (for  $x > 0$ )?



## Example 4

### Question

What is the antiderivative of  $f(x) = \frac{1}{x}$  (for  $x > 0$ )?

### Solution

$$F(x) = \ln x$$

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

### Solution

Note that  $f(x) = (1+x)^{-2}$ . So

$$F(x) = \frac{1}{1+x}$$

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

### Solution

Note that  $f(x) = (1+x)^{-2}$ . So

$$F(x) = -\frac{1}{1+x}$$

## Example 6

### Question

What is the antiderivative of  $f(x) = 2x \cos x^2$ ?

## Example 6

### Question

What is the antiderivative of  $f(x) = 2x \cos x^2$ ?

### Solution

$$F(x) = \sin x^2$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$



## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$

$$F(x) = x^{\frac{1}{2}}$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$


$$F(x) = 2x^{\frac{1}{2}}$$

# Slope fields

In some cases it is impossible to find the antiderivative (without special functions). E.g.

$$f(x) = e^{-x^2}$$

But we can still (approximately) graph the antiderivative! First we draw the slope field



# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.

## Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?

# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$

# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$



# Slope fields (how to draw)


How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$
5. Do this for a grid of points on the  $xy$ -plane.


## Example 1

$$f(x) = e^{-x^2}$$




## Example 1

$$f(x) = e^{-x^2}$$




## Example 2

$$f(x) = \sin(x^2)$$



## Example 2

$$f(x) = \sin(x^2)$$



# Accumulated change

Often we encounter problems involving **accumulated change**.

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?



# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

These problems involve finding the area under some curve.

## Example 1


If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

## Example 1

If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

### Solution

We model the car's speed using the function  $s(t) = 30$ . So we can see that the area under this curve



is the distance travelled (75 miles)

## Example 2


If a car accelerates for 20 seconds at a rate of  $2m/s^2$  and then decelerates for 30 seconds at a rate of  $1m/s^2$ , how far has it travelled?

## Example 2

If a car accelerates for 20 seconds at a rate of  $2\text{m/s}^2$  and then decelerates for 30 seconds at a rate of  $1\text{m/s}^2$ , how far has it travelled?


### Solution

The car's speed is given by  $s(t) = 2t$  when  $0 \leq t \leq 20$  and  $s(t) = 60 - t$  when  $20 \leq t \leq 50$ . So the graph looks like



## More complicated areas

How do we calculate the area under more complicated curves?



## More complicated areas

How do we calculate the area under more complicated curves?

