

Midterm 2 practice

UCLA: Math 3B, Fall 2016

Instructor: Noah White
Date: Monday, November 21, 2016
Version: *practice*.

- This exam has 3 questions, for a total of 40 points.
- Please print your working and answers neatly.
- Write your solutions in the space provided showing working.
- Indicate your final answer clearly.
- You may write on the reverse of a page or on the blank pages found at the back of the booklet however these will not be graded unless very clearly indicated.
- Non programmable and non graphing calculators are allowed.

Name: _____

ID number: _____

Discussion section: _____

| Question | Points | Score |
|----------|--------|-------|
| 1 | 14 | |
| 2 | 13 | |
| 3 | 13 | |
| Total: | 40 | |

1. A hemispherical hole needs to be dug in order to plant a tree. The hole should be 2 m in diameter. Recall that the volume of a sphere, radius r , is given by

$$V = \frac{4}{3}\pi r^3.$$

- (a) (1 point) What is the total volume of the hole?

Solution:

$$\frac{1}{2} \cdot \frac{4}{3}\pi = \frac{2}{3}\pi \text{ m}^3$$

- (b) (1 point) Suppose we divide the hole into n slices, horizontally, of constant thickness Δh . What is Δh in terms of n ?

Solution:

$$\Delta h = \frac{h}{n}$$

- (c) (1 point) Let $h = 0$ be the top of the hole and $h = 1$ be the bottom. Let h_k be the depth of the top of the k^{th} slice (where the first slice is the 0^{th} slice), this $h_0 = 0$. What is h_k in terms of Δh and k ?

Solution:

$$h_k = k\Delta h$$

- (d) (2 points) At a depth of $h = h_k$, what is the radius of the hole?

Solution:

$$\sqrt{1 - h_k^2}$$

- (e) (2 points) If n is large enough, we can approximate each slice by a cylinder of height Δh and radius as in part (d). Using this approximation, what is the volume of the k^{th} slice?

Solution:

$$\pi(1 - h_k^2)\Delta h \text{ m}^3$$

- (f) (3 points) The work needed to lift m kg of material, d meters up is given by

$$W = 10md$$

where we have approximated the acceleration due to gravity as 10 m/s^2 . How much work needs to be done in order to lift the k^{th} slice out of the hole if we assume 1 m^3 of dirt weighs 1000 kg ?

Solution:

$$10000\pi(1 - h_k^2)h_k\Delta h$$

- (g) (2 points) Write a Riemann sum which represents the total amount of work needed to dig the dirt out of the hole.

Solution:

$$W = \lim_{n \rightarrow \infty} \Delta h \sum_{k=0}^{n-1} 10000\pi(1 - h_k^2)h_k$$

- (h) (2 points) Use an integral to evaluate the Riemann sum above.

Solution:

$$\begin{aligned} W &= \int_0^1 10000\pi(1 - h^2)h \, dh \\ &= \left[10000\pi \left(h - \frac{1}{3}h^3 \right) \right]_0^1 \\ &= 10000\pi \left(1 - \frac{1}{3} \right) - 10000\pi(0) \\ &= \frac{20000}{3}\pi \end{aligned}$$

2. Solve the following differential equations. If no initial condition is given, find the general solution.

(a) (4 points) $\frac{dy}{dx} = \frac{y}{2x+4}$.

Solution:

$$y = C\sqrt{2x-4}$$

(b) (4 points) $\frac{dy}{dt} = \frac{y}{2t+4}$ where $y(1) = 0$.

Solution:

$$y = \sqrt{\ln\left(\frac{1}{3-2t}\right)}$$

(c) (5 points) $\frac{dy}{dx} = e^{x+y}$ where $y(0) = 0$.

Solution:

$$y(x) = -\ln(2 - e^e)$$

3. A river flows into a small lake and another river flows out of the lake such that the lake has a constant volume of 2000 m^3 (the rate of water flowing in equals the rate of water flowing out). The river flowing into the lake contains a pollutant present at 0.5 g/m^3 . In this question you will model the total amount of pollutant, $y(t)$, present at time t (Note that $y(t)$ is the total amount of pollutant in the lake and not a concentration).

- (a) (1 point) Assume that the river flowing in, flows at a constant rate of $20 \text{ m}^3/\text{h}$. At what rate is the pollutant flowing into the lake (in mg/h)?

Solution: Every hour there is 0.5 grams of pollutant entering the lake *per meter cubed* of water. Since there are 20 m^3 of water entering the lake every hour, there is 10 g/h of pollutant entering the lake.

- (b) (3 points) Under the above assumption, write a differential equation describing the change in the level of pollution in the lake.

Solution: The differential equation will take the form

$$\frac{dy}{dt} = \text{rate in} - \text{rate out}.$$

Thus we need to find the rate out. There are 20 m^3 flowing out every hour. At time t the concentration of pollutant in the lake is

$$\frac{y(t)}{2000} \text{ g/m}^3.$$

Thus at time t there is

$$\frac{20y(t)}{2000} = \frac{y(t)}{100} \text{ g/h}$$

of pollutant leaving the lake. Thus our differential equation is

$$\frac{dy}{dt} = 10 - \frac{y}{100}$$

- (c) (3 points) Assuming that initially there is no pollutant in the lake, solve this differential equation.

Solution: We can use the general solution of the linear model to get

$$y(t) = 1000 - Ce^{-0.01t}.$$

We assume that $y(0) = 0$ to get that

$$0 = 1000 - C$$

so the final solution is

$$y(t) = 1000 (1 - e^{-0.01t})$$

- (d) (5 points) Now assume that there is some seasonal variability and that the river flowing in (and thus also the river flowing out), flow at a rate of $40 \sin^2 t \text{ m}^3/\text{h}$. Write and solve a differential equation to model this situation, assuming there is initially no pollution in the lake.

Solution: Here we repeat the analysis above with the changed assumption. At time t , there is $40 \sin^2 t \text{ m}^3$ of water entering the lake every hour. Thus there is $20 \sin^2 t \text{ g/h}$ of pollutant entering the lake at time t .

Now, at time t , there is $y(t)$ grams of pollutant in the lake and thus the concentration of pollutant is

$$\frac{y(t)}{2000} \text{ g/m}^3.$$

Thus there is

$$\frac{40y(t) \sin^2 t}{2000} = \frac{y(t) \sin^2 t}{50} \text{ g/h}$$

flowing out of the lake. The differential equation is

$$\frac{dy}{dt} = 20 \sin^2 t - \frac{y(t) \sin^2 t}{50} = \left(20 - \frac{y}{50}\right) \sin^2 t.$$

To solve this we separate variables and integrate

$$\int \frac{50}{1000 - y} dy = \int \sin^2 t dt.$$

We use the hint to obtain

$$-50 \ln(1000 - y) = \frac{1}{2}(t - \sin(t) \cos(t)) + C.$$

Rearranging we get that the solution is

$$y(t) = 1000 - C \exp(-0.01(t - \sin(t) \cos(t)))$$

We can use the fact that $y(0) = 0$ to get

$$C = 1000$$

so the final solution is

$$y(t) = 1000 \left(1 - e^{-0.01(t - \sin(t) \cos(t))}\right).$$

- (e) (1 point) Compare the long term behaviour of the two solutions.

Solution: In the long term, both solution approach 1000 as the $\sin(t) \cos(t)$ term becomes insignificant.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.