

This weeks problem set focuses on the ideas of linear combinations, linear dependence and bases. A question marked with a  $\dagger$  is difficult and probably too hard for an exam (though still illustrates a useful point). A question marked with a  $*$  is especially important.

**Homework:** due Friday 10 April, uploaded to Gradescope before 11:59pm: questions 3,4, and 5 below.

1. From section 1.4, problems 1, 7, 8 ( $P_n(F)$  is the set of polynomials of degree less than or equal to  $n$ ), 11, 12, 13\*.
2. From section 1.5, problems 1, 2a, c, e, 4\*, 5, 9\*, 15, 18\*.

**Quotient spaces:** Let  $V$  be a vector space over a field  $\mathbb{F}$  and  $W$  a subspace of  $V$ . For any  $v \in V$ , consider the set  $\{v\} + W = \{v + w \mid w \in W\}$ . We will denote it simply as  $v + W$ . Now consider the set

$$V/W = \{v + W \mid v \in V\}.$$

We can define addition and scalar multiplication on this set by

$$(v + W) + (w + W) = (v + w) + W \quad \text{and} \quad \lambda(v + W) = \lambda v + W.$$

It turns out this is a vector space, it is called the *quotient* of  $V$  by  $W$ . See below.

3. Let  $V = \mathbb{R}^2$  and  $W = \text{span} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$ . List all the elements of  $V/W$ , making sure not to list any element twice.

**Solution:** Consider the set

$$\begin{aligned} \begin{pmatrix} a \\ b \end{pmatrix} + W &= \left\{ \begin{pmatrix} a \\ b \end{pmatrix} + \begin{pmatrix} \lambda \\ \lambda \end{pmatrix} \mid \lambda \in \mathbb{R} \right\} \\ &= \left\{ \begin{pmatrix} a + \lambda \\ b + \lambda \end{pmatrix} \mid \lambda \in \mathbb{R} \right\} \\ &= \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \mid x - y = a - b \right\} \end{aligned}$$

So  $\begin{pmatrix} a \\ b \end{pmatrix} + W$  is the line  $y = x + (b - a)$ . Thus we can say  $V/W$  is the set of lines with slope 1. The line  $y = x + b$  is the element  $\begin{pmatrix} 0 \\ b \end{pmatrix} + W$  and is clearly not equal to any other line.

*Solution 2:* Another way that one could say the above is that  $\begin{pmatrix} a \\ b \end{pmatrix} + W = \begin{pmatrix} 0 \\ b - a \end{pmatrix} + W$ . Furthermore,  $\begin{pmatrix} 0 \\ b \end{pmatrix} + W = \begin{pmatrix} 0 \\ b' \end{pmatrix} + W$  only if  $b = b'$ . Thus every element in  $V/W$  is equal to  $\begin{pmatrix} 0 \\ b \end{pmatrix} + W$  for some unique  $b$ .

- 4\* Prove that  $V/W$  is a vector space.

**Solution:** We simply check each axiom one at a time.

**VS1** Clearly  $(v + W) + (u + W) = (v + u) + W = (u + v) + W = (u + W) + (v + W)$  since  $v + u = u + v$  in  $V$ .

**VS2** This holds in exactly the same way as above since  $(v + u) + w = v + (u + w)$  in  $V$ .

**VS3** The zero element is  $0 + W = W$ . Indeed

$$(0 + W) + (v + W) = (0 + v) + W = v + W$$

Note that  $0$  is the zero element of  $V$  and we denote the zero element of  $V/W$  by  $0 + W$  or  $W$  for clarity.

**VS4** The additive inverse of  $v + W$  is  $-v + W$ . Indeed

$$(v + W) + (-v + W) = (v - v) + W = 0 + W$$

since  $v - v = 0$  in  $V$ .

**VS5** It is clear that  $1 \cdot (v + W) = (1v) + W = v + W$  since  $1v = v$  in  $V$ .

**VS6-8** These all follow in exactly the same way as above. The relations are true in  $V$  so they are true in  $V/W$ .

5. Let  $\mathbb{C}[x]$  be the vector space of polynomials and let  $W = \text{span}\{x^{2a} \mid a \geq 0\}$ .

(a) Find a set of 3 linearly independent elements of  $\mathbb{C}[x]/W$ .

**Solution:** Note that  $x^a + W = W$  if  $a$  is even. Thus we can choose  $x + W, x^3 + W, x^5 + W$ . These are linearly independent since if

$$a(x + W) + b(x^3 + W) + c(x^5 + W) = W$$

then

$$(ax + bx^3 + cx^5) + W = W$$

and thus  $ax + bx^3 + cx^5 \in W$ , but this is impossible unless  $a = b = c = 0$ .

(b) Find 2 nonzero elements  $p, q \in \mathbb{C}[x]$  that are linearly independent and such that  $p + W$  and  $q + W$  are linearly dependant and nonzero. *Note: you can only receive full points for this problem if your polynomials  $p$  and  $q$  are different from everyone else's! If you understand the problem then this will be easy to ensure.*

**Solution:** We want two elements  $p, q$  such that they are linearly independent. The easiest way to ensure this is to pick two polynomials with different degrees. We also want  $p + W$  and  $q + W$  to be non-zero. That is, we need to make sure  $p, q \notin W$ . The easiest way to ensure this is to make sure they have, for example, a linear term. Thirdly we want to make sure  $p + W$  and  $q + W$  are linearly dependent. The easiest way to ensure this is to make sure  $p + W = q + W$ , i.e.  $(p - q) + W = W$ , that is we want  $p - q \in W$ . This means their odd degree terms should agree. Lets summarise the conditions we want  $p, q$  to have.

- They should have different degrees.
- They should have a linear term
- Their odd degree terms should be the same.

So we could pick, for example,  $p = x$  and  $q = 1 + x$ .

**Note on quotient spaces:** The astute reader might be worried that the addition and scalar multiplication might not be well defined. What do I mean by this? Well, it is entirely possible that  $v + W = v' + W$  for two different elements  $v, v' \in V$ . This means we could calculate a sum in two different ways. As

$$(v + W) + (u + W) = (v + u) + W$$

or as

$$(v + W) + (u + W) = (v' + W) + (u + W) = (v' + u) + W$$

(since  $v + W = v' + W$ ). So we need to check that  $(v + u) + W = (v' + u) + W$ . I will show you how to do this below. You might like to try to prove that the scalar multiplication is unambiguous for yourself.

*Proof that  $(v + u) + W = (v' + u) + W$ :* Note that  $(v + u) + W = \{(v + u) + w \mid w \in W\}$  and  $(v' + u) + W = \{(v' + u) + w \mid w \in W\}$ . Also note that  $v \in v + W$  since  $v = v + 0$  and  $0 \in W$ .

Since  $v + W = v' + W$  we see that  $v \in v' + W$  and thus  $v = v' + x$  for some  $x \in W$ . Now let's take an arbitrary element  $s \in (v + u) + W$ , it will be of the form  $s = v + u + w$ . We know

$$s = v + u + w = v' + x + u + w = (v' + u) + (x + w).$$

Since  $x + u \in W$  we see that  $s = (v' + u) + (x + w) \in (v' + u) + W$ . We have just shown that  $(v + u) + W \subset (v' + u) + W$ . To complete the proof we need to show the opposite containment.

We do this in almost the same way. Take an arbitrary element  $t \in (v' + u) + W$ . We have that  $t = v' + u + w$  for some  $w \in W$ . Then

$$t = v' + u + w = v - x + u + w = (v + u) + (w - x) \in (v + u) + W.$$

Thus we have shown  $(v' + u) + W \subset (v + u) + W$  and hence  $(v + u) + W = (v' + u) + W$ .