

# Math 3B: Lecture 4

Noah White

October 14, 2019

## Last time

Last time, we spoke about

- Graphing using calculus

## Last time

Last time, we spoke about

- Graphing using calculus
- Slanted asymptotes

## Last time

Last time, we spoke about

- Graphing using calculus
- Slanted asymptotes
- Examples

# Functions

A function is three pieces of information

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .



# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

are all different functions!

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R} \longrightarrow \mathbb{R}_{\geq 0}; x \mapsto x^2$

are all different functions!

# Global Maximums and minimums

## Definition (Global maximum)

A function  $f : D \longrightarrow R$  has a global maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$

# Global Maximums and minimums

## Definition (Global maximum)

A function  $f : D \longrightarrow R$  has a global maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$


## Definition (Global minimum)

A function  $f : D \longrightarrow R$  has a global minimum at  $a$  if

$$f(x) \geq f(a) \quad \text{for all } x \in D$$

## Example of a global minimum


$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$  has a min at  $x = 0$





## Example of a global maximum

$f : (-\infty, 0] \rightarrow \mathbb{R}; f(x) = x^3$  has a max at  $x = 0$



# Local Maximums and minimums

## Definition (local maximum)

A function  $f : D \longrightarrow R$  has a local maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$

# Local Maximums and minimums

## Definition (local maximum)

A function  $f : D \longrightarrow R$  has a local maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$


## Definition (local minimum)

A function  $f : D \longrightarrow R$  has a local minimum at  $a$  if

$$f(x) \geq f(a) \quad \text{for all } x \text{ near } a$$


## Example of a local minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$  has a min at  $x = 0$




## Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$  has a local max at  $x = -\frac{1}{3}$



## Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$  has a local max at  $x = -\frac{1}{3}$



# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples



# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )
- $f(x) = \sin x$  has a critical point at  $x = \frac{\pi}{2}$  (since  $f'(x) = \cos x$ )

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )
- $f(x) = \sin x$  has a critical point at  $x = \frac{\pi}{2}$  (since  $f'(x) = \cos x$ )
- $f(x) = e^x$  doesn't have any critical points since  $f'(x) = e^x$  can never be zero

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

**Note:** All extrema are critical points, but not all critical points are extrema!



# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

**Note:** All extrema are critical points, but not all critical points are extrema!


## Example

$f : (-\infty, 1] \rightarrow \mathbb{R}; f(x) = x^3$  has critical points at

$$x = 0 \text{ and } 1$$

## Example

$f'(x) = 3x^2$  so  $f'(0) = 0$  and  $f'(1)$  is undefined.



## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

First derivative test (minimums)

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)


- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .





## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

First derivative test (maximums)

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)


- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .



## Second derivative test

Suppose  $x = a$  is a critical point of the function  $f(x)$

Second derivative test

If

## Second derivative test

Suppose  $x = a$  is a critical point of the function  $f(x)$

### Second derivative test

If

- $f''(a) > 0$  then  $f$  has a minimum at  $a$



## Second derivative test

Suppose  $x = a$  is a critical point of the function  $f(x)$

### Second derivative test

If

- $f''(a) > 0$  then  $f$  has a minimum at  $a$
- $f''(a) < 0$  then  $f$  has a maximum at  $a$

## Second derivative test

Suppose  $x = a$  is a critical point of the function  $f(x)$

### Second derivative test

If

- $f''(a) > 0$  then  $f$  has a minimum at  $a$
- $f''(a) < 0$  then  $f$  has a maximum at  $a$

## Second derivative test

Suppose  $x = a$  is a critical point of the function  $f(x)$

### Second derivative test

If

- $f''(a) > 0$  then  $f$  has a minimum at  $a$
- $f''(a) < 0$  then  $f$  has a maximum at  $a$

**Note:** If  $f''(a) = 0$  then we cannot conclude anything! E.g  $x^3$  or  $x^4$ .