

# Midterm 2 practice

## UCLA: Math 32B, Winter 2017

*Instructor:* Noah White

*Date:* February, 2017

*Version:* practice

- This exam has 4 questions, for a total of 40 points.
- Please print your working and answers neatly.
- Write your solutions in the space provided showing working.
- Indicate your final answer clearly.
- You may write on the reverse of a page or on the blank pages found at the back of the booklet however these will not be graded unless very clearly indicated.
- Non programmable and non graphing calculators are allowed.

Name: Solutions.


ID number: \_\_\_\_\_

Discussion section (please circle):

Day/TA	Ben	Gyu Eun	Robbie
Tuesday	3A	3C	3E
Thursday	3B	3D	3F

Question	Points	Score
1	10	
2	8	
3	8	
4	14	
Total:	40	

1. (a) (5 points) Compute the center of mass of the region in the  $xy$ -plane above the  $x$ -axis and below the curve  $y = 1 - x^2$ . Assume a constant mass density of 1.



$$(x_{cm}, y_{cm}) = \frac{1}{\text{Area}} \iint_D (x, y) \cdot \overset{\text{density}}{1} dA$$

$$D = \{(x, y) \mid -1 \leq x \leq 1, 0 \leq y \leq 1 - x^2\}.$$

$$\text{Area}(D) = \iint_D 1 dA = \int_{-1}^1 \int_0^{1-x^2} dy dx$$

$$= \int_{-1}^1 (1 - x^2) dx = \left[ x - \frac{1}{3} x^3 \right]_{-1}^1 = 2 - \frac{2}{3}$$

$$= \frac{4}{3}$$

$$\iint_D (x, y) dA = \int_{-1}^1 \int_0^{1-x^2} (x, y) dy dx$$

$$= \int_{-1}^1 \left[ xy, \frac{1}{2} y^2 \right]_0^{1-x^2} dx$$

$$= \int_{-1}^1 \left( x - x^3, \frac{1}{2} (1 - x^2)^2 \right) dx$$

$$= \int_{-1}^1 \left[ \frac{1}{2} x^2 - \frac{1}{4} x^4, \frac{1}{2} \left( x - \frac{2}{3} x^3 + \frac{1}{5} x^5 \right) \right]_{-1}^1 dx$$

$$= \left( 0, \frac{1}{2} \left( 1 - \frac{2}{3} + \frac{1}{5} \right) + \frac{1}{2} \left( 1 - \frac{2}{3} + \frac{1}{5} \right) \right) = \left( 0, \frac{8}{15} \right)$$

So

$$(x_{cm}, y_{cm}) = \left( 0, \frac{8}{15} \right).$$

(b) (5 points) Determine the surface area of the paraboloid

$$x^2 + y^2 = 2z, \quad 0 \leq z \leq 1.$$

$z = \frac{1}{2}(x^2 + y^2)$  so we parametrise by

$$G(u, v) = \left(u, v, \frac{1}{2}(u^2 + v^2)\right).$$

if  $z \in [0, 1]$  then  $0 \leq x^2 + y^2 \leq 2$  so

$(u, v) \in \text{Disk radius } \sqrt{2} = D.$

$$\underline{T}_u = (1, 0, u) \quad \underline{N}(u, v) = (-u, -v, 1)$$

$$\underline{T}_v = (0, 1, v) \quad \|\underline{N}(u, v)\| = \sqrt{u^2 + v^2 + 1}$$

Thus

$$\iint_S 1 \, dS = \iint_D \sqrt{u^2 + v^2 + 1} \, dA_{uv}$$

$$= \int_0^{\sqrt{2}} \int_0^{2\pi} r \sqrt{r^2 + 1} \, dr \, d\theta$$

← change to polar.

$$= \int_0^{2\pi} \left[ \frac{1}{3} (r^2 + 1)^{3/2} \right]_0^{\sqrt{2}} d\theta$$

$$= \int_0^{2\pi} \left( \frac{1}{3} 3^{3/2} - \frac{1}{3} \right) d\theta$$

$$= 2\pi \left( \sqrt{3} - \frac{1}{3} \right)$$

2. (8 points) Consider the region  $\mathcal{E}$  given by

$$0 \leq z \leq (y - x^2)^2, \quad x^2 \leq y \leq x.$$

Use the change of variables

$$x = u, y = v + u^2, z = wv^2,$$

to evaluate

$$\int_{\mathcal{E}} \frac{1}{y - x^2} dV.$$

The region becomes

$$0 \leq wv^2 \leq (y - x^2)^2$$

$$0 \leq wv^2 \leq v^2 \quad \text{i.e.} \quad 0 \leq w \leq 1$$

$$u^2 \leq v + u^2 \leq u \quad \text{i.e.} \quad 0 \leq v \leq u - u^2$$

Thus  $0 \leq u \leq 1$ .

$$\Rightarrow \mathcal{E}_0: \quad 0 \leq u, w \leq 1, \quad 0 \leq v \leq u - u^2$$

$$\iiint_{\mathcal{E}} \frac{1}{y - x^2} dV = \text{Jac} \det \begin{pmatrix} 1 & 0 & 0 \\ zu & 1 & 0 \\ 0 & 2uv & v^2 \end{pmatrix} = v^2 > 0$$

so

$$\iiint_{\mathcal{E}} \frac{1}{y - x^2} dV = \iiint_{\mathcal{E}_0} \frac{1}{v} \cancel{dV_{uvw}} \cdot v^2 dV_{uvw} = \iiint_{\mathcal{E}_0} v dV_{uvw}$$

$$= \int_0^1 \int_0^1 \int_0^{u - u^2} v dv du dw$$

$$= \int_0^1 \int_0^1 \left[ \frac{1}{2} v^2 \right]_0^{u - u^2} du dw = \int_0^1 \int_0^1 \frac{1}{2} (u^2 - 2u^3 + u^4) du dw$$

$$= \int_0^1 \left[ \frac{1}{2} \left( \frac{1}{3} u^3 - \frac{1}{2} u^4 + \frac{1}{5} u^5 \right) \right]_0^1 dw = \int_0^1 \frac{1}{2} \left( \frac{1}{3} - \frac{1}{2} + \frac{1}{5} \right) dw$$

$$= \int_0^1 \frac{1}{60} dw = \frac{1}{60}$$

3. Let  $\mathbf{F}$  be a vector field given by

$$\mathbf{F}(x, y, z) = (y \cos z - yze^x, x \cos z - ze^x, -xy \sin z - ye^x).$$

(a) (4 points) Show that  $\mathbf{F}$  is conservative.

(b) (4 points) Find a potential function for  $\mathbf{F}$ .

a)  $\mathbf{F}$  is def. on all of  $\mathbb{R}^3$  which is simply connected.

$$\text{curl}(\mathbf{F}) = \nabla \times \mathbf{F} = \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ F_1 & F_2 & F_3 \end{pmatrix}$$

$$= \left\langle (-x \sin z - e^x) - (-x \sin z - e^x), -(-y \sin z - ye^x) + (-y \sin z - ye^x), (\cos z - ze^x) - (\cos z - ze^x) \right\rangle$$

$$= \mathbf{0}$$

Thus  $\mathbf{F}$  is conservative!

b) Suppose  $\nabla \varphi = \mathbf{F}$ . Then

$$\varphi_x = y \cos z - yze^x \Rightarrow \varphi = xy \cos z - yze^x + \alpha(y, z)$$

$$\varphi_y = x \cos z - ze^x \Rightarrow \varphi = xy \cos z - yze^x + \beta(x, z)$$

$$\varphi_z = -xy \sin z - ye^x \Rightarrow \varphi = xy \cos z - yze^x + \gamma(xy)$$

so  $\alpha = \beta = \gamma$  is a const. so

$$\varphi = xy \cos z - yze^x + C.$$

4. In this question we will calculate the surface area of the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{a^2} + z^2 = 1$ .

- (a) (4 points) Find a parameterisation of the ellipsoid given above.  
 (b) (5 points) Express the surface area as a double iterated integral.  
 (c) (5 points) Evaluate the surface area when  $a = 2$ . You may use the fact that

$$\int \sqrt{1+x^2} dx = \frac{1}{2}\sqrt{1+x^2} + \frac{1}{2}\ln(\sqrt{1+x^2}+x) + C.$$

a) The sphere is parametrised by  
 $G' = (\cos \theta \sin \phi, \sin \theta \sin \phi, \cos \phi)$

so the ellipsoid is parametr. by

$$G(\theta, \phi) = (a \cos \theta \sin \phi, a \sin \theta \sin \phi, \cos \phi)$$

$$(\theta, \phi) \in [0, 2\pi] \times [0, \pi].$$

$$b) T_\theta = (-a \sin \theta \sin \phi, a \cos \theta \sin \phi, 0)$$

$$T_\phi = (a \cos \theta \cos \phi, a \sin \theta \cos \phi, -\sin \phi)$$

$$\begin{aligned} \text{so } N(\theta, \phi) &= \begin{pmatrix} -a \cos \theta \sin^2 \phi, -a \sin \theta \sin^2 \phi, -a^2 \sin^2 \theta \sin \phi \cos \phi \\ \quad \quad \quad -a^2 \cos^2 \theta \sin \phi \cos \phi \end{pmatrix} \\ &= (-a \cos \theta \sin^2 \phi, -a \sin \theta \sin^2 \phi, -a^2 \sin \phi \cos \phi) \end{aligned}$$

$$\begin{aligned} \text{so } \|N(\theta, \phi)\|^2 &= a^2 \cos^2 \theta \sin^4 \phi + a^2 \sin^2 \theta \sin^4 \phi + a^4 \sin^2 \phi \cos^2 \phi \\ &= a^2 \sin^4 \phi + a^4 \sin^2 \phi \cos^2 \phi \\ &= a^2 \sin^2 \phi (\sin^2 \phi + a^2 \cos^2 \phi) \\ &= \cancel{a^2 \sin^2 \phi} \cancel{a^2 (\sin^2 \phi)} \end{aligned}$$

PTO

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

$$= a^2 \sin^2 \phi (1 - \cos^2 \phi + a^2 \cos^2 \phi)$$

$$= a^2 \sin^2 \phi (1 + (a^2 - 1) \cos^2 \phi)$$

$$\text{so } \| \underline{N}(\theta, \phi) \| = a \sin \phi \sqrt{1 + (a^2 - 1) \cos^2 \phi}$$

$$\iint_S 1 \, dS = \iint_D a \sin \phi \sqrt{1 + (a^2 - 1) \cos^2 \phi} \, d\theta \, d\phi$$

$$\text{where } D = \{(\theta, \phi) \mid 0 \leq \theta \leq 2\pi, 0 \leq \phi \leq \pi\}.$$

$$\text{so } S.A. = \int_0^\pi \int_0^{2\pi} a \sin \phi \sqrt{1 + (a^2 - 1) \cos^2 \phi} \, d\theta \, d\phi$$

$$c) \text{ Let } x = \cos \phi. \text{ Then } \frac{dx}{d\phi} = -\sin \phi \text{ so}$$

$$S.A. = \int_0^{2\pi} - \int_{-1}^1 a \sqrt{1 + (a^2 - 1)x^2} \, dx = 2\pi a \int_{-1}^1 \sqrt{1 + (a^2 - 1)x^2} \, dx$$

$$\text{let } u = \sqrt{a^2 - 1} x \text{ then } \frac{du}{dx} = \sqrt{a^2 - 1} \text{ so}$$

$$S.A. = \frac{2\pi a}{\sqrt{a^2 - 1}} \int_{x=-1}^{x=1} \sqrt{1 + u^2} \, du$$

$$= \frac{2\pi a}{\sqrt{a^2 - 1}} \left[ \frac{1}{2} \sqrt{1 + u^2} + \frac{1}{2} \ln(\sqrt{1 + u^2} + u) \right]_{x=-1}^{x=1}$$

PTO

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

$$\begin{aligned}
 &= \frac{2\pi a}{\sqrt{a^2-1}} \left[ \frac{1}{2} \sqrt{1+(a^2-1)x^2} + \frac{1}{2} \ln \left( \sqrt{1+(a^2-1)x^2} + \sqrt{a^2-1} x \right) \right]_{x=-1}^{x=1} \\
 &= \frac{2\pi a}{\sqrt{a^2-1}} \left[ a + \frac{1}{2} \ln(a + \sqrt{a^2-1}) - a - \frac{1}{2} \ln(a - \sqrt{a^2-1}) \right] \\
 &= \frac{2\pi a}{\sqrt{a^2-1}} \ln \left( \frac{a + \sqrt{a^2-1}}{a - \sqrt{a^2-1}} \right)
 \end{aligned}$$

When  $a = 2$  :

$$S.A. = \frac{2\pi}{\sqrt{3}} \ln \left( \frac{2+\sqrt{3}}{2-\sqrt{3}} \right)$$