

Math 3B: Lecture 3

Noah White

September 28, 2016

Academic Advancement Program

AAP Peer Learning Sessions now available for Math 3B

- ONLY FOR AAP STUDENTS - apply if you qualify

Academic Advancement Program

AAP Peer Learning Sessions now available for Math 3B

- ONLY FOR AAP STUDENTS - apply if you qualify
- Sign up and view session times on myUCLA via the academics tab

Academic Advancement Program

AAP Peer Learning Sessions now available for Math 3B

- ONLY FOR AAP STUDENTS - apply if you qualify
- Sign up and view session times on myUCLA via the academics tab
- Sessions will begin Week 2

Last time

Last time, we spoke about

- Graphing using calculus

Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes

Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes

Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

Last time

Last time, we spoke about


- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

Note: The quiz will start at the beginning of the discussion section next time.

Example time

... On the board.

Slanted asymptotes



Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote $y = mx + b$.

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote $y = mx + b$.
- To find b :

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote $y = mx + b$.
- To find b :

Slanted asymptotes

- An asymptote is a straight line which the function approaches as $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote $y = mx + b$.
- To find b :

$$b = \lim_{x \rightarrow \pm\infty} (f(x) - mx)$$

Example time

... On the board.

Functions

A function is three pieces of information

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Example

The functions

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

Functions

A function is three pieces of information

- A domain, $D \subset \mathbb{R}$
- A range, $R \subset \mathbb{R}$, and
- A rule $f : D \longrightarrow R$ that assigns to every element of D an element of R .

Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R} \longrightarrow \mathbb{R}_{\geq 0}; x \mapsto x^2$

are all different functions!

Global Maximums and minimums

Definition (Global maximum)

A function $f : D \longrightarrow R$ has a global maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$

Global Maximums and minimums

Definition (Global maximum)

A function $f : D \longrightarrow R$ has a global maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$


Definition (Global minimum)

A function $f : D \longrightarrow R$ has a global minimum at a if

$$f(x) \geq f(a) \quad \text{for all } x \in D$$


Example of a global minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$ has a min at $x = 0$



Example of a global maximum

$f : (-\infty, 0] \rightarrow \mathbb{R}; f(x) = x^3$ has a max at $x = 0$



Local Maximums and minimums

Definition (local maximum)

A function $f : D \longrightarrow R$ has a local maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$

Local Maximums and minimums

Definition (local maximum)

A function $f : D \longrightarrow R$ has a local maximum at a if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$


Definition (local minimum)

A function $f : D \longrightarrow R$ has a local minimum at a if

$$f(x) \geq f(a) \quad \text{for all } x \text{ near } a$$


Example of a local minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$ has a min at $x = 0$




Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$ has a local max at $x = -\frac{1}{3}$



Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$ has a local max at $x = -\frac{1}{3}$



Critical point

Definition (Critical point)

A function $f(x)$ has a critical point at $x = a$ if $f'(a) = 0$ or if $f'(a)$ is undefined.

Critical point

Definition (Critical point)

A function $f(x)$ has a critical point at $x = a$ if $f'(a) = 0$ or if $f'(a)$ is undefined.

Examples

Critical point

Definition (Critical point)

A function $f(x)$ has a critical point at $x = a$ if $f'(a) = 0$ or if $f'(a)$ is undefined.

Examples

- $f(x) = x^2$ has a critical point at $x = 0$ (since $f'(x) = 2x$)

Critical point

Definition (Critical point)

A function $f(x)$ has a critical point at $x = a$ if $f'(a) = 0$ or if $f'(a)$ is undefined.

Examples

- $f(x) = x^2$ has a critical point at $x = 0$ (since $f'(x) = 2x$)
- $f(x) = \sin x$ has a critical point at $x = \frac{\pi}{2}$ (since $f'(x) = \cos x$)

Critical point

Definition (Critical point)

A function $f(x)$ has a critical point at $x = a$ if $f'(a) = 0$ or if $f'(a)$ is undefined.

Examples

- $f(x) = x^2$ has a critical point at $x = 0$ (since $f'(x) = 2x$)
- $f(x) = \sin x$ has a critical point at $x = \frac{\pi}{2}$ (since $f'(x) = \cos x$)
- $f(x) = e^x$ doesn't have any critical points since $f'(x) = e^x$ can never be zero

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

Note: All extrema are critical points, but not all critical points are extrema!

How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

Note: All extrema are critical points, but not all critical points are extrema!


Example

$f : (-\infty, 1] \rightarrow \mathbb{R}; f(x) = x^3$ has critical points at

$$x = 0 \text{ and } 1$$

Example

$f'(x) = 3x^2$ so $f'(0) = 0$ and $f'(1)$ is undefined.



First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)


- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (minimums)

- If $f'(x) < 0$ for x less than and close to a , and
- $f'(x) > 0$ for x greater than and close to a , then
- $f(x)$ has a minimum at a .



First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)


- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .

First derivative test

Suppose $x = a$ is a critical point for the function $f(x)$.

First derivative test (maximums)

- If $f'(x) > 0$ for x less than and close to a , and
- $f'(x) < 0$ for x greater than and close to a , then
- $f(x)$ has a maximum at a .



Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Second derivative test

Suppose $x = a$ is a critical point of the function $f(x)$

Second derivative test

If

- $f''(a) > 0$ then f has a minimum at a
- $f''(a) < 0$ then f has a maximum at a

Note: If $f''(a) = 0$ then we cannot conclude anything! E.g x^3 or x^4 .

Example

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

Example

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

Example

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

| a | $f''(a)$ |
|-----|----------|
| 0 | 6 |
| 2 | -6 |

Example

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

| a | $f''(a)$ |
|-----|----------|
| 0 | 6 |
| 2 | -6 |

- Local minimum at $x = 0$

Example

$$f(x) = -x^3 + 3x^2 - 2 \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -3x^2 + 6x = -3x(x - 2)$$

$$f''(x) = -6x + 6$$

The critical points are

| a | $f''(a)$ |
|-----|----------|
| 0 | 6 |
| 2 | -6 |

- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example

- Local minimum at $x = 0$

Example


- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example

- Local minimum at $x = 0$
- Local maximum at $x = 2$

Example

- Local minimum at $x = 0$
- Local maximum at $x = 2$



Example

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

Example

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

Example

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

The critical points are

| a | $f''(a)$ |
|-----|----------|
| 0 | -16 |

Example

$$f(x) = e^{-8x^2} \text{ (domain: } \mathbb{R}\text{)}.$$

$$f'(x) = -16xe^{-8x^2}$$

$$f''(x) = 16(16x^2 - 1)e^{-8x^2}$$

The critical points are

| a | $f''(a)$ |
|-----|----------|
| 0 | -16 |

- Local minimum at $x = 0$

Example

- Local and global minimum at $x = 0$

Example

- Local and global minimum at $x = 0$

Example

- Local and global minimum at $x = 0$

