

# Math 3B: Lecture 7

Noah White

January 25, 2017

# Introduction

Last time

- Accumulated change

This time

# Introduction

Last time

- Accumulated change
- Area under a curve

This time

# Introduction

Last time

- Accumulated change
- Area under a curve
- The definite integral

This time

# Introduction

Last time

- Accumulated change
- Area under a curve
- The definite integral
- The fundamental theorem of calculus

This time

# Introduction

Last time

- Accumulated change
- Area under a curve
- The definite integral
- The fundamental theorem of calculus

This time

- Midterm discussion

# Introduction

Last time

- Accumulated change
- Area under a curve
- The definite integral
- The fundamental theorem of calculus

This time

- Midterm discussion
- Some properties of definite integrals

# Introduction

Last time

- Accumulated change
- Area under a curve
- The definite integral
- The fundamental theorem of calculus

This time

- Midterm discussion
- Some properties of definite integrals
- The FTC



# Introduction

## Last time

- Accumulated change
- Area under a curve
- The definite integral
- The fundamental theorem of calculus

## This time

- Midterm discussion
- Some properties of definite integrals
- The FTC
- Substitution

## Midterm discussion

- Average score was 21.5.

## Midterm discussion

- Average score was 21.5.
- Exam was long

## Midterm discussion

- Average score was 21.5.
- Exam was long
- Questions were not overly difficult

## Midterm discussion

- Average score was 21.5.
- Exam was long
- Questions were not overly difficult
- Approximate letter grades

## Midterm discussion

- Average score was 21.5.
- Exam was long
- Questions were not overly difficult
- Approximate letter grades

## Midterm discussion

- Average score was 21.5.
- Exam was long
- Questions were not overly difficult
- Approximate letter grades


Grade	Range
A	26+
B	21-25
C	14-20

# Reminder

## Defintion

The definite integral of a function  $f(x)$  is defined to be

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(a + k\Delta x) \Delta x$$





# Properties of definite integrals

Zero area

$$\int_a^a f(x) \, dx = 0$$

# Properties of definite integrals

Zero area

$$\int_a^a f(x) \, dx = 0$$

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

# Properties of definite integrals

## Zero area

$$\int_a^a f(x) \, dx = 0$$

## Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

## Adding areas

$$\int_a^c f(x) \, dx = \int_a^b f(x) \, dx + \int_b^c f(x) \, dx$$

# More properties of definite integrals

## Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

# More properties of definite integrals

## Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

## Linearity (scalars factor out)

$$\int_a^b \alpha f(x) \, dx = \alpha \int_a^b f(x) \, dx$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note



# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .
- every input  $x$  produces a number as an output.

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\frac{d}{dx}F(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right]\end{aligned}$$


## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$


## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx$



## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx hf(x)$





## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x)\end{aligned}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x) \\&= f(x)\end{aligned}$$

## A consequence (corollary)

### Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Why?

Well  $F(x) = \int_a^x f(t) \, dt + C$  for some  $a$  and  $C$ . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of  $a$ .



## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of  $a$ .

- Lets use  $a = 0$ .

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of  $a$ .


- Lets use  $a = 0$ .
- How should we calculate  $F(x)$ ?

## Example

Use the definition!

$$F(x) = \int_0^x |t| \, dt$$

is the area under  $|t|$ !




$$F(x) = \frac{1}{2}x^2 \quad \text{if } x \geq 0$$

## Example

If  $x \leq 0$  then

$$F(x) = \int_0^x |t| \, dt = - \int_x^0 |t| \, dt$$

is the negative of the area under  $|t|$ !



$$F(x) = -\frac{1}{2}x^2 \quad \text{if } x \leq 0$$

## Example

In summary

$$F(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } x \geq 0 \\ -\frac{1}{2}x^2 & \text{if } x \leq 0 \end{cases}$$

or

$$F(x) = \frac{1}{2}x|x|$$

# The indefinite integral

We also use the following notation for the general antiderivative:

**Definition**

$$\int f(x) \, dx = F(x) + C$$

# The indefinite integral

We also use the following notation for the general antiderivative:

## Definition

$$\int f(x) \, dx = F(x) + C$$

## Example

$$\int \sin(x) - x \, dx = -\cos(x) - \frac{1}{2}x^2 + C$$

## Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,



# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.

# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

## Substitution

Suppose  $u = g(x)$ , then

$$\int f(g(x))g'(x) \, dx = \int f(u) \, du$$

# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

## Substitution

Suppose  $u = g(x)$ , then

$$\int f(g(x)) \frac{du}{dx} dx = \int f(g(x)) g'(x) dx = \int f(u) du$$

## Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx = 2 \int \sqrt{u} \, du$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\begin{aligned} \int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C \end{aligned}$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\begin{aligned}\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C \\ &= \frac{4}{3} (x^2 + 1)^{\frac{3}{2}} + C\end{aligned}$$



# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

## Example

$$\int_0^1 4x\sqrt{x^2+1} \, dx = 2 \int_1^2 \sqrt{u} \, du$$

# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

## Example

$$\begin{aligned}\int_0^1 4x\sqrt{x^2+1} \, dx &= 2 \int_1^2 \sqrt{u} \, du \\ &= 2 \left[ \frac{2}{3} u^{\frac{3}{2}} \right]_1^2\end{aligned}$$

# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

## Example

$$\begin{aligned}\int_0^1 4x\sqrt{x^2+1} \, dx &= 2 \int_1^2 \sqrt{u} \, du \\ &= 2 \left[ \frac{2}{3} u^{\frac{3}{2}} \right]_1^2 \\ &= 2 \left( \frac{2}{3} 2^{\frac{3}{2}} - \frac{2}{3} 1^{\frac{3}{2}} \right) = \frac{4}{3} (2\sqrt{2} - 1)\end{aligned}$$