#### Math 3B: Lecture 3

Noah White

January 11, 2019

Last time, we spoke about

• Graphing using calculus

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes

Last time, we spoke about


- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

## Example time

... On the board.



- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$ 

- $\bullet$  An asymptote is a straight line which the function approaches as  $x\to\pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

• We then know the function has a slanted asymptote y = mx + b.

- An asymptote is a straight line which the function approaches as  $x \to \pm \infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x\to\pm\infty}f'(x)=m$$

- We then know the function has a slanted asymptote y = mx + b.
- To find *b*:

$$b = \lim_{x \to \pm \infty} (f(x) - mx)$$

## Example time

... On the board.