

Math 3B: Lecture 19

Noah White

February 27, 2019

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Examples

$$\frac{dy}{dt} = ay, \quad \frac{dy}{dt} = -\lambda y.$$

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

$$\frac{dT}{dt} = k(A - T)$$

General solution

$$T(t) = A - Ce^{-kt}.$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.
- Thus

$$90 = 70 - C \quad \text{so} \quad C = -20.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

- Rearranging we get $20e^{-0.01t} = 5$ i.e.

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

- So we get

$$t = -100 \ln\left(\frac{1}{4}\right) \approx 138 = 2 \text{ hours } 18 \text{ minutes.}$$

Slope fields

We want to study differential equations of the form

$$\frac{dy}{dt} = f(t, y)$$

These could be quite complicated. Most of the time, they are not solvable!

Slope fields

We want to study differential equations of the form

$$\frac{dy}{dt} = f(t, y)$$

These could be quite complicated. Most of the time, they are not solvable!

Aim

Get a qualitative understanding for how a solution behaves, given an initial condition $y(t_0) = y_0$.

Slope fields

We want to study differential equations of the form

$$\frac{dy}{dt} = f(t, y)$$

These could be quite complicated. Most of the time, they are not solvable!

Aim

Get a qualitative understanding for how a solution behaves, given an initial condition $y(t_0) = y_0$.

Key tool

Slope fields. At every point on the yt -plane we draw a small line segment (a vector) with slope $f(y, t)$.

Examples

Note

If we want to draw a slope field, we cannot actually draw a line segment for **every** point. Instead we pick a grid of points in the plane.

Examples

Note

If we want to draw a slope field, we cannot actually draw a line segment for **every** point. Instead we pick a grid of points in the plane.

Examples

Note

If we want to draw a slope field, we cannot actually draw a line segment for **every** point. Instead we pick a grid of points in the plane.

Examples

Lets use Geogebra! Here is the command we will use:

`SlopeField[f(x,y)]` will produce a slope field for the equation

$$\frac{dy}{dx} = f(x,y)$$

Sketching solutions

Using the slope field we can sketch rough pictures of the solution, given a starting point (an initial condition).

Note

These pictures are not supposed to be perfect. But they will hopefully give you an idea of

Sketching solutions

Using the slope field we can sketch rough pictures of the solution, given a starting point (an initial condition).

Note

These pictures are not supposed to be perfect. But they will hopefully give you an idea of

- when does the solution increase/decrease?

Sketching solutions

Using the slope field we can sketch rough pictures of the solution, given a starting point (an initial condition).

Note

These pictures are not supposed to be perfect. But they will hopefully give you an idea of

- when does the solution increase/decrease?
- what does the solution do in the long term?

Sketching solutions

Using the slope field we can sketch rough pictures of the solution, given a starting point (an initial condition).

Note

These pictures are not supposed to be perfect. But they will hopefully give you an idea of

- when does the solution increase/decrease?
- what does the solution do in the long term?
- is the solution ever above to below a certain value?

Sketching solutions

Using the slope field we can sketch rough pictures of the solution, given a starting point (an initial condition).

Note

These pictures are not supposed to be perfect. But they will hopefully give you an idea of

- when does the solution increase/decrease?
- what does the solution do in the long term?
- is the solution ever above to below a certain value?

Examples

Lets use Geogebra again.

Nullclines

Definition

The **nullcline** for $\frac{dy}{dt} = f(t, y)$ is the set of points (t, y) where $f(t, y) = 0$

Nullclines

Definition

The **nullcline** for $\frac{dy}{dt} = f(t, y)$ is the set of points (t, y) where $f(t, y) = 0$

Examples

Lets use Geogebra!

Drawing slope fields by hand

Drawing slope fields by hand can be difficult! But we can use the nullclines to get an approximate picture

Examples

Lets draw some on the board.