

Math 3B: Lecture 9

Noah White

October 16, 2018

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .

The fundamental theorem of calculus

Theorem

For any a ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .
- every input x produces a number as an output.

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Why?

Well $F(x) = \int_a^x f(t) \, dt + C$ for some a and C . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Solution

An antiderivative of $x^2 - 4$ is $\frac{1}{3}x^3 - 4x$ so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Solution

An antiderivative of $\sin x$ is $-\cos x$ so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\frac{d}{dx} F(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \end{aligned}$$

Why is the FTC true?


$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that $\frac{d}{dx} F(x) = f(x)$.

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt \end{aligned}$$


Why is the FTC true?

When h is very small $\int_x^{x+h} f(t) \approx$



Why is the FTC true?

When h is very small $\int_x^{x+h} f(t) \approx hf(x)$



Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$

Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} hf(x)\end{aligned}$$

Why is the FTC true?

We want to show that $\frac{d}{dx}F(x) = f(x)$.

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[\int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x) \\&= f(x)\end{aligned}$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Why?

Well $F(x) = \int_a^x f(t) \, dt + C$ for some a and C . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

The indefinite integral

We also use the following notation for the general antiderivative:

Definition

$$\int f(x) \, dx := F(x) + C$$

The indefinite integral

We also use the following notation for the general antiderivative:

Definition

$$\int f(x) \, dx := F(x) + C$$

Example

$$\int \sin(x) - x \, dx = -\cos(x) - \frac{1}{2}x^2 + C$$

Example

Question

Find an antiderivative of $f(x) = |x|$?

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of a .

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of a .

- Lets use $a = 0$.

Example

Question

Find an antiderivative of $f(x) = |x|$?

Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of a .


- Lets use $a = 0$.
- How should we calculate $F(x)$?

Example

Use the definition!

$$F(x) = \int_0^x |t| \, dt$$

is the area under $|t|$!




$$F(x) = \frac{1}{2}x^2 \quad \text{if } x \geq 0$$

Example

If $x \leq 0$ then

$$F(x) = \int_0^x |t| \, dt = - \int_x^0 |t| \, dt$$

is the negative of the area under $|t|$!



$$F(x) = -\frac{1}{2}x^2 \quad \text{if } x \leq 0$$

Example

In summary

$$F(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } x \geq 0 \\ -\frac{1}{2}x^2 & \text{if } x \leq 0 \end{cases}$$

or

$$F(x) = \frac{1}{2}x|x|$$

Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,

Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.

Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

Substitution

Suppose $u = g(x)$, then

$$\int f(g(x))g'(x) \, dx = \int f(u) \, du$$

Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

Substitution

Suppose $u = g(x)$, then

$$\int f(g(x)) \frac{du}{dx} dx = \int f(g(x)) g'(x) dx = \int f(u) du$$

Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

Solution

We use the substitution $u = x^2 + 1$, so $\frac{du}{dx} = 2x$, we can write the integral

$$\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx = 2 \int \sqrt{u} \, du$$

Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

Solution

We use the substitution $u = x^2 + 1$, so $\frac{du}{dx} = 2x$, we can write the integral

$$\begin{aligned} \int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C \end{aligned}$$

Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

Solution

We use the substitution $u = x^2 + 1$, so $\frac{du}{dx} = 2x$, we can write the integral

$$\begin{aligned}\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C \\ &= \frac{4}{3} (x^2 + 1)^{\frac{3}{2}} + C\end{aligned}$$

Integration by substitution (definite integrals)

Substitution for definite integrals

Suppose $u = g(x)$, then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

Integration by substitution (definite integrals)

Substitution for definite integrals

Suppose $u = g(x)$, then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

Example

$$\int_0^1 4x\sqrt{x^2+1} \, dx = 2 \int_1^2 \sqrt{u} \, du$$

Integration by substitution (definite integrals)

Substitution for definite integrals

Suppose $u = g(x)$, then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

Example

$$\begin{aligned}\int_0^1 4x\sqrt{x^2+1} \, dx &= 2 \int_1^2 \sqrt{u} \, du \\ &= 2 \left[\frac{2}{3} u^{\frac{3}{2}} \right]_1^2\end{aligned}$$

Integration by substitution (definite integrals)

Substitution for definite integrals

Suppose $u = g(x)$, then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

Example

$$\begin{aligned}\int_0^1 4x\sqrt{x^2+1} \, dx &= 2 \int_1^2 \sqrt{u} \, du \\ &= 2 \left[\frac{2}{3} u^{\frac{3}{2}} \right]_1^2 \\ &= 2 \left(\frac{2}{3} 2^{\frac{3}{2}} - \frac{2}{3} 1^{\frac{3}{2}} \right) = \frac{4}{3} (2\sqrt{2} - 1)\end{aligned}$$

The product rule

Just like integration by substitution reverses the chain rule, integration by parts "reverses" the product rule:

$$\frac{d}{dx}f(x)g(x) = f'(x)g(x) + f(x)g'(x)$$

The product rule

Just like integration by substitution reverses the chain rule, integration by parts "reverses" the product rule:

$$\frac{d}{dx} f(x)g(x) = f'(x)g(x) + f(x)g'(x)$$

written another way

$$(uv)' = u'v + uv'$$

Integration by parts

$$(uv)' = u'v + uv'$$

Integration by parts

$$(uv)' = u'v + uv'$$

Lets integrate both sides

$$\int (uv)' \, dx = \int u'v \, dx + \int uv' \, dx$$

Integration by parts

$$(uv)' = u'v + uv'$$

Lets integrate both sides

$$\int (uv)' \, dx = \int u'v \, dx + \int uv' \, dx$$

By the fundamental theorem of calculus

$$uv = \int u'v \, dx + \int uv' \, dx$$

Integration by parts

$$(uv)' = u'v + uv'$$

Lets integrate both sides

$$\int (uv)' \, dx = \int u'v \, dx + \int uv' \, dx$$

By the fundamental theorem of calculus

$$uv = \int u'v \, dx + \int uv' \, dx$$

Rearranging...

Integration by parts

The integration by parts formula

$$\int uv' \, dx = uv - \int u'v \, dx$$

Integration by parts

The integration by parts formula

$$\int uv' \, dx = uv - \int u'v \, dx$$

Alternative statement

$$\int u \, dv = uv - \int v \, du$$

Examples

One the board...