

Math 3B: Lecture 11

Noah White

February 6, 2017

Introduction

Quiz this week

- Quiz 3 questions will be drawn from problem set 5
- Notes on long division and partial fractions online

Survey

I have put a survey online:

`math.ucla.edu/~noah/3Bsurvey`

Last time

- Integration by parts

Last time

- Integration by parts
- Polynomial long division

How to deal with rational functions?

How can we integrate something like

$$\int \frac{3x^3 - 7x^2 + 17x - 3}{x^2 - 2x - 8} dx$$

or

$$\int \frac{x + 2}{x^3 - x} dx?$$

Long division of polynomials

For the first example we can rewrite it in the form

$$\frac{3x^3 - 7x^2 + 17x - 3}{x^2 - 2x - 8} = 3x - 1 + \frac{39x - 11}{x^2 - 2x - 8}$$

using polynomial long division.

Long division of polynomials

For the first example we can rewrite it in the form

$$\frac{3x^3 - 7x^2 + 17x - 3}{x^2 - 2x - 8} = 3x - 1 + \frac{39x - 11}{x^2 - 2x - 8}$$

using polynomial long division.

This is still not something we can integrate so we need to go further.

Partial fractions

When we are faced with a sum of the form

$$\frac{1}{x+1} + \frac{3}{2-3x} + \cdots$$

Partial fractions

When we are faced with a sum of the form

$$\frac{1}{x+1} + \frac{3}{2-3x} + \cdots = \frac{P(x)}{Q(x)}$$

Partial fractions

When we are faced with a sum of the form

$$\frac{1}{x+1} + \frac{3}{2-3x} + \cdots = \frac{P(x)}{Q(x)}$$

How do we reverse this process?

Answer: partial fractions

When the denominator is $(ax + b)(cx + d) \cdots$

We want to rewrite $\frac{P(x)}{Q(x)}$ as a sum. Let

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$$

When the denominator is $(ax + b)(cx + d) \cdots$

We want to rewrite $\frac{P(x)}{Q(x)}$ as a sum. Let

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$$

- If the degree of $Q(x)$ is larger than the degree of $P(x)$

When the denominator is $(ax + b)(cx + d) \cdots$

We want to rewrite $\frac{P(x)}{Q(x)}$ as a sum. Let

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$$

- If the degree of $Q(x)$ is larger than the degree of $P(x)$
- $Q(x)$ has no repeated factors. E.g. $Q(x) = (x - 1)(x + 2)$ but not $Q(x) = (x - 1)^2(x + 2)$, then

When the denominator is $(ax + b)(cx + d) \cdots$

We want to rewrite $\frac{P(x)}{Q(x)}$ as a sum. Let

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$$

- If the degree of $Q(x)$ is larger than the degree of $P(x)$
- $Q(x)$ has no repeated factors. E.g. $Q(x) = (x - 1)(x + 2)$ but not $Q(x) = (x - 1)^2(x + 2)$, then

When the denominator is $(ax + b)(cx + d) \cdots$

We want to rewrite $\frac{P(x)}{Q(x)}$ as a sum. Let

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n)$$

- If the degree of $Q(x)$ is larger than the degree of $P(x)$
- $Q(x)$ has no repeated factors. E.g. $Q(x) = (x - 1)(x + 2)$ but not $Q(x) = (x - 1)^2(x + 2)$, then

we can always find constants A_1, A_2, \dots, A_n so that

$$\frac{P(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_n}{a_nx + b_n}$$

Example 1

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x+1} + \frac{B}{x-1}$$

Example 1

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x+1} + \frac{B}{x-1}$$

Multiplying both sides by $(x-1)(x+1)$

$$\begin{aligned} 1 &= \frac{A(x-1)(x+1)}{x+1} + \frac{B(x-1)(x+1)}{x-1} \\ &= A(x-1) + B(x+1) \\ &= (A+B)x + (B-A) \end{aligned}$$

Example 1

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x+1} + \frac{B}{x-1}$$

Multiplying both sides by $(x-1)(x+1)$

$$\begin{aligned} 0x + 1 &= \frac{A(x-1)(x+1)}{x+1} + \frac{B(x-1)(x+1)}{x-1} \\ &= A(x-1) + B(x+1) \\ &= (A+B)x + (B-A) \end{aligned}$$

Comparing coefficients

$$A + B = 0 \quad \text{and} \quad B - A = 1$$

Example 1

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x+1} + \frac{B}{x-1}$$

Multiplying both sides by $(x-1)(x+1)$

$$\begin{aligned} 0x + 1 &= \frac{A(x-1)(x+1)}{x+1} + \frac{B(x-1)(x+1)}{x-1} \\ &= A(x-1) + B(x+1) \\ &= (A+B)x + (B-A) \end{aligned}$$

Comparing coefficients

$$A + B = 0 \quad \text{and} \quad B - A = 1$$

So

$$-2A = 1 \quad \text{hence} \quad A = -\frac{1}{2} \quad \text{and} \quad B = \frac{1}{2}.$$

Repeated factors

What if $q(x)$ contains repeated factors? E.g. if $q(x) = (x - 1)^2$ or $q(x) = (x - 1)(x + 2)^3$?

Repeated factors

What if $q(x)$ contains repeated factors? E.g. if $q(x) = (x - 1)^2$ or $q(x) = (x - 1)(x + 2)^3$?

For every factor $(ax + b)^k$ in $q(x)$, the partial fraction expansion has terms of the form

$$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \frac{A_3}{(ax + b)^3} + \cdots + \frac{A_k}{(ax + b)^k}.$$

Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

Comparing coefficients

$$A = 1 \quad \text{and} \quad B - A = 0$$

Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

Comparing coefficients

$$A = 1 \quad \text{and} \quad B - A = 0$$

So

$$A = 1 \quad \text{and} \quad B = 1.$$

Side note: integrating $\frac{1}{x}$.

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$

Side note: integrating $\frac{1}{x}$.

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$

Side note: integrating $\frac{1}{x}$.

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$

Using substitution this gives the formula

$$\int \frac{1}{ax + b} dx = \frac{1}{a} \ln |ax + b| + C.$$

Side note: integrating $\frac{1}{x^k}$.

Recall that if $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Side note: integrating $\frac{1}{x^k}$.

Recall that if $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Side note: integrating $\frac{1}{x^k}$.

Recall that if $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Using substitution this gives the formula

$$\int \frac{1}{(ax+b)^k} dx = -\frac{1}{a(k-1)(ax+c)^{k-1}} + C.$$

Integrating rational functions $p(x)/q(x)$

Action plan

Integrating rational functions $p(x)/q(x)$

Action plan

1. Express $\frac{p(x)}{q(x)}$ in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using polynomial long division.

Integrating rational functions $p(x)/q(x)$

Action plan

1. Express $\frac{p(x)}{q(x)}$ in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using polynomial long division.

2. Write $\frac{r(x)}{q(x)}$ as a sum of fractions of the form

$$\frac{A}{(ax + b)^k}$$

using partial fractions

Integrating rational functions $p(x)/q(x)$

Action plan

1. Express $\frac{p(x)}{q(x)}$ in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using polynomial long division.

2. Write $\frac{r(x)}{q(x)}$ as a sum of fractions of the form

$$\frac{A}{(ax + b)^k}$$

using partial fractions

3. Integrate all these pieces separately.

Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

Solution

Using long division

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1}$$

Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

Solution

Using long division and partial fractions

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1} = x^2 - 2 + \frac{1}{2(x - 1)} - \frac{1}{2(x + 1)}$$

Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

Solution

Using long division and partial fractions

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1} = x^2 - 2 + \frac{1}{2(x - 1)} - \frac{1}{2(x + 1)}$$

So

$$I = \frac{1}{3}x^3 - 2x + \frac{1}{2} \ln |x - 1| - \frac{1}{2} \ln |x + 1| + C.$$

Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

Solution

Using long division

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3}$$

Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

Solution

Using long division and partial fractions

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3} = 1 + \frac{1}{x-1} + \frac{3}{(x-1)^2} + \frac{3}{(x-1)^3}$$

Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

Solution

Using long division and partial fractions

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3} = 1 + \frac{1}{x-1} + \frac{3}{(x-1)^2} + \frac{3}{(x-1)^3}$$

So

$$I = x + \ln|x-1| - \frac{3}{x-1} - \frac{3}{2(x-1)^2} + C.$$