

# Math 3B: Lecture 19

Noah White

November 4, 2016

## Last time

- Modelling using differential equations

## Last time

- Modelling using differential equations
- Exponential population growth

## Last time

- Modelling using differential equations
- Exponential population growth
- Logistic population growth

# Logistic growth

The equation

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

is called the **Logistic equation** and  $K$  is the **carrying capacity**.

## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 1.  $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

# Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 1.  $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

Case 2.  $N(0) = K$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains  $K$ .

# Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

## Case 1. $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

## Case 2. $N(0) = K$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains  $K$ .

## Key takeaway

Both  $N(t) = 0$  and  $N(t) = K$  are solutions to the ODE. They are called **equilibrium solutions**.



## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 3.  $0 \leq N(0) \leq K$

In this case,  $N$  is initially increasing and so becomes more positive, slowing down as it gets close to  $K$ .

## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 3.  $0 \leq N(0) \leq K$

In this case,  $N$  is initially increasing and so becomes more positive, slowing down as it gets close to  $K$ .

Case 4.  $N(0) \geq K$

In this case  $N$  is initially decreasing but decreases slower and slower as it gets close to  $K$ .

## Logistic growth with outside effects

We can also modify the logistic equation to get something which models an outside effect. For example harvesting.

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N - h(t)$$

## Logistic growth with outside effects

We can also modify the logistic equation to get something which models an outside effect. For example harvesting.

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N - h(t)$$

This would model a population growing logistically but where we are harvesting at a rate of  $h(N)$ . E.g. we decide to continually harvest 3% of the population then

$$h(N) = 0.03N.$$

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

$$\frac{dy}{dx} = e^{\sin x} \cos x$$

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

$$\begin{aligned}\frac{dy}{dx} &= e^{\sin x} \cos x \\ y \cos x &= e^{\sin x} \cos x\end{aligned}$$



## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1$$

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \\ \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) &= 0\end{aligned}$$

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \\ \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) &= 0 \\ 2x + 2y'y &= 0\end{aligned}$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$



## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$

$$3 - y' \sin y = y + xy'.$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$

$$3 - y' \sin y = y + xy'.$$

### Note

We can rearrange this to get

$$y' = \frac{3 - y}{x + \sin y}$$

a differential equation. Whatever  $y$  is, as long as it obeys the above relation, it is a solution to this ODE!

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

4. solve for  $y$ !

## Examples

On the board...