

Math 3B: Lecture 1

Noah White

September 23, 2016

Syllabus

Take a copy of the syllabus as you walk in

or

find it online at math.ucla.edu/~noah

Class website

There are a few places where you will find/receive information about Math 3B:

- The class website: `www.math.ucla.edu/~noah`
- CCLE
- Email
- Piazza

Instructor and TAs

Instructor Noah White
office hours MS 6304, M 3:30pm-5pm, R 9-10:30am

TA Kyung Ha
office hours MS 6943, W 4-5pm

Robert Houseden
MS 3915B, T 3-4pm

Dustan Levenstein
MS 3965, R 3-4

Instructor and TAs

Instructor Noah White
office hours MS 6304, M 3:30pm-5pm, R 9-10:30am

TA Kyung Ha
office hours MS 6943, W 4-5pm

Robert Houseden
MS 3915B, T 3-4pm


Dustan Levenstein
MS 3965, R 3-4

Note!

Change of room for Discussion Section 2D (Thursday with Dustan).
You are now in Boelter 5419.

Textbook

S. J. Schreiber, *Calculus for the Life Sciences*, Wiley



Problem sets, homework, and quizzes

Problem sets

Assigned every week. Long list of problems. Not graded, but recommended!

Problem sets, homework, and quizzes

Problem sets

Assigned every week. Long list of problems. Not graded, but recommended!

Homework

Due every second week. A small number of questions drawn from the problem sets. There will be 5.

Problem sets, homework, and quizzes

Problem sets

Assigned every week. Long list of problems. Not graded, but recommended!

Homework

Due every second week. A small number of questions drawn from the problem sets. There will be 5.

Quizzes

Administered every other week in discussion session. There will be 4.

Homework; collection and collaboration

- Homework will be collected at the **start** of the lecture on the dues date.

Homework; collection and collaboration

- Homework will be collected at the **start** of the lecture on the dues date.
- Late homework will not be accepted.

Homework; collection and collaboration

- Homework will be collected at the **start** of the lecture on the dues date.
- Late homework will not be accepted.
- You may collaborate in small groups if:

Homework; collection and collaboration

- Homework will be collected at the **start** of the lecture on the dues date.
- Late homework will not be accepted.
- You may collaborate in small groups if:
 - ▶ you acknowledge your collaborators,

Homework; collection and collaboration

- Homework will be collected at the **start** of the lecture on the dues date.
- Late homework will not be accepted.
- You may collaborate in small groups if:
 - ▶ you acknowledge your collaborators,
 - ▶ write up your own solutions, in your own words.

Exams

There will be two midterms and a final exam

- Midterm 1 2-2:50pm Monday, 17 October, 2016

Exams

There will be two midterms and a final exam

- Midterm 1 2-2:50pm Monday, 17 October, 2016
- Midterm 2 2-2:50pm Monday, 21 November, 2016

Exams

There will be two midterms and a final exam

- Midterm 1 2-2:50pm Monday, 17 October, 2016
- Midterm 2 2-2:50pm Monday, 21 November, 2016
- Final 3-6pm Monday, 5 December, 2016

Exams

There will be two midterms and a final exam

- Midterm 1 2-2:50pm Monday, 17 October, 2016
- Midterm 2 2-2:50pm Monday, 21 November, 2016
- Final 3-6pm Monday, 5 December, 2016

Cheatsheets and calculators

You will be allowed a small cheatsheet in each exam. Must be self-written and one side, half a letter size piece of paper. You are also allowed to use non-programmable, non-graphing calculators.

Grading

Your final grade will be calculated using the maximum of the following two grading schemes.

15% (best 7 quizzes/hw) + 40% (midterms) + 45% (final)

or

15% (best 7 quizzes/hw) + 30% (best midterm) + 55% (final)

Schedule

| | Dates | Monday | Tuesday | Wednesday | Thursday | Friday |
|----|------------|--------------------|---------|--------------------|--------------|----------------------|
| 0 | 9/19-23 | No classes | | | | <i>Intro/4.1</i> |
| 1 | 9/26-30 | <i>4.1</i> | Quiz 1 | <i>4.2</i> | Quiz 1 | <i>4.3-4.4</i> |
| 2 | 10/3-7 | <i>Review</i> | | <i>5.1</i> | | HW 1 due/ <i>5.2</i> |
| 3 | 10/10-14 | <i>5.2 (cont.)</i> | Quiz 2 | <i>5.3</i> | Quiz 2 | <i>Review</i> |
| 4 | 10/17-21 | Midterm 1 | | <i>5.4</i> | | HW 2 due/ <i>5.5</i> |
| 5 | 10/24-28 | <i>5.6</i> | Quiz 3 | <i>5.6 (cont.)</i> | Quiz 3 | <i>5.8</i> |
| 6 | 10/31-11/4 | <i>Review</i> | | <i>6.1</i> | | HW 3 due/ <i>6.2</i> |
| 7 | 11/7-11 | <i>6.2 (cont.)</i> | Quiz 4 | <i>6.3</i> | Quiz 4 | Vet's day |
| 8 | 11/14-18 | <i>6.4</i> | | <i>6.4-6.5</i> | | HW 4 due/ <i>6.5</i> |
| 9 | 11/21-25 | Midterm 2 | | <i>6.6</i> | Thanksgiving | |
| 10 | 11/28-12/2 | HW 5 due | | <i>Review</i> | | <i>Review</i> |

Where to get help

Piazza

Here you can ask questions and answer others' questions. Lets take a look. . .

Where to get help

Piazza

Here you can ask questions and answer others' questions. Lets take a look. . .

Office hours

Use them wisely! Prepare and seek information first. If you learn something, post it to Piazza!

Where to get help

Piazza

Here you can ask questions and answer others' questions. Lets take a look. . .

Office hours

Use them wisely! Prepare and seek information first. If you learn something, post it to Piazza!

Student Math Center (SMC)

Location: MS 3974, times: M-R 9am-3pm.

The SMC offers free, individual and group tutoring for all lower division math courses. This service is available on a walk-in basis; no appointment is necessary. Students may ask any of the TAs in attendance for assistance with math problems.

Differentiation

You should have a good feel for what the derivative means. I.e. derivative at a point = tangent slope. You need to understand differentiation algebraically **as well as** geometrically.

Differentiation

You should have a good feel for what the derivative means. I.e. derivative at a point = tangent slope. You need to understand differentiation algebraically **as well as** geometrically.

You should also know the definition of the derivative

$$\frac{d}{dx}f(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Differentiation

You should be able to differentiate many of the standard functions we will see in this course. This includes:

- polynomials/power functions

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

Differentiation

You should be able to differentiate many of the standard functions we will see in this course. This includes:

- polynomials/power functions

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

- exponentials

$$\frac{d}{dx} (e^x) = e^x$$

Differentiation

You should be able to differentiate many of the standard functions we will see in this course. This includes:

- polynomials/power functions

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

- exponentials

$$\frac{d}{dx} (e^x) = e^x$$

- logarithms

$$\frac{d}{dx} (\ln x) = \frac{1}{x}$$

Differentiation

You should be able to differentiate many of the standard functions we will see in this course. This includes:

- polynomials/power functions

$$\frac{d}{dx} (x^n) = nx^{n-1}$$

- exponentials

$$\frac{d}{dx} (e^x) = e^x$$

- logarithms

$$\frac{d}{dx} (\ln x) = \frac{1}{x}$$

- trig functions

$$\frac{d}{dx} (\sin x) = \cos x$$

Product rule

The product rule lets us differentiate functions of the form $f(x) = g(x)h(x)$. It says

$$\frac{d}{dx}f(x) = g'(x)h(x) + g(x)h'(x)$$

Product rule

The product rule lets us differentiate functions of the form $f(x) = g(x)h(x)$. It says

$$\frac{d}{dx}f(x) = g'(x)h(x) + g(x)h'(x)$$

Example

Let's differentiate the function $f(x) = e^x \sin x$.

$$\frac{d}{dx}f(x) = \left(\frac{d}{dx}e^x\right) \sin x + e^x \left(\frac{d}{dx}\sin x\right)$$

Product rule

The product rule lets us differentiate functions of the form $f(x) = g(x)h(x)$. It says

$$\frac{d}{dx}f(x) = g'(x)h(x) + g(x)h'(x)$$

Example

Let's differentiate the function $f(x) = e^x \sin x$.

$$\begin{aligned}\frac{d}{dx}f(x) &= \left(\frac{d}{dx}e^x\right)\sin x + e^x\left(\frac{d}{dx}\sin x\right) \\ &= e^x \sin x + e^x \cos x\end{aligned}$$

Product rule

The product rule lets us differentiate functions of the form $f(x) = g(x)h(x)$. It says

$$\frac{d}{dx}f(x) = g'(x)h(x) + g(x)h'(x)$$

Example

Let's differentiate the function $f(x) = e^x \sin x$.

$$\begin{aligned}\frac{d}{dx}f(x) &= \left(\frac{d}{dx}e^x\right)\sin x + e^x\left(\frac{d}{dx}\sin x\right) \\ &= e^x \sin x + e^x \cos x \\ &= e^x(\sin x + \cos x)\end{aligned}$$

The chain rule

The chain rule is very important! It allows us to differentiate functions of the form $f(x) = g(h(x))$. It says

$$\frac{d}{dx}f(x) = h'(x)g'(h(x))$$

The chain rule

The chain rule is very important! It allows us to differentiate functions of the form $f(x) = g(h(x))$. It says

$$\frac{d}{dx}f(x) = h'(x)g'(h(x))$$

Example

Let's differentiate $f(x) = \sin(e^x)$. In this example

$$h(x) = e^x \quad \text{and} \quad g(x) = \sin x$$

The chain rule

The chain rule is very important! It allows us to differentiate functions of the form $f(x) = g(h(x))$. It says

$$\frac{d}{dx}f(x) = h'(x)g'(h(x))$$

Example

Let's differentiate $f(x) = \sin(e^x)$. In this example

$$h(x) = e^x \quad \text{and} \quad g(x) = \sin x$$

The chain rule

The chain rule is very important! It allows us to differentiate functions of the form $f(x) = g(h(x))$. It says

$$\frac{d}{dx}f(x) = h'(x)g'(h(x))$$

Example

Let's differentiate $f(x) = \sin(e^x)$. In this example

$$h(x) = e^x \quad \text{and} \quad g(x) = \sin x$$

so

$$h'(x) = e^x \quad \text{and} \quad g'(x) = \cos x$$

The chain rule

The chain rule is very important! It allows us to differentiate functions of the form $f(x) = g(h(x))$. It says

$$\frac{d}{dx}f(x) = h'(x)g'(h(x))$$

Example

Let's differentiate $f(x) = \sin(e^x)$. In this example

$$h(x) = e^x \quad \text{and} \quad g(x) = \sin x$$

so

$$h'(x) = e^x \quad \text{and} \quad g'(x) = \cos x$$

so

$$f'(x) = e^x \cos(e^x)$$

The quotient rule is stupid

The quotient rule says

$$\frac{d}{dx} \left(\frac{g(x)}{h(x)} \right) = \frac{g'(x)h(x) - g(x)h'(x)}{h(x)^2}$$

The quotient rule is stupid

The quotient rule says

$$\frac{d}{dx} \left(\frac{g(x)}{h(x)} \right) = \frac{g'(x)h(x) - g(x)h'(x)}{h(x)^2}$$

This is annoying to remember (where does that minus sign go again?). Luckily we can notice

$$\frac{g(x)}{h(x)} = g(x)k(x) \quad \text{where} \quad k(x) = (h(x))^{-1}$$

So we can just use the product rule!

Example

Question

Differentiate

$$f(x) = \sin \frac{1}{x}$$

Example

Question

Differentiate

$$f(x) = \sin \frac{1}{x}$$

Solution

We should use the chain rule. Notice $f(x) = g(h(x))$ where

$$h(x) = x^{-1} \quad \text{and} \quad g(x) = \sin x$$

Example

Question

Differentiate

$$f(x) = \sin \frac{1}{x}$$

Solution

We should use the chain rule. Notice $f(x) = g(h(x))$ where

$$h(x) = x^{-1} \quad \text{and} \quad g(x) = \sin x$$

Example

Question

Differentiate

$$f(x) = \sin \frac{1}{x}$$

Solution

We should use the chain rule. Notice $f(x) = g(h(x))$ where

$$h(x) = x^{-1} \quad \text{and} \quad g(x) = \sin x$$

so

$$h'(x) = -x^{-2} \quad \text{and} \quad g'(x) = \cos x$$

Example

Question

Differentiate

$$f(x) = \sin \frac{1}{x}$$

Solution

We should use the chain rule. Notice $f(x) = g(h(x))$ where

$$h(x) = x^{-1} \quad \text{and} \quad g(x) = \sin x$$

so

$$h'(x) = -x^{-2} \quad \text{and} \quad g'(x) = \cos x$$

so

$$\begin{aligned} f'(x) &= h'(x)g'(h(x)) \\ &= -\frac{1}{x^2} \cos(x^{-1}) \end{aligned}$$

Example

Question

Differentiate

$$f(x) = \frac{x-1}{x+1}$$

Example

Question

Differentiate

$$f(x) = \frac{x-1}{x+1} = (x-1)(x+1)^{-1}$$

Solution

We should use the product/quotient rule. Notice $f(x) = g(x)h(x)$ where

$$h(x) = (x+1)^{-1} \quad \text{and} \quad g(x) = x-1$$

$$h'(x) = -(x+1)^{-2} \quad \text{and} \quad g'(x) = 1$$

Example

Question

Differentiate

$$f(x) = \frac{x-1}{x+1} = (x-1)(x+1)^{-1}$$

Solution

We should use the product/quotient rule. Notice $f(x) = g(x)h(x)$ where

$$h(x) = (x+1)^{-1} \quad \text{and} \quad g(x) = x-1$$

$$h'(x) = -(x+1)^{-2} \quad \text{and} \quad g'(x) = 1$$

Example

Question

Differentiate

$$f(x) = \frac{x-1}{x+1} = (x-1)(x+1)^{-1}$$

Solution

We should use the product/quotient rule. Notice $f(x) = g(x)h(x)$ where

$$h(x) = (x+1)^{-1} \quad \text{and} \quad g(x) = x-1$$

$$h'(x) = -(x+1)^{-2} \quad \text{and} \quad g'(x) = 1$$

so

$$f'(x) = g'(x)h(x) + g(x)h'(x)$$

Example

Question

Differentiate

$$f(x) = \frac{x-1}{x+1} = (x-1)(x+1)^{-1}$$

Solution

We should use the product/quotient rule. Notice $f(x) = g(x)h(x)$ where

$$h(x) = (x+1)^{-1} \quad \text{and} \quad g(x) = x-1$$

$$h'(x) = -(x+1)^{-2} \quad \text{and} \quad g'(x) = 1$$

so

$$\begin{aligned} f'(x) &= g'(x)h(x) + g(x)h'(x) \\ &= \frac{1}{x+1} - \frac{x-1}{(x+1)^2} = \frac{2}{(x+1)^2} \end{aligned}$$

Example

Question

Differentiate

$$F(x) = \frac{\sin x^2 - 1}{\sin x^2 + 1}$$

Example

Question

Differentiate

$$F(x) = \frac{\sin x^2 - 1}{\sin x^2 + 1}$$

Solution

We should notice that $F(x) = f(g(x))$ so we can use the chain rule!

$$f(x) = \frac{x - 1}{x + 1} \quad \text{and} \quad g(x) = \sin x^2$$

$$f'(x) = \frac{2}{(x + 1)^2} \quad \text{and} \quad g'(x) = 2x \cos x^2$$

Example

Question

Differentiate

$$F(x) = \frac{\sin x^2 - 1}{\sin x^2 + 1}$$

Solution

We should notice that $F(x) = f(g(x))$ so we can use the chain rule!

$$f(x) = \frac{x - 1}{x + 1} \quad \text{and} \quad g(x) = \sin x^2$$

$$f'(x) = \frac{2}{(x + 1)^2} \quad \text{and} \quad g'(x) = 2x \cos x^2$$

Example

Question

Differentiate

$$F(x) = \frac{\sin x^2 - 1}{\sin x^2 + 1}$$

Solution

We should notice that $F(x) = f(g(x))$ so we can use the chain rule!

$$f(x) = \frac{x - 1}{x + 1} \quad \text{and} \quad g(x) = \sin x^2$$

$$f'(x) = \frac{2}{(x + 1)^2} \quad \text{and} \quad g'(x) = 2x \cos x^2$$

so

$$f'(x) = g'(x)f'(g(x))$$

Example

Question

Differentiate

$$F(x) = \frac{\sin x^2 - 1}{\sin x^2 + 1}$$

Solution

We should notice that $F(x) = f(g(x))$ so we can use the chain rule!

$$f(x) = \frac{x - 1}{x + 1} \quad \text{and} \quad g(x) = \sin x^2$$

$$f'(x) = \frac{2}{(x + 1)^2} \quad \text{and} \quad g'(x) = 2x \cos x^2$$

so

$$\begin{aligned} f'(x) &= g'(x)f'(g(x)) \\ &= 2x \cos x^2 \frac{2}{(\sin x^2 + 1)^2} \end{aligned}$$

Example

Question

A bacterial colony is estimated to have a population of P thousand individuals, where

$$P(t) = \frac{24t + 10}{t^2 + 1}$$

and t is the number of hours after a toxin is introduced.

Example

Question

A bacterial colony is estimated to have a population of P thousand individuals, where

$$P(t) = \frac{24t + 10}{t^2 + 1}$$

and t is the number of hours after a toxin is introduced.

1. At what rate is the population changing when $t = 1$?

Example

Question

A bacterial colony is estimated to have a population of P thousand individuals, where

$$P(t) = \frac{24t + 10}{t^2 + 1}$$

and t is the number of hours after a toxin is introduced.

1. At what rate is the population changing when $t = 1$?
2. Is the rate increasing or decreasing at this time?

Example

Question

A bacterial colony is estimated to have a population of P thousand individuals, where

$$P(t) = \frac{24t + 10}{t^2 + 1}$$

and t is the number of hours after a toxin is introduced.

1. At what rate is the population changing when $t = 1$?
2. Is the rate increasing or decreasing at this time?
3. At what time does the population begin to decrease?

Example (cont.)

1. At what rate is the population changing when $t = 1$?

We want to know how many individuals are being added per hour at $t = 1$. We need to find the derivative:

$$P(t) = (24t + 10)(t^2 + 1)^{-1}$$

Example (cont.)

1. At what rate is the population changing when $t = 1$?

We want to know how many individuals are being added per hour at $t = 1$. We need to find the derivative:

$$P(t) = (24t + 10)(t^2 + 1)^{-1} \quad \text{and} \quad \frac{d}{dx}(t^2 + 1)^{-1} = -2t(t^2 + 1)^{-2}$$

Example (cont.)

1. At what rate is the population changing when $t = 1$?

We want to know how many individuals are being added per hour at $t = 1$. We need to find the derivative:

$$P(t) = (24t + 10)(t^2 + 1)^{-1} \quad \text{and} \quad \frac{d}{dx}(t^2 + 1)^{-1} = -2t(t^2 + 1)^{-2}$$

$$P'(t) = 24(t^2 + 1)^{-1} - 2t(24t + 10)(t^2 + 1)^{-2}$$

Example (cont.)

1. At what rate is the population changing when $t = 1$?

We want to know how many individuals are being added per hour at $t = 1$. We need to find the derivative:

$$P(t) = (24t + 10)(t^2 + 1)^{-1} \quad \text{and} \quad \frac{d}{dx}(t^2 + 1)^{-1} = -2t(t^2 + 1)^{-2}$$

$$\begin{aligned} P'(t) &= 24(t^2 + 1)^{-1} - 2t(24t + 10)(t^2 + 1)^{-2} \\ &= -\frac{4(3t - 2)(2t + 3)}{(t^2 + 1)^2} \end{aligned}$$

Example (cont.)

1. At what rate is the population changing when $t = 1$?

We want to know how many individuals are being added per hour at $t = 1$. We need to find the derivative:

$$P(t) = (24t + 10)(t^2 + 1)^{-1} \quad \text{and} \quad \frac{d}{dx}(t^2 + 1)^{-1} = -2t(t^2 + 1)^{-2}$$

$$\begin{aligned} P'(t) &= 24(t^2 + 1)^{-1} - 2t(24t + 10)(t^2 + 1)^{-2} \\ &= -\frac{4(3t - 2)(2t + 3)}{(t^2 + 1)^2} \end{aligned}$$

$$P'(1) = -\frac{4 \cdot 1 \cdot 5}{4} = -5$$

Example (cont.)

1. Is the rate increasing or decreasing at this time?

We need to figure out when the derivative $P'(t)$ goes from being positive to negative. First let's ask when does

$$P'(t) = 0$$

Example (cont.)

1. Is the rate increasing or decreasing at this time?
2. At what time does the population begin to decrease?

We need to figure out when the derivative $P'(t)$ goes from being positive to negative. First let's ask when does

$$P'(t) = 0$$

Example (cont.)

1. Is the rate increasing or decreasing at this time?
2. At what time does the population begin to decrease?

We need to figure out when the derivative $P'(t)$ goes from being positive to negative. First let's ask when does

$$P'(t) = 0$$

Example (cont.)

1. Is the rate increasing or decreasing at this time?
2. At what time does the population begin to decrease?

We need to figure out when the derivative $P'(t)$ goes from being positive to negative. First let's ask when does

$$P'(t) = 0$$

$$-\frac{4(3t-2)(2t+3)}{(t^2+1)^2} = 0$$

Example (cont.)

1. Is the rate increasing or decreasing at this time?
2. At what time does the population begin to decrease?

We need to figure out when the derivative $P'(t)$ goes from being positive to negative. First let's ask when does

$$P'(t) = 0$$

$$-\frac{4(3t-2)(2t+3)}{(t^2+1)^2} = 0$$

$$t = \frac{2}{3} \quad \text{or} \quad -\frac{3}{2}$$