

# Math 3B: Lecture 14

Noah White

November 2, 2018

## Repeated factors

What if  $q(x)$  contains repeated factors? E.g. if  $q(x) = (x - 1)^2$  or  $q(x) = (x - 1)(x + 2)^3$ ?

## Repeated factors

What if  $q(x)$  contains repeated factors? E.g. if  $q(x) = (x - 1)^2$  or  $q(x) = (x - 1)(x + 2)^3$ ?

For every factor  $(ax + b)^k$  in  $q(x)$ , the partial fraction expansion has terms of the form

$$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \frac{A_3}{(ax + b)^3} + \cdots + \frac{A_k}{(ax + b)^k}.$$

## Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

## Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by  $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

## Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by  $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

Comparing coefficients

$$A = 1 \quad \text{and} \quad B - A = 0$$

## Example 1

$$\frac{x}{(x-1)^2} = \frac{A}{x-1} + \frac{B}{(x-1)^2}$$

Multiplying both sides by  $(x-1)^2$

$$\begin{aligned}x &= A(x-1) + B \\ &= Ax + (B-A)\end{aligned}$$

Comparing coefficients

$$A = 1 \quad \text{and} \quad B - A = 0$$

So

$$A = 1 \quad \text{and} \quad B = 1.$$

Side note: integrating  $\frac{1}{x}$ .

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$



Side note: integrating  $\frac{1}{x}$ .

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$

Side note: integrating  $\frac{1}{x}$ .

Recall that

Fact

$$\int \frac{1}{x} dx = \ln |x| + C$$

Using substitution this gives the formula

$$\int \frac{1}{ax + b} dx = \frac{1}{a} \ln |ax + b| + C.$$

Side note: integrating  $\frac{1}{x^k}$ .

Recall that if  $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Side note: integrating  $\frac{1}{x^k}$ .

Recall that if  $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Side note: integrating  $\frac{1}{x^k}$ .

Recall that if  $k > 1$

Fact

$$\int \frac{1}{x^k} dx = -\frac{1}{(k-1)x^{k-1}} + C$$

Using substitution this gives the formula

$$\int \frac{1}{(ax+b)^k} dx = -\frac{1}{a(k-1)(ax+c)^{k-1}} + C.$$

# Integrating rational functions $p(x)/q(x)$

Action plan

# Integrating rational functions $p(x)/q(x)$

## Action plan

1. Express  $\frac{p(x)}{q(x)}$  in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using polynomial long division.

# Integrating rational functions $p(x)/q(x)$

## Action plan

1. Express  $\frac{p(x)}{q(x)}$  in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using polynomial long division.

2. Write  $\frac{r(x)}{q(x)}$  as a sum of fractions of the form

$$\frac{A}{(ax + b)^k}$$

using partial fractions



# Integrating rational functions $p(x)/q(x)$

## Action plan

1. Express  $\frac{p(x)}{q(x)}$  in the form

$$d(x) + \frac{r(x)}{q(x)}$$

using **polynomial long division**.

2. Write  $\frac{r(x)}{q(x)}$  as a sum of fractions of the form

$$\frac{A}{(ax + b)^k}$$

using **partial fractions**

3. Integrate all these pieces separately.

## Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

## Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

### Solution

Using long division

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1}$$

## Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

### Solution

Using long division and partial fractions

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1} = x^2 - 2 + \frac{1}{2(x - 1)} - \frac{1}{2(x + 1)}$$

## Example 1

$$I = \int \frac{x^4 - 3x^2 + 3}{x^2 - 1} dx$$

### Solution

Using long division and partial fractions

$$\frac{x^4 - 3x^2 + 3}{x^2 - 1} = x^2 - 2 + \frac{1}{x^2 - 1} = x^2 - 2 + \frac{1}{2(x - 1)} - \frac{1}{2(x + 1)}$$

So

$$I = \frac{1}{3}x^3 - 2x + \frac{1}{2} \ln |x - 1| - \frac{1}{2} \ln |x + 1| + C.$$

## Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

## Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

### Solution

Using long division

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3}$$

## Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

### Solution

Using long division and partial fractions

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3} = 1 + \frac{1}{x-1} + \frac{3}{(x-1)^2} + \frac{3}{(x-1)^3}$$



## Example 2

$$I = \int \frac{x^3 - 2x^2 + 4x}{(x-1)^3} dx$$

### Solution

Using long division and partial fractions

$$\frac{x^3 - 2x^2 + 4x}{(x-1)^3} = 1 + \frac{x^2 + x + 1}{(x-1)^3} = 1 + \frac{1}{x-1} + \frac{3}{(x-1)^2} + \frac{3}{(x-1)^3}$$

So

$$I = x + \ln|x-1| - \frac{3}{x-1} - \frac{3}{2(x-1)^2} + C.$$

# Differential equations (motivation)

An (ordinary) **differential equation** (or **ODE**) is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

# Differential equations (motivation)

An (ordinary) **differential equation** (or **ODE**) is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

The challenge is to find all the functions  $y = f(x)$  (or even just one) that satisfy a given equation.

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

If it involves only first and second derivatives it is a **second order** differential equation. E.g.

$$y'' + y' = x$$

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

If it involves only first and second derivatives it is a **second order** differential equation. E.g.

$$y'' + y' = x$$

And so on.

# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$



# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$

We just need to find the antiderivative of  $f(x)$ .

# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$

We just need to find the antiderivative of  $f(x)$ .

## Note

The right hand side of the equation does not have any  $y$ 's.

## More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs
- classify the behaviour of many ODEs (e.g. does the solution go to zero or infinity?)



# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs
- classify the behaviour of many ODEs (e.g. does the solution go to zero or infinity?)
- understand how sensitive ODEs are to their parameters.

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".
- E.g.  $y(0) = 2$ .

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".
- E.g.  $y(0) = 2$ .
- Then we see that  $y(0) = 1 + C$ , so  $C = 1$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.



# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .
- If we want to draw the graph of  $y(t)$  then we look at  $g(0, 1)$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .
- If we want to draw the graph of  $y(t)$  then we look at  $g(0, 1)$ .
- If this is positive we go up, negative we go down!