

Math 3B: Lecture 7

Noah White

October 12, 2018

Midterm information

Midterm 1

- Practice midterms online

Midterm information

Midterm 1

- Practice midterms online
- arrive on time

Midterm information

Midterm 1

- Practice midterms online
- arrive on time
- calculators, cheat sheets

Midterm information

Midterm 1

- Practice midterms online
- arrive on time
- calculators, cheat sheets
- expected average, grades

Accumulated change

Often we encounter problems involving **accumulated change**.

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

Accumulated change

Often we encounter problems involving **accumulated change**.

Example

A rocket is accelerating at a rate of $a(t) = 0.3t^2$ metres per second squared. What is the rocket's velocity at $t = 30$?

Example

A population grows at a rate of $0.5P(t)$ people per year. How much does the population increase over 10 years?

These problems involve finding the area under some curve.

Example 1


If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

Example 1

If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

Solution

We model the car's speed using the function $s(t) = 30$. So we can see that the area under this curve



is the distance travelled (75 miles)

Example 2


If a car accelerates for 20 seconds at a rate of $2m/s^2$ and then decelerates for 30 seconds at a rate of $1m/s^2$, how far has it travelled?

Example 2

If a car accelerates for 20 seconds at a rate of 2m/s^2 and then decelerates for 30 seconds at a rate of 1m/s^2 , how far has it travelled?


Solution

The car's speed is given by $s(t) = 2t$ when $0 \leq t \leq 20$ and $s(t) = 60 - t$ when $20 \leq t \leq 50$. So the graph looks like




More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).

Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between $t = a$ and $t = b$?

Accumulated change

- Suppose we know the rate of change $f(t)$, of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between $t = a$ and $t = b$?
- Answer: area under $f(t)$ between a and b .

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

A first approach to calculating the area under a curve is to approximate using rectangles:

Areas under general curves

We would like to calculate the area between a function $f(x)$ and the x -axis, between $x = a$ and $x = b$.

A first approach to calculating the area under a curve is to approximate using rectangles:

(Too hard to draw, lets look at an animation)

The definite integral

Defintion

The definite integral of a function $f(x)$ is defined to be

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$

where $\Delta x = \frac{b-a}{n}$.

The fundamental theorem of calculus

Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

The fundamental theorem of calculus

Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

The fundamental theorem of calculus

Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

The fundamental theorem of calculus

Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .

The fundamental theorem of calculus

Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is, $F(x) = \int_a^x f(t) dt$ is an antiderivative of $f(x)$!

Note

- $F(x) = \int_a^x f(t) dt$ is a function of x .
- every input x produces a number as an output.

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

A consequence (corollary)

Corollary

For **any** antiderivative $F(x)$ of $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

Why?

Well $F(x) = \int_a^x f(t) \, dt + C$ for some a and C . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Example 1

Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

Solution

An antiderivative of $x^2 - 4$ is $\frac{1}{3}x^3 - 4x$ so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Example 2

Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

Solution

An antiderivative of $\sin x$ is $-\cos x$ so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$