

# Math 3B: Lecture 9

Noah White

January 30, 2019

# Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

# Differential equations (motivation)

A **differential equation** is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

The **challenge** is to find all the functions  $y = f(x)$  (or even just one) that satisfy a given equation.

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If  $h(t)$  measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

## Newton's second law (motivation)

The original differential equation!

$$F = ma$$

If  $h(t)$  measures the height of an object (maybe an apple?) above the earth then

$$a = h''(t)$$

The force due to gravity is roughly  $-10m$  Newtons, so

$$-10m = mh''(t)$$

## Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*



# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*

# Population growth (motivation)

A very rudimentary (but sometimes surprisingly accurate) way to model population growth is

## Assumption

*The rate of growth of a population is proportional to its current size*

If  $P(t)$  is the population at time  $t$ :

$$\frac{dP}{dt} = rP(t)$$

## Some more examples of differential equations

$$\frac{dy}{dx} = f(x)$$

$$\frac{dy}{dx} = y(1 - y)$$

$$y'' = \sqrt{a^2 - (y')^2}$$

$$\frac{dy}{dt} = k(A - y)^2$$

# Antiderivatives

We have been solving differential equations of the form

$$\frac{dy}{dx} = f(x).$$


Any antiderivative  $y = F(x)$  of  $f(x)$  is a solutions to this differential equation!

# Slope fields

In some cases it is impossible to find the antiderivative (without special functions). E.g.

$$f(x) = e^{-x^2}$$

But we can still (approximately) graph the antiderivative! First we draw the slope field



# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.

## Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?

# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$



# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$

# Slope fields (how to draw)


How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$
5. Do this for a grid of points on the  $xy$ -plane.


## Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




## Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




## Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



## Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



# Accumulated change

Often we encounter problems involving **accumulated change**.

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?



# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

These problems involve finding the area under some curve.

## Example 1


If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

## Example 1

If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

### Solution

We model the car's speed using the function  $s(t) = 30$ . So we can see that the area under this curve



is the distance travelled (75 miles)

## Example 2


If a car accelerates for 20 seconds at a rate of  $2m/s^2$  and then decelerates for 30 seconds at a rate of  $1m/s^2$ , how far has it travelled?

## Example 2

If a car accelerates for 20 seconds at a rate of  $2\text{m/s}^2$  and then decelerates for 30 seconds at a rate of  $1\text{m/s}^2$ , how far has it travelled?


### Solution

The car's speed is given by  $s(t) = 2t$  when  $0 \leq t \leq 20$  and  $s(t) = 60 - t$  when  $20 \leq t \leq 50$ . So the graph looks like



## More complicated rates of change


Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?





## More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



## Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).

# Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between  $t = a$  and  $t = b$ ?

# Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between  $t = a$  and  $t = b$ ?
- Answer: area under  $f(t)$  between  $a$  and  $b$ .

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

A first approach to calculating the area under a curve is to approximate using rectangles:

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

A first approach to calculating the area under a curve is to approximate using rectangles:

(Too hard to draw, lets look at an animation)

# The definite integral

## Defintion

The definite integral of a function  $f(x)$  is defined to be

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$

where  $\Delta x = \frac{b-a}{n}$ .




# The definite integral

## Defintion

The definite integral of a function  $f(x)$  is defined to be

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$


where  $\Delta x = \frac{b-a}{n}$ .



# Properties of definite integrals

## Zero area


$$\int_a^a f(x) \, dx = 0$$



# Properties of definite integrals

## Adding areas

$$\int_a^c f(x) \, dx = \int_a^b f(x) \, dx + \int_b^c f(x) \, dx$$



## More properties of definite integrals

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

# More properties of definite integrals

## Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

## Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

# More properties of definite integrals

Reversing the area

$$\int_a^b f(x) \, dx = - \int_b^a f(x) \, dx$$

Additivity

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$

Linearity (scalars factor out)

$$\int_a^b \alpha f(x) \, dx = \alpha \int_a^b f(x) \, dx$$

# The fundamental theorem of calculus

## Theorem

For any  $a$ ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

For any  $a$ ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$



# The fundamental theorem of calculus

## Theorem

For any  $a$ ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

# The fundamental theorem of calculus

## Theorem

For any  $a$ ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .

# The fundamental theorem of calculus

## Theorem

For any  $a$ ,

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .
- every input  $x$  produces a number as an output.

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Why?

Well  $F(x) = \int_a^x f(t) \, dt + C$  for some  $a$  and  $C$ . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

### Solution

An antiderivative of  $x^2 - 4$  is  $\frac{1}{3}x^3 - 4x$  so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$



## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

### Solution

An antiderivative of  $\sin x$  is  $-\cos x$  so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$

## Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that  $\frac{d}{dx} F(x) = f(x)$ .

## Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that  $\frac{d}{dx} F(x) = f(x)$ .

$$\frac{d}{dx} F(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

## Why is the FTC true?

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that  $\frac{d}{dx} F(x) = f(x)$ .

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \end{aligned}$$

## Why is the FTC true?


$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

We want to show that  $\frac{d}{dx} F(x) = f(x)$ .

$$\begin{aligned} \frac{d}{dx} F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) dt \end{aligned}$$


## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx$



## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx hf(x)$



## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$



## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} hf(x)\end{aligned}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x) \\&= f(x)\end{aligned}$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Why?

Well  $F(x) = \int_a^x f(t) \, dt + C$  for some  $a$  and  $C$ . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

# The indefinite integral

We also use the following notation for the general antiderivative:

**Definition**

$$\int f(x) \, dx := F(x) + C$$

# The indefinite integral

We also use the following notation for the general antiderivative:

## Definition

$$\int f(x) \, dx := F(x) + C$$

## Example

$$\int \sin(x) - x \, dx = -\cos(x) - \frac{1}{2}x^2 + C$$

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution



## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) \, dt$$

is an antiderivative for any choice of  $a$ .

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) dt$$

is an antiderivative for any choice of  $a$ .

- Lets use  $a = 0$ .

## Example

### Question

Find an antiderivative of  $f(x) = |x|$ ?

### Solution

- The FTC tells us that

$$F(x) = \int_a^x f(t) dt$$

is an antiderivative for any choice of  $a$ .


- Lets use  $a = 0$ .
- How should we calculate  $F(x)$ ?

## Example

Use the definition!

$$F(x) = \int_0^x |t| \, dt$$

is the area under  $|t|$ !




$$F(x) = \frac{1}{2}x^2 \quad \text{if } x \geq 0$$

## Example

If  $x \leq 0$  then

$$F(x) = \int_0^x |t| \, dt = - \int_x^0 |t| \, dt$$

is the negative of the area under  $|t|$ !



$$F(x) = -\frac{1}{2}x^2 \quad \text{if } x \leq 0$$

## Example

In summary

$$F(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } x \geq 0 \\ -\frac{1}{2}x^2 & \text{if } x \leq 0 \end{cases}$$

or

$$F(x) = \frac{1}{2}x|x|$$