

Math 3B: Lecture 5

Noah White

January 16, 2019

Last time

Last time, we spoke about

- global and local extrema

Last time

Last time, we spoke about

- global and local extrema
- critical points

Last time

Last time, we spoke about

- global and local extrema
- critical points
- how to find local extrema

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.
2. Classify each critical point as local max or min using

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.
2. Classify each critical point as local max or min using
 - 2.1 First derivative test

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.
2. Classify each critical point as local max or min using
 - 2.1 First derivative test
 - 2.2 Second derivative test

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.
2. Classify each critical point as local max or min using
 - 2.1 First derivative test
 - 2.2 Second derivative test
3. Find the global max (if one exists) using

Recipe for classifying maxima and minima

We have a function $f : D \longrightarrow R$. How do we find all local/global extrema?

1. Find all critical points.
2. Classify each critical point as local max or min using
 - 2.1 First derivative test
 - 2.2 Second derivative test
3. Find the global max (if one exists) using
 - 3.1 ???

Closed interval method

To find the global extrema of $f(x)$ defined on a closed interval $[a, b]$:

1. Find all critical points.

Closed interval method

To find the global extrema of $f(x)$ defined on a closed interval $[a, b]$:

1. Find all critical points.
2. Evaluate $f(x)$ at all the critical points and at a and b

Closed interval method

To find the global extrema of $f(x)$ defined on a closed interval $[a, b]$:

1. Find all critical points.
2. Evaluate $f(x)$ at all the critical points and at a and b
3. The smallest value is the global min

Closed interval method

To find the global extrema of $f(x)$ defined on a closed interval $[a, b]$:

1. Find all critical points.
2. Evaluate $f(x)$ at all the critical points and at a and b
3. The smallest value is the global min
4. The largest value is the global max

Open interval method

To find the global extrema of $f(x)$ defined on a **open** interval (a, b) :

Note: a could be $-\infty$ and b could be ∞ .

1. Find all critical points.

Open interval method

To find the global extrema of $f(x)$ defined on a **open** interval (a, b) :

Note: a could be $-\infty$ and b could be ∞ .

1. Find all critical points.
2. Find the limits

$$L = \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad M = \lim_{x \rightarrow b^-} f(x)$$

Open interval method

To find the global extrema of $f(x)$ defined on a **open** interval (a, b) :

Note: a could be $-\infty$ and b could be ∞ .

1. Find all critical points.
2. Find the limits

$$L = \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad M = \lim_{x \rightarrow b^-} f(x)$$

3. Evaluate $f(x)$ at all the critical points

Open interval method

To find the global extrema of $f(x)$ defined on a **open** interval (a, b) :

Note: a could be $-\infty$ and b could be ∞ .

1. Find all critical points.
2. Find the limits

$$L = \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad M = \lim_{x \rightarrow b^-} f(x)$$

3. Evaluate $f(x)$ at all the critical points
4. The smallest value is the global min unless L or M are smaller, in which case there is no global min

Open interval method

To find the global extrema of $f(x)$ defined on a **open** interval (a, b) :

Note: a could be $-\infty$ and b could be ∞ .

1. Find all critical points.
2. Find the limits

$$L = \lim_{x \rightarrow a^+} f(x) \quad \text{and} \quad M = \lim_{x \rightarrow b^-} f(x)$$

3. Evaluate $f(x)$ at all the critical points
4. The smallest value is the global min unless L or M are smaller, in which case there is no global min
5. The largest value is the global max unless L or M are larger, in which case there is no global max

Example

$f(x)$ defined on $(-\infty, \infty)$ with

$$L = \lim_{x \rightarrow -\infty} f(x) = M = \lim_{x \rightarrow \infty} f(x) = 0$$

