

Math 3B: Lecture 18

Noah White

November 13, 2017

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

$$\frac{dy}{dx} = e^{\sin x} \cos x$$

Checking solutions

The most straightforward way of checking a function $y = f(x)$ is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

Example

The function $y = e^{\sin x}$ is a solution of $\frac{dy}{dx} = y \cos x$. To check note that

$$\begin{aligned}\frac{dy}{dx} &= e^{\sin x} \cos x \\ y \cos x &= e^{\sin x} \cos x\end{aligned}$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the y 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

4. solve for y !

Examples

On the board...

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Examples

$$\frac{dy}{dt} = ay, \quad \frac{dy}{dt} = -\lambda y.$$

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream

Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream
- pollutant in water supply

General solution

Using separation of variables, we can show that the general solution to

$$\frac{dy}{dt} = a - by$$

is

$$y(t) = \frac{a}{b} - Ce^{-bt}$$

where C is an arbitrary constant.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = M e^{-0.5t \ln(2)} \text{ mg left}$$

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = Me^{-0.5t \ln(2)} \text{ mg left}$$

- Thus the rate at which the drug is leaving (at time t) is given by

$$0.5 \ln(2) Me^{-0.5t \ln(2)} = 0.5 \ln(2)(\text{current concentration}) \text{ mg/h.}$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

- Thus at time t the concentration is

$$y(t) = 28.9 - 28.9e^{-0.3t} = 28.9(1 - e^{-0.3t})$$

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

$$\frac{dT}{dt} = k(A - T)$$

General solution

$$T(t) = A - Ce^{-kt}.$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.
- Thus

$$90 = 70 - C \quad \text{so} \quad C = -20.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

- Rearranging we get $20e^{-0.01t} = 5$ i.e.

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

- So we get

$$t = -100 \ln\left(\frac{1}{4}\right) \approx 138 = 2 \text{ hours } 18 \text{ minutes.}$$