

Math 3B: Lecture 16

Noah White

February 17, 2017

Last time

- Checking solutions to differential equations.

Last time

- Checking solutions to differential equations.
- Implicit differentiation.

Last time

- Checking solutions to differential equations.
- Implicit differentiation.
- Separation of variables.

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Linear models

Definition

A first order ODE is **linear** if it is of the form

$$\frac{dy}{dt} = a + by$$

for constants a and b . In other words, the right hand side is a linear function of y .

Examples

$$\frac{dy}{dt} = ay, \quad \frac{dy}{dt} = -\lambda y.$$

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

Mixing models

A mixing model describes the concentration of something over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream

Mixing models

A mixing model describes the concentration of **something** over time, if

- rate in is constant
- rate out is proportional to concentration

so

$$\begin{aligned}\frac{dy}{dt} &= \text{rate in} - \text{rate out} \\ &= a - by\end{aligned}$$

Note

Something could mean (for example)

- concentration of a drug in bloodstream
- pollutant in water supply

General solution

Using separation of variables, we can show that the general solution to

$$\frac{dy}{dt} = a - by$$

is

$$y(t) = \frac{a}{b} - Ce^{-bt}$$

where C is an arbitrary constant.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an infusion rate of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = M e^{-0.5t \ln(2)} \text{ mg left}$$

Example 1

A drug with a half-life of 2 hours is injected into the bloodstream with an **infusion rate** of 10 mg/h. Determine the concentration $y(t)$ at time t .

Solution

- Ignoring infusion, every 2 hours the amount of drug halves.
- Starting with M mg, after t hours there will be

$$M \left(\frac{1}{2} \right)^{t/2} = Me^{-0.5t \ln(2)} \text{ mg left}$$

- Thus the rate at which the drug is leaving (at time t) is given by

$$0.5 \ln(2) Me^{-0.5t \ln(2)} = 0.5 \ln(2)(\text{current concentration}) \text{ mg/h.}$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

Example 1

- If we infuse the drug at a rate of 10 mg/h we have

$$\frac{dy}{dt} = 10 - 0.5 \ln(2)y$$

- The general solution to this is

$$y(t) = \frac{10}{0.5 \ln(2)} - Ce^{-0.5 \ln(2)t}.$$

- Since there was initially no drug in the bloodstream, $y(0) = 0$,

$$0 = \frac{20}{\ln(2)} - C \approx 28.9 - C$$

- Thus at time t the concentration is

$$y(t) = 28.9 - 28.9e^{-0.3t} = 28.9(1 - e^{-0.3t})$$

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

Newton's Law of Cooling

Isaac Newton stated that

Newton's Law of Cooling

The temperature T of a body changes at a rate proportional to the difference between the ambient temperature A and T .

$$\frac{dT}{dt} = k(A - T)$$

General solution

$$T(t) = A - Ce^{-kt}.$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.

Example 2

An object takes 20 minutes to cool from 90° to 86° in a room which is 70° . At what time will it be 75° ?

Solution

- The temp is described by the equation

$$\frac{dT}{dt} = k(70 - T).$$

- The solution is given by

$$T(t) = 70 - Ce^{-kt}$$

- We know $T(0) = 90$ and $T(20) = 86$.
- Thus

$$90 = 70 - C \quad \text{so} \quad C = -20.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln \left(\frac{4}{5} \right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

Example 2

- $T(t) = 70 + 20e^{-kt}$
- To find k we use $T(20) = 86$

$$86 = 70 + 20e^{-20k}$$

- Thus

$$e^{-20k} = \frac{86 - 70}{20} = \frac{4}{5} \quad \text{so} \quad k = -\frac{1}{20} \ln\left(\frac{4}{5}\right) \approx -0.01.$$

- The model is thus $T(t) = 70 + 20e^{-0.01t}$. We want to solve

$$75 = 70 + 20e^{-0.01t}.$$

- Rearranging we get $20e^{-0.01t} = 5$ i.e.

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

Example 2

- $20e^{-0.01t} = 5$ becomes

$$e^{-0.01t} = \frac{1}{4}$$

- Applying a logarithm

$$-0.01t = \ln\left(\frac{1}{4}\right)$$

- So we get

$$t = -100 \ln\left(\frac{1}{4}\right) \approx 138 = 2 \text{ hours } 18 \text{ minutes.}$$

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

$$\frac{dM}{dt} = aS - bM.$$

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

$$\frac{dM}{dt} = aS - bM.$$

We can make an additional assumption

- The organism is (roughly) a cube of length L cm.

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

$$\frac{dM}{dt} = aS - bM.$$

We can make an additional assumption

- The organism is (roughly) a cube of length L cm.
- This means the mass $M = L^3$ g (if we assume the organism is mostly water).

Von Bertalanffy growth model

Ludwig von Bertalanffy estimated the growth of an organism by assuming that it

- ingests food at a rate proportional to its surface area
- respire proportional to its mass

$$\frac{dM}{dt} = aS - bM.$$

We can make an additional assumption

- The organism is (roughly) a cube of length L cm.
- This means the mass $M = L^3$ g (if we assume the organism is mostly water).
- Surface area is $S = 6L^2$

Von Bertalanffy growth model

- We can differentiate the relationship $M = L^3$

$$\frac{dM}{dt} = 3L^2 \frac{dL}{dt}$$

Von Bertalanffy growth model

- We can differentiate the relationship $M = L^3$

$$\frac{dM}{dt} = 3L^2 \frac{dL}{dt}$$

- Combining this with

$$\frac{dM}{dt} = 6aL^2 - bL^3$$

Von Bertalanffy growth model

- We can differentiate the relationship $M = L^3$

$$\frac{dM}{dt} = 3L^2 \frac{dL}{dt}$$

- Combining this with

$$\frac{dM}{dt} = 6aL^2 - bL^3$$

- We get

$$3L^2 \frac{dL}{dt} = 6aL^2 - bL^3$$

Von Bertalanffy growth model

- Dividing by $3L^2$ gives

$$\frac{dL}{dt} = \frac{6a}{3} - \frac{b}{3}L = \frac{b}{3} \left(\frac{6a}{b} - L \right)$$

Von Bertalanffy growth model

- Dividing by $3L^2$ gives

$$\frac{dL}{dt} = \frac{6a}{3} - \frac{b}{3}L = \frac{b}{3} \left(\frac{6a}{b} - L \right)$$

Von Bertalanffy growth model

The growth of an organism is governed by

$$\frac{dL}{dt} = k(L_{\infty} - L)$$

where k and L_{∞} are constants.