

# Math 3B: Lecture 16

Noah White

February 15, 2019

# Differential equations (motivation)

An (ordinary) **differential equation** (or **ODE**) is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

# Differential equations (motivation)

An (ordinary) **differential equation** (or **ODE**) is an equation that involves derivatives of an unknown function.

$$\frac{d^2y}{dx^2} = y - 3y^2$$

or

$$x^2y'' + xy' + x^2y = 0$$

The challenge is to find all the functions  $y = f(x)$  (or even just one) that satisfy a given equation.

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

If it involves only first and second derivatives it is a **second order** differential equation. E.g.

$$y'' + y' = x$$

## First, second, ... order

If a differential equation involves only first derivatives it is a **first order** differential equation. E.g.

$$\frac{dy}{dx} = 2y$$

If it involves only first and second derivatives it is a **second order** differential equation. E.g.

$$y'' + y' = x$$

And so on.

# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$



# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$

We just need to find the antiderivative of  $f(x)$ .

# Integration

We already know how to solve ODEs of the form

$$\frac{dy}{dx} = f(x)$$

We just need to find the antiderivative of  $f(x)$ .

## Note

The right hand side of the equation does not have any  $y$ 's.

## More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs

# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs
- classify the behaviour of many ODEs (e.g. does the solution go to zero or infinity?)



# More types of differential equations

In this course we focus on first order ODEs. You will be able to solve equations like

- $\frac{dy}{dx} = -3y + 5$
- $\frac{dy}{dx} = f(x)g(y)$
- $y' = x^{-5} \sin y$

And you'll be able to

- draw solutions for many other ODEs
- classify the behaviour of many ODEs (e.g. does the solution go to zero or infinity?)
- understand how sensitive ODEs are to their parameters.

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".
- E.g.  $y(0) = 2$ .

# Initial value problems

- If we try to solve the differential equation

$$\frac{dy}{dt} = 3t^2 - \sin t$$

we get (by integrating)

$$y(t) = t^3 + \cos t + C.$$

- This doesn't tell us exactly what  $y(t)$  is, we still need to know what  $C$  is!
- The extra piece of data we need is called an "initial value".
- E.g.  $y(0) = 2$ .
- Then we see that  $y(0) = 1 + C$ , so  $C = 1$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.



# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .
- If we want to draw the graph of  $y(t)$  then we look at  $g(0, 1)$ .

# How to think about differential equations

- Suppose you are given a differential equation, and an initial value:

$$\frac{dy}{dt} = g(t, y) \quad y(0) = 1$$

- We think of  $t$  as time and  $y$  as describing how some quantity changes over time.
- We think about  $y(t)$  "evolving" over time.
- Imagine a point starting at  $(t = 0, y = 1)$ .
- If we want to draw the graph of  $y(t)$  then we look at  $g(0, 1)$ .
- If this is positive we go up, negative we go down!

# Modelling using differential equations

## Question

Why are differential equations important?

# Modelling using differential equations

## Question

Why are differential equations important?

## Answer

They describe nature!

# Modelling using differential equations

## Question

Why are differential equations important?

## Answer

They describe nature!

# Modelling using differential equations

## Question

Why are differential equations important?

## Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function  $y(t)$  that describes something we are interested in (e.g. population/mass/etc)



# Modelling using differential equations

## Question

Why are differential equations important?

## Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function  $y(t)$  that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)

# Modelling using differential equations

## Question

Why are differential equations important?

## Answer

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function  $y(t)$  that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)
- We can't do this directly, but we can write down an ODE that  $y$  satisfies instead.

## Example 1

### The senario

Want to find a function  $N(t)$ , descibing the size of a population at time  $t$  (in years). Assume

## Example 1

### The senario

Want to find a function  $N(t)$ , descibing the size of a population at time  $t$  (in years). Assume

- The population is 100 people when  $t = 0$

## Example 1

### The senario

Want to find a function  $N(t)$ , descibing the size of a population at time  $t$  (in years). Assume

- The population is 100 people when  $t = 0$
- No immigration or emmigration

## Example 1

### The senario

Want to find a function  $N(t)$ , descibing the size of a population at time  $t$  (in years). Assume

- The population is 100 people when  $t = 0$
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

$bN(t)$  births per year, for some  $b$

## Example 1

### The senario

Want to find a function  $N(t)$ , descibing the size of a population at time  $t$  (in years). Assume

- The population is 100 people when  $t = 0$
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

$bN(t)$  births per year, for some  $b$

- Number of deaths is proportional to the total number of people. So

$dN(t)$  deaths per year, for some  $d$

## Example 1

The total change in population at time  $t$  is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$



## Example 1

The total change in population at time  $t$  is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$

In real life we would determine  $b$  and  $d$  experimentally.

## Example 1

The total change in population at time  $t$  is

$$\begin{aligned}\frac{dN}{dt} &= \text{births at } t - \text{deaths at } t \\ &= bN(t) - dN(t) \\ &= (b - d)N(t).\end{aligned}$$

In real life we would determine  $b$  and  $d$  experimentally. Let  $r = b - d$ . the **instinsic growth rate**. So our model is

$$\frac{dN}{dt} = rN.$$

and we know  $N(0) = 100$ .

## Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

### Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so  $N(t) = 100$  for all  $t$ ).

## Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

### Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so  $N(t) = 100$  for all  $t$ ).

### Case 2: $r > 0$

The population is increasing indefinitely.

# Behaviour of solutions

$$\frac{dN}{dt} = rN.$$

## Case 1: $r = 0$

The population never grows or shrinks, it always stays the same (so  $N(t) = 100$  for all  $t$ ).

## Case 2: $r > 0$

The population is increasing indefinitely.

## Case 3: $r < 0$

The population is decreasing indefinitely.

# Solution to a simple ODE

## Theorem

For any constant  $a$ , if  $y$  is a solution to the ODE

$$\frac{dy}{dx} = ay$$

then  $y$  is given by

$$y = Ce^{ax}$$

for some constant  $C$ .

# Solution to a simple ODE

## Theorem

For any constant  $a$ , if  $y$  is a solution to the ODE

$$\frac{dy}{dx} = ay$$

then  $y$  is given by

$$y = Ce^{ax}$$

for some constant  $C$ .

## Next time

We will see why, but for now we can verify it is actually a solution:

$$\frac{dy}{dx} = \frac{d}{dx} Ce^{ax} = C \frac{d}{dx} e^{ax} = Cae^{ax} = ay.$$

## Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$



## Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant  $C$ .

## Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant  $C$ . To determine  $C$ , we need one extra piece of information,  $N(0) = 100$ .

## Back to example 1

We know our population model was governed by

$$\frac{dN}{dt} = (b - d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant  $C$ . To determine  $C$ , we need one extra piece of information,  $N(0) = 100$ .

$$100 = Ce^0 \quad \text{so} \quad C = 100.$$

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d \propto N(t)).$$

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d + kN(t)) N(t).$$

- This means

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d + kN(t)) N(t).$$

- This means



## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d + kN(t)) N(t).$$

- This means

$$\frac{dN}{dt} = bN - (d + kN)N$$

Where  $K = r/k$ .

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d + kN(t)) N(t).$$

- This means

$$\begin{aligned}\frac{dN}{dt} &= bN - (d + kN)N \\ &= (b - d - kN)N = (r - kN)N\end{aligned}$$

Where  $K = r/k$ .

## Logistic growth

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constraints, e.g. amount of space, food, etc.

- When the population becomes large, the death rate should increase!
- Let's assume this is linear, i.e. number of deaths at time  $t$

$$(d + kN(t)) N(t).$$

- This means

$$\begin{aligned}\frac{dN}{dt} &= bN - (d + kN)N \\ &= (b - d - kN)N = (r - kN)N \\ &= r \left(1 - \frac{kN}{r}\right) N = r \left(1 - \frac{N}{K}\right) N\end{aligned}$$

Where  $K = r/k$ .

# Logistic growth

The equation

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

is called the **Logistic equation** and  $K$  is the **carrying capacity**.

## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 1.  $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

# Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 1.  $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

Case 2.  $N(0) = K$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains  $K$ .

# Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 1.  $N(0) = 0$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains 0.

Case 2.  $N(0) = K$

In this case the growth rate is 0 initially, so  $N(t)$  does not increase or decrease, so remains  $K$ .

Key takeaway

Both  $N(t) = 0$  and  $N(t) = K$  are solutions to the ODE. They are called **equilibrium solutions**.

## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 3.  $0 \leq N(0) \leq K$

In this case,  $N$  is initially increasing and so becomes more positive, slowing down as it gets close to  $K$ .



## Behaviour of logistic growth

Assume that  $r > 0$  and  $K > 0$ .

$$\frac{dN}{dt} = r \left( 1 - \frac{N}{K} \right) N$$

Case 3.  $0 \leq N(0) \leq K$

In this case,  $N$  is initially increasing and so becomes more positive, slowing down as it gets close to  $K$ .

Case 4.  $N(0) \geq K$

In this case  $N$  is initially decreasing but decreases slower and slower as it gets close to  $K$ .

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

$$\frac{dy}{dx} = e^{\sin x} \cos x$$

## Checking solutions

The most straightforward way of checking a function  $y = f(x)$  is a solution to a differential equation

$$\frac{dy}{dx} = g(x, y)$$

is to simply plug it in to both sides.

### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

$$\begin{aligned}\frac{dy}{dx} &= e^{\sin x} \cos x \\ y \cos x &= e^{\sin x} \cos x\end{aligned}$$

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\frac{d}{dx}(x^2 + y^2) = \frac{d}{dx}1$$



## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \\ \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) &= 0\end{aligned}$$

## Reminder: Implicit differentiation

If we have an equation relating variables  $y$  and  $x$ , e.g.

$$x^2 + y^2 = 1$$

we can **differentiate implicitly** by applying  $\frac{d}{dx}$  to both side.

### Example

In the above example we get

$$\begin{aligned}\frac{d}{dx}(x^2 + y^2) &= \frac{d}{dx}1 \\ \frac{d}{dx}(x^2) + \frac{d}{dx}(y^2) &= 0 \\ 2x + 2y'y &= 0\end{aligned}$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$

$$3 - y' \sin y = y + xy'.$$

## Another example

Lets differentiate

$$3x + \cos y = xy$$

To do this we apply  $\frac{d}{dx}$  to both sides:

$$\frac{d}{dx}(3x + \cos y) = \frac{d}{dx}(xy)$$

$$\frac{d}{dx}(3x) + \frac{d}{dx}(\cos y) = y \frac{d}{dx}(x) + x \frac{d}{dx}(y)$$

$$3 - y' \sin y = y + xy'.$$

### Note

We can rearrange this to get

$$y' = \frac{3 - y}{x + \sin y}$$

a differential equation. Whatever  $y$  is, as long as it obeys the above relation, it is a solution to this ODE!

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$



## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

## Separation of variables

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{dy}{dx} = f(y)g(x)$$

It works by

1. moving all the  $y$ 's over to the left:

$$\frac{1}{f(y)} \frac{dy}{dx} = g(x)$$

2. integrating both sides (with respect to  $x$ ):

$$\int \frac{1}{f(y)} \frac{dy}{dx} dx = \int g(x) dx.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} dy = \int g(x) dx.$$

4. solve for  $y$ !

## Examples

On the board...