

# Math 3B: Lecture 3

Noah White

September 28, 2016

## Last time

Last time, we spoke about

- Graphing using calculus

## Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes

## Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes

## Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

## Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative

# Last time

Last time, we spoke about

- Graphing using calculus
- Horizontal asymptotes
- Vertical asymptotes
- Role of the first/second derivative


**Note:** The quiz will start at the beginning of the discussion section next time.

## Example time

... On the board.



# Slanted asymptotes



# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote  $y = mx + b$ .

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote  $y = mx + b$ .
- To find  $b$ :

# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote  $y = mx + b$ .
- To find  $b$ :



# Slanted asymptotes

- An asymptote is a straight line which the function approaches as  $x \rightarrow \pm\infty$
- If a function has a slanted asymptote its gradient must approach a constant
- So we should find

$$\lim_{x \rightarrow \pm\infty} f'(x) = m$$

- We then know the function has a slanted asymptote  $y = mx + b$ .
- To find  $b$ :

$$b = \lim_{x \rightarrow \pm\infty} (f(x) - mx)$$

## Example time

... On the board.

# Functions

A function is three pieces of information

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

are all different functions!



# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$

are all different functions!

# Functions

A function is three pieces of information

- A domain,  $D \subset \mathbb{R}$
- A range,  $R \subset \mathbb{R}$ , and
- A rule  $f : D \longrightarrow R$  that assigns to every element of  $D$  an element of  $R$ .

## Example

The functions

- $f : \mathbb{R} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}; x \mapsto x^2$
- $f : \mathbb{R} \longrightarrow \mathbb{R}_{\geq 0}; x \mapsto x^2$

are all different functions!

# Global Maximums and minimums

## Definition (Global maximum)

A function  $f : D \longrightarrow R$  has a global maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$

# Global Maximums and minimums

## Definition (Global maximum)

A function  $f : D \longrightarrow R$  has a global maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \in D$$


## Definition (Global minimum)

A function  $f : D \longrightarrow R$  has a global minimum at  $a$  if

$$f(x) \geq f(a) \quad \text{for all } x \in D$$


## Example of a global minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$  has a min at  $x = 0$



## Example of a global maximum

$f : (-\infty, 0] \rightarrow \mathbb{R}; f(x) = x^3$  has a max at  $x = 0$



# Local Maximums and minimums

## Definition (local maximum)

A function  $f : D \longrightarrow R$  has a local maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$



# Local Maximums and minimums

## Definition (local maximum)

A function  $f : D \longrightarrow R$  has a local maximum at  $a$  if

$$f(x) \leq f(a) \quad \text{for all } x \text{ near } a$$


## Definition (local minimum)

A function  $f : D \longrightarrow R$  has a local minimum at  $a$  if

$$f(x) \geq f(a) \quad \text{for all } x \text{ near } a$$


## Example of a local minimum

$f : \mathbb{R} \rightarrow \mathbb{R}; x \mapsto x^2$  has a min at  $x = 0$




## Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$  has a local max at  $x = -\frac{1}{3}$



## Example of a local maximum

$f : \mathbb{R} \rightarrow \mathbb{R}; f(x) = x^3 - 4x^2 - 3x + 13$  has a local max at  $x = -\frac{1}{3}$



# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )

# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )
- $f(x) = \sin x$  has a critical point at  $x = \frac{\pi}{2}$  (since  $f'(x) = \cos x$ )



# Critical point

## Definition (Critical point)

A function  $f(x)$  has a critical point at  $x = a$  if  $f'(a) = 0$  or if  $f'(a)$  is undefined.

## Examples

- $f(x) = x^2$  has a critical point at  $x = 0$  (since  $f'(x) = 2x$ )
- $f(x) = \sin x$  has a critical point at  $x = \frac{\pi}{2}$  (since  $f'(x) = \cos x$ )
- $f(x) = e^x$  doesn't have any critical points since  $f'(x) = e^x$  can never be zero

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

**Note:** All extrema are critical points, but not all critical points are extrema!

# How to find minimums and maximums

Local maximums and minimums (extrema) occur at

- critical points
- end points of the domain (are also critical points!)

**Note:** All extrema are critical points, but not all critical points are extrema!


## Example

$f : (-\infty, 1] \rightarrow \mathbb{R}; f(x) = x^3$  has critical points at

$$x = 0 \text{ and } 1$$

## Example

$f'(x) = 3x^2$  so  $f'(0) = 0$  and  $f'(1)$  is undefined.





## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

First derivative test (minimums)

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)


- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (minimums)

- If  $f'(x) < 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) > 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a minimum at  $a$ .



## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

First derivative test (maximums)

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and



# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .

## First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

### First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .

# First derivative test

Suppose  $x = a$  is a critical point for the function  $f(x)$ .

## First derivative test (maximums)

- If  $f'(x) > 0$  for  $x$  less than and close to  $a$ , and
- $f'(x) < 0$  for  $x$  greater than and close to  $a$ , then
- $f(x)$  has a maximum at  $a$ .

