## Math 3B: Lecture 16

Noah White

November 5, 2018

### Question

Why are differential equations important?

### Question

Why are differential equations important?

### **Answer**

They describe nature!

### Question

Why are differential equations important?

### **Answer**

They describe nature!

### Question

Why are differential equations important?

#### **Answer**

They describe nature!

Today we will see an number of real world situations

• The goal is to write down a function y(t) that describes something we are interested in (e.g. population/mass/etc)

### Question

Why are differential equations important?

### **Answer**

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function y(t) that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)

### Question

Why are differential equations important?

#### **Answer**

They describe nature!

Today we will see an number of real world situations

- The goal is to write down a function y(t) that describes something we are interested in (e.g. population/mass/etc)
- as some other variable changes (usually time)
- We can't do this directly, but we can write down an ODE that y satisfies instead.

### The senario

Want to find a function N(t), descibing the size of a population at time t (in years). Assume

#### The senario

Want to find a function N(t), descibing the size of a population at time t (in years). Assume

• The population is 100 people when t=0

### The senario

Want to find a function N(t), descibing the size of a population at time t (in years). Assume

- The population is 100 people when t=0
- No immigration or emmigration

### The senario

Want to find a function N(t), descibing the size of a population at time t (in years). Assume

- The population is 100 people when t = 0
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

bN(t) births per year, for some b

#### The senario

Want to find a function N(t), descibing the size of a population at time t (in years). Assume

- The population is 100 people when t = 0
- No immigration or emmigration
- Number of births is proportional to the total number of people. So

bN(t) births per year, for some b

 Number of deaths is proportional to the total number of people. So

dN(t) deaths per year, for some d

The total change in population at time t is

$$\frac{\mathrm{d}N}{\mathrm{d}t} = \text{ births at } t - \text{ deaths at } t$$
$$= bN(t) - dN(t)$$
$$= (b - d)N(t).$$

The total change in population at time t is

$$\frac{\mathrm{d}N}{\mathrm{d}t} = \text{ births at } t - \text{ deaths at } t$$
$$= bN(t) - dN(t)$$
$$= (b - d)N(t).$$

In real life we would determine b and d experimentally.

The total change in population at time t is

$$\frac{\mathrm{d}N}{\mathrm{d}t} = \text{ births at } t - \text{ deaths at } t$$
$$= bN(t) - dN(t)$$
$$= (b - d)N(t).$$

In real life we would determine b and d experimentally. Let r=b-d. the instinsic growth rate. So our model is

$$\frac{\mathrm{d}N}{\mathrm{d}t}=rN.$$

and we know N(0) = 100.

### Behaviour of solutions

$$\frac{\mathrm{d}N}{\mathrm{d}t} = rN.$$

### Case 1: r = 0

The population never grows or shrinks, it always stays the same (so N(t) = 100 for all t).

### Behaviour of solutions

$$\frac{\mathrm{d}N}{\mathrm{d}t} = rN.$$

### Case 1: r = 0

The population never grows or shrinks, it always stays the same (so N(t) = 100 for all t).

### Case 2: r > 0

The population is increasing indefinitely.

### Behaviour of solutions

$$\frac{\mathrm{d}N}{\mathrm{d}t} = rN.$$

### Case 1: r = 0

The population never grows or shrinks, it always stays the same (so N(t) = 100 for all t).

Case 2: r > 0

The population is increasing indefinitely.

Case 3: r < 0

The population is decreasing indefinitely.

# Solution to a simple ODE

### **Theorem**

For any constant a, if y is a solution to the ODE

$$\frac{\mathrm{d}y}{\mathrm{d}x} = ay$$

then y is given by

$$y = Ce^{ax}$$

for some constant C.

## Solution to a simple ODE

#### Theorem

For any constant a, if y is a solution to the ODE

$$\frac{\mathrm{d}y}{\mathrm{d}x} = ay$$

then y is given by

$$y = Ce^{ax}$$

for some constant C.

#### Next time

We will see why, but for now we can verify it is actually a solution:

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{d}}{\mathrm{d}x}Ce^{ax} = C\frac{\mathrm{d}}{\mathrm{d}x}e^{a}x = Cae^{ax} = ay.$$

We know our population model was governed by

$$\frac{\mathrm{d}N}{\mathrm{d}t}=(b-d)N.$$

We know our population model was governed by

$$\frac{\mathrm{d}N}{\mathrm{d}t}=(b-d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C.

We know our population model was governed by

$$\frac{\mathrm{d}N}{\mathrm{d}t}=(b-d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C. To detirmine C, we need one extra piece of information, N(0)=100.

We know our population model was governed by

$$\frac{\mathrm{d}N}{\mathrm{d}t}=(b-d)N.$$

So the general solution is

$$N(t) = Ce^{(b-d)t}$$

for some constant C. To detirmine C, we need one extra piece of information, N(0)=100.

$$100 = Ce^0$$
 so  $C = 100$ .

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

 When the popluation becomes large, the death rate should increase!

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

$$(d+kN(t))N(t).$$

This means

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

$$(d+kN(t))N(t).$$

This means

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

$$(d+kN(t))N(t).$$

This means

$$\frac{\mathrm{d}N}{\mathrm{d}t} = bN - (d + kN)N$$

Where K = r/k.

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

$$(d+kN(t))N(t).$$

This means

$$\frac{\mathrm{d}N}{\mathrm{d}t} = bN - (d + kN)N$$
$$= (b - d - kN)N = (r - kN)N$$

Where K = r/k.

The previous example is a good first approximation but it is not very realistic in the long term. Usually there are constrains, e.g amount of space, food, etc.

- When the popluation becomes large, the death rate should increase!
- Lets assume this is linear, i.e. number of deaths at time t

$$(d+kN(t))N(t).$$

This means

$$\frac{\mathrm{d}N}{\mathrm{d}t} = bN - (d + kN)N$$

$$= (b - d - kN)N = (r - kN)N$$

$$= r\left(1 - \frac{kN}{r}\right)N = r\left(1 - \frac{N}{K}\right)N$$

Where K = r/k.

The equation

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

is called the Logistic equation and K is the carrying capacity.

Assume that r > 0 and K > 0.

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

Case 1. 
$$N(0) = 0$$

In this case the growth rate is 0 initially, so N(t) does not increase or decrease, so remains 0.

Assume that r > 0 and K > 0.

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

### Case 1. N(0) = 0

In this case the growth rate is 0 initially, so N(t) does not increase or decrease, so remains 0.

Case 2. 
$$N(0) = K$$

In this case the growth rate is 0 initially, so N(t) does not increase or decrease, so remains K.

Assume that r > 0 and K > 0.

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

### Case 1. N(0) = 0

In this case the growth rate is 0 initially, so N(t) does not increase or decrease, so remains 0.

Case 2. 
$$N(0) = K$$

In this case the growth rate is 0 initially, so N(t) does not increase or decrease, so remains K.

### Key takeaway

Both N(t) = 0 and N(t) = K are solutions to the ODE. They are called equalibrium solutions.

Assume that r > 0 and K > 0.

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

Case 3. 
$$0 \le N(0) \le K$$

In this case, N is initially increasing and so becomes more positive, slowing down as it gets close to K.

Assume that r > 0 and K > 0.

$$\frac{\mathrm{d}N}{\mathrm{d}t} = r\left(1 - \frac{N}{K}\right)N$$

### Case 3. $0 \le N(0) \le K$

In this case, N is initially increasing and so becomes more positive, slowing down as it gets close to K.

Case 4. 
$$N(0) \ge K$$

In this case N is initially decreasing but decreases slower and slower as it gets close to K.