

# Midterm 2 practice

UCLA: Math 31B, Spring 2017

*Instructor:* Noah White

*Date:*

*Version:* practice

- This exam has 4 questions, for a total of 40 points.
- Please print your working and answers neatly.
- Write your solutions in the space provided showing working.
- All final answers should be exact values. Decimal approximations will not be given credit.
- Indicate your final answer clearly.
- Full points will only be awarded for solutions with correct working.
- You may write on the reverse of a page or on the blank pages found at the back of the booklet however these will not be graded unless very clearly indicated.
- Non programmable and non graphing calculators are allowed.

Name: \_\_\_\_\_

ID number: \_\_\_\_\_

Discussion section: \_\_\_\_\_

Question	Points	Score
1	8	
2	10	
3	15	
4	7	
Total:	40	

1. Calculate the following limits using any technique you like.

(a) (5 points)

$$\lim_{x \rightarrow \infty} \left( \frac{x+1}{x} \right)^{x^2}$$

(b) (3 points)

$$\lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} + e^{\frac{1}{x}} \right)^{\arctan x}$$

2. For each of the following improper integrals say whether it converges or diverges. If the integral converges, you should say what value it converges to.

(a) (5 points)

$$\int_{-\frac{\pi}{2}}^0 \frac{\cos x}{\sin x} dx$$

(b) (5 points)

$$\int_1^{\infty} \frac{\ln x}{x^2} dx$$

3. For each of the following series say whether it converges or diverges. You do NOT need to justify your answer.

Grading scheme: 0 points for wrong, 1 point for no response, 3 points for correct.

(a) (3 points)  $\sum_{n=1}^{\infty} \frac{1}{n}$ .

(b) (3 points)  $\sum_{n=1}^{\infty} \frac{(n+3)^2}{n^4}$ .

(c) (3 points)  $\sum_{n=45}^{\infty} \frac{(-2)^n + 8^n}{3^{2n}}$ .

(d) (3 points)  $\sum_{n=1}^{\infty} a_n$  where the sequence of partial sums  $(s_N)_{N=1}^{\infty}$  is described by

$$s_N = \sum_{n=1}^N \frac{1}{\sqrt{n}}.$$

(e) (3 points)  $\sum_{n=1}^{\infty} a_n$  where the sequence of partial sums  $(s_N)_{N=1}^{\infty}$  is described by

$$s_N = \frac{1}{\sqrt{N}}.$$

4. (7 points) Let  $(F_n)$  be the *Fibonacci* sequence, that is

$$F_0 = 0, F_1 = 1, \text{ and } F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 2.$$

Consider the sequence  $(R_n)$  for  $n \geq 1$  defined by  $R_n = \frac{F_{n+1}}{F_n}$ , i.e.

$$\frac{1}{1}, \frac{2}{1}, \frac{3}{2}, \frac{5}{3}, \frac{8}{5}, \frac{13}{8}, \dots$$

You may assume that  $(R_n)$  converges to a limit  $L$  such that  $1 < L < 2$ . Find  $L$ .

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.

This page has been left intentionally blank. You may use it as scratch paper. It will not be graded unless indicated very clearly here and next to the relevant question.