

# Math 3B: Lecture 9

Noah White

October 12, 2016

# Review lecture

- Poll on Piazza

# Review lecture

- Poll on Piazza
- Please vote for what you want covered in the review lecture

## Last time

- Finding area under curves

## Last time

- Finding area under curves
- Definition of definite integral using Reimann Sum

## Last time

- Finding area under curves
- Definition of definite integral using **Reimann Sum**
- Using summation identities to calculate definite integrals

# The definite integral

## Defintion

The definite integral of a function  $f(x)$  is defined to be

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \frac{b-a}{n} \sum_{k=0}^{n-1} f(a + k \cdot \Delta x)$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$



## Example 2

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

### Solution

$$\int_0^1 x^2 - 4 \, dx = \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right)$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

### Solution

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \frac{k^2}{n^2} - 4\end{aligned}$$

## Example 2

$$\int_0^1 x^2 - 4 \, dx = \lim_{n \rightarrow \infty} \frac{1 - 0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right)$$

## Example 2

$$\int_0^1 x^2 - 4 \, dx = \lim_{n \rightarrow \infty} \frac{1 - 0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right)$$

## Example 2

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \lim_{n \rightarrow \infty} \frac{1 - 0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right) \\&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \frac{k^2}{n^2} - 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=0}^{n-1} k^2 - \frac{1}{n} \sum_{k=0}^{n-1} 4\end{aligned}$$

## Example 2

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right) \\&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \frac{k^2}{n^2} - 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=0}^{n-1} k^2 - \frac{1}{n} \sum_{k=0}^{n-1} 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \cdot \frac{1}{6}(n-1)n(2n-1) - \frac{4n}{n}\end{aligned}$$

## Example 2

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right) \\&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \frac{k^2}{n^2} - 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=0}^{n-1} k^2 - \frac{1}{n} \sum_{k=0}^{n-1} 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \cdot \frac{1}{6} (n-1)n(2n-1) - \frac{4n}{n} \\&= \lim_{n \rightarrow \infty} \frac{2n^2 - 3n + 1}{6n^2} - 4\end{aligned}$$

## Example 2

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{k=0}^{n-1} f\left(0 + k \cdot \frac{1}{n}\right) \\&= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \frac{k^2}{n^2} - 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=0}^{n-1} k^2 - \frac{1}{n} \sum_{k=0}^{n-1} 4 \\&= \lim_{n \rightarrow \infty} \frac{1}{n^3} \cdot \frac{1}{6} (n-1)n(2n-1) - \frac{4n}{n} \\&= \lim_{n \rightarrow \infty} \frac{2n^2 - 3n + 1}{6n^2} - 4 \\&= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$



# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .
- every input  $x$  produces a number as an output.

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\frac{d}{dx}F(x) = \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right]\end{aligned}$$




## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$


## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx$



## Why is the FTC true?

When  $h$  is very small  $\int_x^{x+h} f(t) \approx hf(x)$



## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt\end{aligned}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x)\end{aligned}$$

## Why is the FTC true?

We want to show that  $\frac{d}{dx}F(x) = f(x)$ .

$$\begin{aligned}\frac{d}{dx}F(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_a^{x+h} f(t) \, dt - \int_a^x f(t) \, dt \right] \\&= \lim_{h \rightarrow 0} \frac{1}{h} \int_x^{x+h} f(t) \, dt \\&= \lim_{h \rightarrow 0} \frac{1}{h} h f(x) \\&= f(x)\end{aligned}$$

## A consequence (corollary)

### Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Why?

Well  $F(x) = \int_a^x f(t) \, dt + C$  for some  $a$  and  $C$ . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$



## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

### Solution

An antiderivative of  $x^2 - 4$  is  $\frac{1}{3}x^3 - 4x$  so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

### Solution

An antiderivative of  $\sin x$  is  $-\cos x$  so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$

# The indefinite integral

We also use the following notation for the general antiderivative:

**Definition**

$$\int f(x) \, dx = F(x) + C$$

# The indefinite integral

We also use the following notation for the general antiderivative:

**Definition**

$$\int f(x) \, dx = F(x) + C$$

**Example**

$$\int \sin(x) - x \, dx = -\cos(x) - \frac{1}{2}x^2 + C$$

## Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,

## Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.



# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

## Substitution

Suppose  $u = g(x)$ , then

$$\int f(g(x))g'(x) \, dx = \int f(u) \, du$$

# Integration by substitution

- As we have seen, some antiderivatives are difficult to guess,
- especially if it involves reversing the chain rule.
- We solve this by introducing a new variable.

## Substitution

Suppose  $u = g(x)$ , then

$$\int f(g(x)) \frac{du}{dx} dx = \int f(g(x)) g'(x) dx = \int f(u) du$$

## Example 1

Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx = 2 \int \sqrt{u} \, du$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\begin{aligned}\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C\end{aligned}$$

## Example 1

### Question

$$\int 4x\sqrt{x^2 + 1} \, dx$$

### Solution

We use the substitution  $u = x^2 + 1$ , so  $\frac{du}{dx} = 2x$ , we can write the integral

$$\begin{aligned}\int 2 \cdot \sqrt{x^2 + 1} \cdot 2x \, dx &= 2 \int \sqrt{u} \, du \\ &= 2 \cdot \frac{2}{3} u^{\frac{3}{2}} + C \\ &= \frac{4}{3} (x^2 + 1)^{\frac{3}{2}} + C\end{aligned}$$

# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$



# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

## Example

$$\int_0^1 4x\sqrt{x^2 + 1} \, dx = 2 \int_1^2 \sqrt{u} \, du$$

# Integration by substitution (definite integrals)

## Substitution for definite integrals

Suppose  $u = g(x)$ , then

$$\int_a^b f(g(x))g'(x) \, dx = \int_{g(a)}^{g(b)} f(u) \, du$$

## Example

$$\begin{aligned}\int_0^1 4x\sqrt{x^2+1} \, dx &= 2 \int_1^2 \sqrt{u} \, du \\ &= 2 \left( \frac{2}{3} 2^{\frac{3}{2}} - \frac{2}{3} 1^{\frac{3}{2}} \right) = \frac{4}{3}(2\sqrt{2} - 1)\end{aligned}$$