### Math 3B: Lecture 18

Noah White

November 13, 2017

The most straighforward way of checking a function y = f(x) is a solution to a differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = g(x,y)$$

is to simply plug it in to both sides.

The most straighforward way of checking a function y = f(x) is a solution to a differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = g(x,y)$$

is to simply plug it in to both sides.

#### Example

The function  $y=e^{\sin x}$  is a solution of  $\frac{\mathrm{d}y}{\mathrm{d}x}=y\cos x$ . To check note that

The most straighforward way of checking a function y = f(x) is a solution to a differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = g(x,y)$$

is to simply plug it in to both sides.

#### Example

The function  $y=e^{\sin x}$  is a solution of  $\frac{\mathrm{d}y}{\mathrm{d}x}=y\cos x$ . To check note that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = e^{\sin x} \cos x$$

The most straighforward way of checking a function y = f(x) is a solution to a differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = g(x,y)$$

is to simply plug it in to both sides.

#### Example

The function  $y = e^{\sin x}$  is a solution of  $\frac{dy}{dx} = y \cos x$ . To check note that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = e^{\sin x} \cos x$$
$$y \cos x = e^{\sin x} \cos x$$

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f(y)g(x)$$

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f(y)g(x)$$

It works by

1. moving all the y's over to the left:

$$\frac{1}{f(y)}\frac{\mathrm{d}y}{\mathrm{d}x} = g(x)$$

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f(y)g(x)$$

It works by

1. moving all the y's over to the left:

$$\frac{1}{f(y)}\frac{\mathrm{d}y}{\mathrm{d}x} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{\mathrm{d}y}{\mathrm{d}x} \, \mathrm{d}x = \int g(x) \, \mathrm{d}x.$$

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f(y)g(x)$$

It works by

1. moving all the y's over to the left:

$$\frac{1}{f(y)}\frac{\mathrm{d}y}{\mathrm{d}x}=g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{\mathrm{d}y}{\mathrm{d}x} \, \mathrm{d}x = \int g(x) \, \mathrm{d}x.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} \, \mathrm{d}y = \int g(x) \, \mathrm{d}x.$$

Separation of variables is a very powerful technique for solving ODEs. It works, whenever we can write out an ODE in the form

$$\frac{\mathrm{d}y}{\mathrm{d}x} = f(y)g(x)$$

It works by

1. moving all the y's over to the left:

$$\frac{1}{f(y)}\frac{\mathrm{d}y}{\mathrm{d}x} = g(x)$$

2. integrating both sides (with respect to x):

$$\int \frac{1}{f(y)} \frac{\mathrm{d}y}{\mathrm{d}x} \, \mathrm{d}x = \int g(x) \, \mathrm{d}x.$$

3. we can use the integration by substitution formula to rewrite the left hand side:

$$\int \frac{1}{f(y)} \, \mathrm{d}y = \int g(x) \, \mathrm{d}x.$$

4. solve for y!

# Examples

On the board...