Math 3B: Lecture 22

Noah White

March 8, 2017

International Womens' Day

Today is International Womens Day



Let's meet some of the women who have influenced the mathematics we are learning.

Hypatia (c.360-415)



Maria Gaetana Agnesi (1718-1799)



Maria Gaetana Agnesi (1718-1799)



Sofia Kovalevskaya (1850-1891)



Sofia Kovalevskaya (1850-1891)




Emmy Noether (1882-1935)



Mary Cartwright (1900-1998)



Nalini Joshi



Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t} = f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

We want points (t, y) such that f(y) = 0.

• Suppose f(a) = 0.

Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions

We want points (t, y) such that f(y) = 0.

- Suppose f(a) = 0.
- Then (t, a) is on the nullcline, for any t.

Deafinition

An ODE of the form

$$\frac{\mathrm{d}y}{\mathrm{d}t} = f(y)$$

i.e. where the right hand side does not depend on t, is called autonomous

Important property

The nullclines of an autonomous equation are horizontal straight lines! Nullclines = equilibrium solutions


We want points (t, y) such that f(y) = 0.

- Suppose f(a) = 0.
- Then (t, a) is on the nullcline, for any t.
- So the line y = a is part of the nullcline, whenever f(a) = 0.


Slope fields and nullclines for autonomous systems

•	

Phase lines/diagram



Phase lines/diagram



Phase lines/diagram



Recipe to draw phase lines

1. Draw a vertical corresponding to y axis

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- 3. Draw up arrows on intervals between dots where the derivative is positive

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- 3. Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition

 An equalibrium is stable if the two arrows are pointing towards it.

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative

Definition


- An equalibrium is stable if the two arrows are pointing towards it.
- It is unstable if the two arrows are pointing away from it.

Recipe to draw phase lines

- 1. Draw a vertical corresponding to y axis
- 2. Draw dots where equilibrim solutions live
- Draw up arrows on intervals between dots where the derivative is positive
- 4. Draw down arrows on intervals between dots where the derivative is negative


Definition


- An equalibrium is stable if the two arrows are pointing towards it.
- It is unstable if the two arrows are pointing away from it.
- It is semistable if the arrows point in the same direction.





$$\frac{\mathrm{d}y}{\mathrm{d}t} = y(y-10)(25-y)$$


$$\frac{\mathrm{d}y}{\mathrm{d}t} = y(y-10)(25-y)$$









































Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t} = f(y)$$

- (i.e. f(a) = 0) then a is
 - stable if f'(a) < 0

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

- stable if f'(a) < 0
- unstable if f'(a) > 0

Classification of equilibria

If a is an equillibrium of

$$\frac{\mathrm{d}y}{\mathrm{d}t}=f(y)$$

(i.e. f(a) = 0) then a is

- stable if f'(a) < 0
- unstable if f'(a) > 0
- indeterminate if f'(a) = 0

