

# Math 3B: Lecture 7

Noah White

October 13, 2017

## Learning assistant office hours

- Learning assistants (usually Nick) will be holding office hours

## Learning assistant office hours

- Learning assistants (usually Nick) will be holding office hours
- Fridays, 6-8pm in Covel Commons 237

## Learning assistant office hours

- Learning assistants (usually Nick) will be holding office hours
- Fridays, 6-8pm in Covell Commons 237
- Big study room, second floor

## Learning assistant office hours

- Learning assistants (usually Nick) will be holding office hours
- Fridays, 6-8pm in Covel Commons 237
- Big study room, second floor
- This Friday only: moved to 5-7pm

Bad news

- I will be away Thurs, Fri next week

Good news

# News

## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled

## Good news

## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled
- Nivedita Bhaskhar will take Friday's review

## Good news



## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled
- Nivedita Bhaskhar will take Friday's review

## Good news

- I will post notes online (no video)

## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled
- Nivedita Bhaskhar will take Friday's review

## Good news

- I will post notes online (no video)
- Rescheduled office hours: Tue 10-11am, Wed 4-6pm

## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled
- Nivedita Bhaskhar will take Friday's review

## Good news

- I will post notes online (no video)
- Rescheduled office hours: Tue 10-11am, Wed 4-6pm
- Office hour Sunday 10/22 TBA

## Bad news

- I will be away Thurs, Fri next week
- office hours cancelled
- Nivedita Bhaskhar will take Friday's review

## Good news

- I will post notes online (no video)
- Rescheduled office hours: Tue 10-11am, Wed 4-6pm
- Office hour Sunday 10/22 TBA
- Practice exams are at the extreme end of the spectrum of difficulty

# Midterm information

## Midterm 1

- Practice midterm

# Midterm information

## Midterm 1

- Practice midterm
- arrive on time

# Midterm information

## Midterm 1

- Practice midterm
- arrive on time
- calculators, cheat sheets

# Midterm information

## Midterm 1

- Practice midterm
- arrive on time
- calculators, cheat sheets
- expected average, grades



# Antiderivatives

We will be concentrating on solving differential equations of the form

$$\frac{dy}{dx} = f(x)$$

# Antiderivatives

We will be concentrating on solving differential equations of the form

$$\frac{dy}{dx} = f(x)$$

A solution  $y = F(x)$  is called an **antiderivative** of  $f(x)$ .

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + 4$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + 8$$

## Example 1

### Question

What is the antiderivative of  $f(x) = 2x$ ?

### Solution

$$F(x) = x^2 + C$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?



## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2 - x$$

## Example 2

### Question

What is the antiderivative of  $f(x) = x^3 + 4x - 1$ ?

### Solution

$$F(x) = \frac{1}{4}x^4 + 2x^2 - x + C$$

## Example 3

### Question

What is the antiderivative of  $f(x) = e^{2x}$ ?

## Example 3

### Question

What is the antiderivative of  $f(x) = e^{2x}$ ?

### Solution

$$F(x) = \frac{1}{2}e^{2x}$$

## Example 3

### Question

What is the antiderivative of  $f(x) = e^{2x}$ ?

### Solution

$$F(x) = \frac{1}{2}e^{2x}$$

## Example 4

### Question

What is the antiderivative of  $f(x) = \frac{1}{x}$  (for  $x > 0$ )?



## Example 4

### Question

What is the antiderivative of  $f(x) = \frac{1}{x}$  (for  $x > 0$ )?

### Solution

$$F(x) = \ln x$$

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

### Solution

Note that  $f(x) = (1+x)^{-2}$ . So

$$F(x) = \frac{1}{1+x}$$

## Example 5

### Question

What is the antiderivative of  $f(x) = \frac{1}{(1+x)^2}$ ?

### Solution

Note that  $f(x) = (1+x)^{-2}$ . So

$$F(x) = -\frac{1}{1+x}$$

## Example 6

### Question

What is the antiderivative of  $f(x) = 2x \cos x^2$ ?

## Example 6

### Question

What is the antiderivative of  $f(x) = 2x \cos x^2$ ?

### Solution

$$F(x) = \sin x^2$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$



## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$

$$F(x) = x^{\frac{1}{2}}$$

## Example 7

### Question

What is the antiderivative of  $f(x) = \frac{1}{\sqrt{x}}$ ?

### Solution

$$f(x) = x^{-\frac{1}{2}}$$


$$F(x) = 2x^{\frac{1}{2}}$$

# Slope fields

In some cases it is impossible to find the antiderivative (without special functions). E.g.

$$f(x) = e^{-x^2}$$

But we can still (approximately) graph the antiderivative! First we draw the slope field



# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.

## Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?

## Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$

# Slope fields (how to draw)

How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$



# Slope fields (how to draw)


How to draw a slope field for

$$\frac{dy}{dx} = f(x)$$

1. Draw the  $xy$ -plane.
2. At every point  $(x, y)$  what would the slope of  $y = F(x)$  be if it passed through that point?
3. Answer given by differential equation above, slope is  $f(x)$
4. Draw a small arrow with slope  $f(x)$  and the point  $(x, y)$
5. Do this for a grid of points on the  $xy$ -plane.


## Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




## Example 1

$$f(x) = e^{-x^2} \text{ with } F(0) = -0.25$$




## Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



## Example 2

$$f(x) = \sin(x^2) \text{ with } F(0) = 0.1$$



# Accumulated change

Often we encounter problems involving **accumulated change**.

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?



# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

# Accumulated change

Often we encounter problems involving **accumulated change**.

## Example

A rocket is accelerating at a rate of  $a(t) = 0.3t^2$  metres per second squared. What is the rocket's velocity at  $t = 30$ ?

## Example

A population grows at a rate of  $0.5P(t)$  people per year. How much does the population increase over 10 years?

These problems involve finding the area under some curve.

## Example 1


If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

## Example 1

If a car travels at a constant speed of 30 miles per hour, how much distance does it cover after 2.5 hours?

### Solution

We model the car's speed using the function  $s(t) = 30$ . So we can see that the area under this curve



is the distance travelled (75 miles)

## Example 2


If a car accelerates for 20 seconds at a rate of  $2m/s^2$  and then decelerates for 30 seconds at a rate of  $1m/s^2$ , how far has it travelled?

## Example 2

If a car accelerates for 20 seconds at a rate of  $2\text{m/s}^2$  and then decelerates for 30 seconds at a rate of  $1\text{m/s}^2$ , how far has it travelled?


### Solution

The car's speed is given by  $s(t) = 2t$  when  $0 \leq t \leq 20$  and  $s(t) = 60 - t$  when  $20 \leq t \leq 50$ . So the graph looks like




## More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?



## More complicated rates of change

Suppose we have a car whose speed is described by the following curve. How far has it travelled in this time?





## Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).

# Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between  $t = a$  and  $t = b$ ?

# Accumulated change

- Suppose we know the rate of change  $f(t)$ , of some quantity (distance, water flow, population, etc).
- How do we find the total amount by which this changes between  $t = a$  and  $t = b$ ?
- Answer: area under  $f(t)$  between  $a$  and  $b$ .

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

A first approach to calculating the area under a curve is to approximate using rectangles:

## Areas under general curves

We would like to calculate the area between a function  $f(x)$  and the  $x$ -axis, between  $x = a$  and  $x = b$ .

A first approach to calculating the area under a curve is to approximate using rectangles:

(Too hard to draw, lets look at an animation)

# The definite integral

## Defintion

The definite integral of a function  $f(x)$  is defined to be

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \Delta x \sum_{k=1}^n f(a + k\Delta x)$$

where  $\Delta x = \frac{b-a}{n}$ .

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$



# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .

# The fundamental theorem of calculus

## Theorem

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

That is,  $F(x) = \int_a^x f(t) dt$  is an antiderivative of  $f(x)$ !

## Note

- $F(x) = \int_a^x f(t) dt$  is a function of  $x$ .
- every input  $x$  produces a number as an output.

## A consequence (corollary)

### Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

# A consequence (corollary)

## Corollary

For **any** antiderivative  $F(x)$  of  $f(x)$

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

## Why?

Well  $F(x) = \int_a^x f(t) \, dt + C$  for some  $a$  and  $C$ . So

$$\begin{aligned} F(b) - F(a) &= \int_a^b f(t) \, dt + C - \int_a^a f(t) \, dt - C \\ &= \int_a^b f(t) \, dt \end{aligned}$$

## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

## Example 1

### Question

Evaluate the definite integral

$$\int_0^1 x^2 - 4 \, dx$$

### Solution

An antiderivative of  $x^2 - 4$  is  $\frac{1}{3}x^3 - 4x$  so

$$\begin{aligned}\int_0^1 x^2 - 4 \, dx &= \frac{1}{3} \cdot 1^3 - 4 - \frac{1}{3} \cdot 0^3 + 4 \cdot 0 \\ &= \frac{1}{3} - 4 = -\frac{11}{3}\end{aligned}$$



## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

## Example 2

### Question

Evaluate the definite integral

$$\int_0^{\pi} \sin x \, dx$$

### Solution

An antiderivative of  $\sin x$  is  $-\cos x$  so

$$\begin{aligned}\int_0^{\pi} \sin x \, dx &= -\cos \pi + \cos 0 \\ &= -(-1) + 1 = 2\end{aligned}$$