Advanced Formalisms and Computational Strategies in Numerical Relativity

The discipline of numerical relativity serves as the definitive framework for exploring the most extreme and dynamic environments in the universe, where gravitational fields are both extraordinarily strong and rapidly varying.[1] At its mathematical core, numerical relativity is the pursuit of solutions to the Einstein field equations—a set of ten coupled, non-linear, second-order partial differential equations—using advanced computational algorithms.[3] These equations, $G_{\mu\nu} = 8\pi G T_{\mu\nu}$, relate the geometry of spacetime to the energy-momentum content within it, but their inherent complexity restricts exact analytical solutions to highly symmetric or idealized cases.[4] Consequently, the modeling of realistic astrophysical phenomena, such as the coalescence of binary black holes, the merger of neutron stars, and the non-spherical collapse of stellar cores, necessitates the use of high-performance computing to resolve the evolution of spacetime in four dimensions.[7]

The evolution of the field has been marked by several distinct eras of progress. In the mid-1960s, pioneering attempts by Hahn and Lindquist aimed to simulate the head-on collision of two black holes, but these efforts were stymied by both the limited computational resources of the time and a lack of understanding regarding the mathematical stability of the evolution equations.[4] It was not until the late 1970s that Smarr and Eppley produced the first successful simulations of black hole collisions, though these remained restricted to axisymmetry and low resolution.[4] The modern era of numerical relativity was inaugurated in 2005 with the "breakthrough" achieved by Frans Pretorius, followed closely by the development of the "moving puncture" approach by groups at the University of Texas at Brownsville and Rochester Institute of Technology.[10] These advancements finally allowed for stable, long-term simulations of orbiting black hole binaries, providing the precise gravitational waveforms necessary for the burgeoning field of gravitational-wave astronomy.[8]

The 3+1 Cauchy Formulation and Spacetime Foliation

To solve the Einstein equations on a computer, one must abandon the covariant four-dimensional perspective in favor of a 3+1 decomposition, which treats the four-dimensional manifold as a sequence of three-dimensional spatial hypersurfaces evolving through time.[3] This process, often referred to as the Cauchy problem or initial value problem, allows the equations to be recast in a form where an initial state is specified and then "pushed" forward into the future.[6]

Geometric Decomposition and Projection

The foliation of spacetime begins by defining a scalar time function t whose level surfaces Σ_t are spacelike hypersurfaces.[5] A unit normal vector n^μ is defined for these slices, representing the four-velocity of "Eulerian" observers who remain orthogonal to the spatial slices.[5] The four-metric $g_{\mu\nu}$ is then decomposed using the induced spatial metric γ_{ij} and two gauge variables: the lapse function α and the shift vector β^i.[5] The lapse function determines the proper time elapsed between slices for an observer moving normally to the hypersurface, while the shift vector describes how spatial coordinates rotate or shift across the slice as time progresses.[6]

The fundamental variables of the 3+1 formulation are the spatial metric γ_{ij} and the extrinsic curvature K_{ij}.[5] While γ_{ij} describes the intrinsic geometry (distances and angles) within the three-dimensional slice, K_{ij} describes how that slice is embedded in the four-dimensional spacetime, effectively acting as the time derivative of the spatial metric.[5] The relationship between the two is defined by the Lie derivative along the normal vector:

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + D_i \beta_j + D_j \beta_i$$

where D_i is the covariant derivative compatible with the spatial metric γ_{ij}.[6]

The Constraint and Evolution System

The Einstein equations, when decomposed, yield two types of equations. The four constraint equations—the Hamiltonian constraint and the three momentum constraints—must be satisfied on every spatial slice.[5] These do not involve second time derivatives and instead represent the consistency requirements of the geometry.[6] The remaining six equations are evolution equations, describing the time-rate-of-change of the extrinsic curvature.[5]

Equation Type	Variable	Mathematical Role	Physical Meaning
Hamiltonian Constraint	γ_{ij}, K_{i}	Elliptic equation	Local energy density must match geometry.

Momentum Constraints	γ_{ij}, K_{i}	Elliptic system	Local momentum flux must match geometric shear.
Kinematic Evolution	γ_{ij}	Hyperbolic system	Relates metric change to extrinsic curvature.
Dynamic Evolution	K_{ij}	Hyperbolic system	Describes how curvature responds to stress-energy.

In a vacuum, the Hamiltonian constraint is $R + K^2 - K_{ij}K^{ij} = 0$, where R is the three-dimensional Ricci scalar and K is the trace of the extrinsic curvature.[5] The momentum constraints are $D_j(K^{ij} - \gamma^{ij}K) = 0$.[5] These equations form the bedrock of the initial data problem: one cannot simply pick arbitrary starting values for γ_{ij} and K_{ij}; they must be solved for simultaneously to ensure the simulation begins in a state allowed by general relativity.[16]

Initial Data and the Conformal Transverse-Traceless Formalism

Constructing initial data for black hole binaries involves specifying the positions, masses, spins, and momenta of the objects while satisfying the constraints.[12] The Conformal Transverse-Traceless (CTT) approach is the most widely utilized method for this purpose.[16] It employs a conformal transformation to separate the physical variables into a background geometry and a conformal factor ψ that encodes the gravitational field's strength.[5]

Conformal Transformations in Initial Data

In the CTT formalism, the physical metric is scaled as $\gamma_{ij} = \psi^4\tilde{\gamma}_{ij}$, where $\tilde{\gamma}_{ij}$ is a freely specifiable conformal metric.[5] The extrinsic curvature is further decomposed into its trace K and a traceless part $\tilde{A}_{ij}$, which is itself split into a longitudinal part and a transverse-traceless part $\tilde{A}_{ij}^{TT}$.[16] This decomposition is highly advantageous because it transforms the non-linear constraints into a set of elliptic partial differential equations that are well-suited for numerical

solvers.[16]

A common simplification used in classrooms and foundational research is to assume a flat conformal metric ($\tilde{\gamma}_{ij} = \delta_{ij}$) and a maximal slicing condition ($K = 0$).[16] Under these assumptions, the Hamiltonian constraint reduces to a non-linear elliptic equation for the conformal factor [16]:

$$\tilde{\nabla}^2 \psi + \frac{1}{8} \psi^{-7} \tilde{A}_{ij} \tilde{A}^{ij} = 0$$

where $\tilde{\nabla}^2$ is the flat-space Laplacian.[16] The momentum constraints become a set of linear equations for a vector potential X^i , where $\tilde{A}_{ij} = (\mathbb{L}X)_{ij} + \tilde{A}^{TT}_{ij}$.[5]

Punctures and the Brandt-Brügmann Method

One of the most significant hurdles in initial data construction was how to represent black holes without encountering the physical singularity at their center.[16] The "puncture" method, proposed by Brandt and Brügmann in 1997, provided a robust solution.[23] Instead of excising a region of the grid, this method factorizes the conformal factor into a singular part (representing the punctures) and a regular part u that is solved for numerically.[16]

In the case of N black holes initially at rest, the conformal factor takes the form of the Brill-Lindquist solution: $\psi = 1 + \sum_{n=1}^{N} \frac{m_n}{2|r - r_n|}$.[16] For black holes with momentum and spin, the Bowen-York extrinsic curvature is added, which provides an analytical solution to the momentum constraints in a conformally flat space.[12] However, Bowen-York data possess an inherent limitation: they cannot represent black holes with spins higher than $\chi \approx 0.928$ because the assumption of conformal flatness becomes physically inconsistent at the extremal limit.[12]

Initial Data Type	Conformal Metric	Topology	Capabilities
Misner	Flat	Wormhole	Equal mass, time-symmetric black holes.
Brill-Lindquist	Flat	Multi-sheet	Multiple black holes

			at rest.
Bowen-York	Flat	Puncture	Boosted and spinning black holes ($\chi <$).
SKS/HiSpID	Non-flat	Puncture	Nearly extremal spins ($\chi >$).

To move beyond the Bowen-York limit, modern simulations use "High-Spin Initial Data" (HiSpID) or superposed Kerr-Schild (SKS) data, which drop the assumption of conformal flatness.[12]

These methods allow for simulations of black holes with spins up to $\chi = 0.9997$, which are essential for modeling astrophysical populations that may be rotating near the extremal limit.[12]

Evolution Systems: Hyperbolicity and Stability

The original ADM evolution equations, while formally correct, are not well-suited for numerical integration because they are "weakly hyperbolic".[14] In a weakly hyperbolic system, the equations admit modes that grow polynomially or exponentially with time due to numerical truncation errors, leading to the eventual collapse of the simulation.[14] Achieving long-term stability requires reformulating the Einstein equations into "strongly hyperbolic" or "symmetric hyperbolic" systems, where the solutions are guaranteed to depend continuously on the initial data.[14]

The BSSN Formulation

The BSSN (Baumgarte-Shapiro-Shibata-Nakamura) formulation is currently the most popular choice for simulations using the moving puncture method.[14] BSSN improves stability by introducing several auxiliary variables and performing a conformal decomposition of the evolution equations.[14] The fundamental variables are:

- A conformal factor $\phi = \frac{1}{12}\ln(\gamma/\bar{\gamma})$ or $\chi = e^{-4\phi}$.[29]
- The conformal metric $\tilde{\gamma}_{ij} = e^{-4\phi}\gamma_{ij}$ with unit determinant.[14]
- The trace of the extrinsic curvature K.[29]
- The conformal traceless extrinsic curvature $\tilde{A}_{ij} = e^{-4\phi}(K_{ij} - \frac{1}{3}\gamma_{ij}K)$.[29]
- The conformal connection functions $\tilde{\Gamma}^i = \tilde{\gamma}^{jk}\tilde{\Gamma}^i_{jk}$.[14]

By evolving the $\tilde{\Gamma}^i$ variables independently, the BSSN system effectively makes the Ricci tensor appear as a wave-like operator, which significantly enhances the hyperbolicity and numerical stability of the code.[14]

The Z4c Formulation and Constraint Damping

While BSSN is robust, it does not have a mechanism to actively correct violations of the constraints that arise from numerical noise.[31] The Z4c formulation (Conformal Z4) addresses this by introducing an additional dynamic variable θ and a damping vector Z_i.[31] In the Z4c scheme, Hamiltonian and momentum constraint violations are treated as propagating waves that are damped away over time.[31]

The Z4c equations for the three-metric and extrinsic curvature include terms proportional to damping parameters κ_1 and κ_2 [31]:

$$\partial_t K_{ij} = \cdots - \kappa_1(1+\kappa_2)\gamma_{ij}\Theta$$

These parameters are tuned to ensure that high-frequency, low-amplitude violations are damped exponentially.[31] Recent studies have shown that Z4c is often superior to BSSN for simulations involving matter, such as binary neutron star mergers, as it maintains Hamiltonian constraint violations that are one to three orders of magnitude lower than BSSN.[32]

Gauge Choices: The Physics of Coordinates

In numerical relativity, the choice of the lapse α and the shift β^i —the gauge—is a critical component of the "art" of the simulation.[16] A poor gauge choice can lead to "slice stretching," where the spatial coordinates become so distorted that the simulation loses resolution and crashes.[14]

Singularity Avoidance and 1+log Slicing

To evolve a black hole spacetime without the coordinates falling into the singularity, one needs a "singularity-avoiding" slicing condition.[14] Maximal slicing ($K = 0$) is one such choice, as it maximizes the volume of the spatial slices and prevents them from reaching the singularity.[14] However, maximal slicing requires solving an elliptic equation at every time step, which is computationally expensive.[14]

A more efficient alternative is "1+log" slicing, part of the Bona-Massó family of hyperbolic slicing

conditions.[16] The evolution of the lapse is governed by:

$$\partial_t \alpha - \beta^i \partial_i \alpha = -2\alpha K$$

As the simulation approaches a region of high curvature, K becomes large, and the lapse function rapidly drops toward zero.[14] This phenomenon, known as "lapse collapse," effectively freezes the flow of coordinate time near the black hole, allowing the simulation to continue in the exterior regions while "waiting" for the high-curvature dynamics to resolve.[16]

Managing Motion with Gamma-Drivers

For simulations where black holes move across the grid, the shift vector must be carefully managed to prevent coordinate distortion.[16] The "gamma-driver" shift conditions are designed to keep the coordinates "attached" to the moving punctures.[14] The hyperbolic version of this condition is:

$$\partial_t \beta^i = \frac{3}{4} B^i, \quad \partial_t B^i = \partial_t \tilde{\Gamma}^i - \eta B^i$$

where η is a damping parameter that prevents the shift from oscillating.[14] This "moving puncture" gauge—combining 1+log slicing with gamma-driver shifts—is the standard for most modern binary black hole simulations, as it allows the singularity to be represented as a point that moves across the Cartesian grid without the need for complex excision algorithms.[8]

The Trumpet Attractor and the $3/2M$ Limiting Surface

An insightful observation in numerical relativity is that when puncture initial data are evolved with the standard moving puncture gauge, the internal geometry of the black hole undergo a coordinate transition.[21] Most initial data, such as Brill-Lindquist, are "wormhole" data, meaning they connect two distinct, asymptotically flat universes.[21] However, the moving puncture gauge is not compatible with the wormhole topology over the long term.[25]

As the evolution progresses, the spatial slices "lose contact" with the second asymptotically flat end and instead settle into a "trumpet" geometry.[25] In this state, the slices smoothly penetrate the event horizon but terminate on a "limiting surface" of finite areal radius.[37] For a Schwarzschild black hole, this surface occurs at an areal radius of $R = 1.5M$, shielding the simulation from the curvature singularity at $R = 0$.[21]

This trumpet geometry is a strong attractor for the 1+log and gamma-driver gauges.[36] Because the coordinates settle into this stationary state, the "trumpet" represents a more natural numerical representation of a black hole than the initial wormhole-like data.[25] Consequently, some researchers now advocate for starting with trumpet initial data from the outset to avoid the coordinate transients that occur during the beginning of a simulation.[25]

Critical Phenomena and the Threshold of Collapse

Beyond the study of stable black holes, numerical relativity has uncovered fundamental behaviors in the "critical" regime of gravitational collapse.[40] In 1993, Matthew Choptuik used adaptive mesh refinement to study the collapse of a massless scalar field in spherical symmetry.[16] By varying an initial parameter p (such as the amplitude of the field), he identified a critical threshold p^* separating fields that disperse from those that form black holes.[16]

Near this threshold, Choptuik discovered several "critical phenomena" analogous to those in statistical mechanics [40]:

1. **Mass Scaling:** For simulations just above the threshold, the mass of the black hole follows a universal power law: $M_{BH} \propto (p - p^*)^\gamma$, where $\gamma \approx 0.37$ for a scalar field.[16]
2. **Self-Similarity:** The critical solution ($p = p^*$) is self-similar, meaning it looks the same when zoomed in by a specific factor in both space and time.[41]
3. **Discrete Echoing:** The scalar field collapse exhibits "discrete self-similarity" (DSS), where the field repeats its structure with an echoing period of $\Delta \approx 3.4$.[42]

This work provided the first evidence that black holes can form with arbitrarily small masses, and that the fine-tuning of initial data can lead to a naked singularity—a region of infinite curvature that is not hidden by an event horizon.[40] This remains one of the most significant theoretical insights ever produced by numerical relativity.[40]

Higher Dimensions and Black Rings

Numerical relativity is not confined to the four-dimensional spacetime of our experience; it is increasingly used to explore gravity in $D > 4$ dimensions, as predicted by string theory and other unified models.[44] In higher dimensions, the uniqueness theorems of four-dimensional general relativity—which state that the Kerr metric is the only stationary black hole solution—do not hold.[44]

The most famous example is the "black ring," discovered by Emparan and Reall in 2001.[44] A black ring is a five-dimensional black hole with an event horizon topology of $S^1 \times$, rather

than the spherical S^3.[44] These objects can rotate along the S^1 direction, with centrifugal force balancing their gravitational self-attraction.[44]

Property	4D Black Hole (Kerr)	5D Black Ring
Topology	S^2	$S^1 \times$
Uniqueness	Unique for given M, J	Multiple solutions possible for same M, J
Stability	Generally stable	Unstable to Gregory-Laflamme mode
Symmetry	Axially symmetric	Axially symmetric (along S^1)

Numerical simulations of perturbed black rings have revealed that they are often unstable.[16] "Thin" black rings are susceptible to a Gregory-Laflamme-type instability, where the ring develops "bulges" and "necks".[16] As the evolution progresses, these necks become thinner and thinner, suggesting that the ring may eventually "pinch off" and break, potentially violating the cosmic censorship conjecture in five dimensions.[16]

Computational Infrastructure and Advanced Methods

The success of numerical relativity is heavily dependent on the available enabling software and hardware.[10] Modern simulations rely on highly scalable infrastructures like the **Einstein Toolkit**, which provides a collection of modules (or "thorns") for solving the 3+1 equations, handling relativistic hydrodynamics, and managing the computational grid.[10]

Adaptive Mesh Refinement (AMR)

One of the most critical components of modern codes is Adaptive Mesh Refinement (AMR).[10] Compact binary systems involve vast differences in scale: the resolution required to track the event horizon of a black hole is orders of magnitude finer than that needed to track the gravitational waves as they propagate to the "wave zone" thousands of kilometers away.[9] AMR algorithms, such as those implemented in the **Carpet** driver, recursively refine the grid in regions of high curvature, allowing simulations to be both accurate and computationally efficient.[10]

Pseudospectral Methods

While most "moving puncture" codes use finite difference methods, the **SXS** (Simulating eXtreme Spacetimes) collaboration utilizes pseudospectral methods in their **SpEC** code.[8] Pseudospectral methods represent the solution as a series of basis functions (such as Chebyshev polynomials or Fourier series), which allows for exponential convergence as the number of basis functions increases.[8] These methods are highly accurate but require smooth, analytic domains; consequently, they typically use excision (removing the black hole interior) rather than the puncture method.[8]

The Scale of Modern Computing

Today's simulations are performed on exascale-class supercomputers, which execute quadrillions of operations per second.[16] A single binary neutron star simulation, involving microphysics like neutrino transport and magnetic fields, can require millions of CPU-hours and generate terabytes of data.[10] The future of the field involves adapting these codes to new hardware architectures, such as GPUs, which offer massive parallelism but require a fundamental rethink of the underlying numerical algorithms.[11]

Synthesis and Final Perspectives

The maturity of numerical relativity has transformed it from a theoretical curiosity into an essential tool for high-energy astrophysics and fundamental physics.[3] By successfully solving the initial data problem through formalisms like CTT and punctures, and stabilizing the evolution problem through BSSN and Z4c, researchers have provided the scientific community with a "gravitational laboratory".[1]

Key synthesis points from the analysis indicate that:

- The transition from wormhole to trumpet topologies is a hallmark of the moving puncture approach, revealing that the "trumpet" is the natural stationary state of a black hole in modern gauges.[21]
- The Z4c formulation provides a superior framework for matter-rich environments by actively damping Hamiltonian constraint violations, which is vital for the multi-messenger era of neutron star observations.[32]
- The discovery of critical phenomena in gravitational collapse demonstrates that numerical relativity is a powerful engine for theoretical discovery, uncovering regimes of gravity that were previously invisible to analytical methods.[40]

As we move toward the era of third-generation gravitational wave detectors, numerical relativity will face new demands for accuracy and physical realism.[11] Addressing these will require a continuous refinement of gauge conditions, the integration of increasingly complex microphysics, and the exploitation of next-generation high-performance computing

architectures.[10] The field remains one of the most dynamic and challenging frontiers of modern science, providing the ultimate testbed for our understanding of Einstein's legacy.[2]

Works cited

1. (PDF) Introduction to Numerical Relativity - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/344268954_Introduction_to_Numerical_Relativity
2. Numerical Relativity: Solving Einstein's Equations on the Computer, accessed March 3, 2026, https://nzdr.ru/data/media/biblio/kolxoz/P/PGr/Baumgarte%20T.W.,%20Shapiro%20S.L.%20Numerical%20Relativity..%20Solving%20Einstein's%20Equations%20on%20the%20Computer%20(CUP,%202010)(ISBN%20052151407X)(O)(718s)_PGr_.pdf
3. Fundamentals of numerical relativity for gravitational wave sources - Physics and Astronomy, accessed March 3, 2026, https://physics.byu.edu/faculty/bergeson/docs/science-numerical-relativity.pdf
4. (PDF) Introduction to 3+1 Numerical Relativity - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/228588827_Introduction_to_31_Numerical_Relativity
5. Initial Data for Numerical Relativity - PMC - NIH, accessed March 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5660886/
6. arXiv:gr-qc/0106072v3 29 Jun 2001, accessed March 3, 2026, https://arxiv.org/pdf/gr-qc/0106072
7. Introduction to Numerical Relativity (1/2) | Vidéo | Carmin.tv, accessed March 3, 2026, https://www.carmin.tv/en/video/introduction-to-numerical-relativity-1-2
8. "Puncture Initial Data and Evolution of Black Hole Binaries with High S" by Ian Ruchlin, accessed March 3, 2026, https://repository.rit.edu/theses/8797/
9. Exploring New Physics Frontiers Through Numerical Relativity - PMC, accessed March 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5255938/
10. The Einstein Toolkit: a community computational infrastructure for relativistic astrophysics - MPG.PuRe, accessed March 3, 2026, https://pure.mpg.de/rest/items/item_1481630/component/file_1481629/content
11. Improved Moving-Puncture Techniques for Compact Binary Simulations - arXiv, accessed March 3, 2026, https://arxiv.org/html/2404.01137v1
12. Puncture initial data for black-hole binaries with high spins and high boosts - The Research Repository @ WVU - West Virginia University, accessed March 3, 2026, https://researchrepository.wvu.edu/cgi/viewcontent.cgi?article=1299&context=faculty_publications
13. Solving the Einstein Equations Numerically - arXiv, accessed March 3, 2026, https://arxiv.org/html/2405.06035v1
14. Binary black hole evolutions in numerical relativity - Moriond, accessed March 3, 2026,

https://moriond.in2p3.fr/J03/transparencies/6_friday/2_afternoon/pollney.pdf
15. Introduction to Numerical Relativity, accessed March 3, 2026, https://www.oit.ac.jp/labs/is/system/shinkai/lecture/winterAPCTP/slideNR_2a_standardF.pdf
16. tactiq-free-transcript-WWaiLRkj2R4.txt
17. (PDF) Status of numerical relativity - Academia.edu, accessed March 3, 2026, https://www.academia.edu/32008274/Status_of_numerical_relativity
18. arXiv:2406.08410v1 [gr-qc] 12 Jun 2024 - MPG.PuRe, accessed March 3, 2026, https://pure.mpg.de/pubman/item/item_3594862_3/component/file_3594863/2406.08410.pdf
19. DYNAMICS OF MIXED BINARY MERGERS USING NUMERICAL RELATIVITY SIMULATIONS - Georgia Tech, accessed March 3, 2026, https://repository.gatech.edu/server/api/core/bitstreams/8d1dad2e-7b78-4d10-9fd9-26e7e499374b/content
20. arXiv:1312.7819v8 [gr-qc] 28 Dec 2017, accessed March 3, 2026, https://arxiv.org/pdf/1312.7819
21. arXiv:0908.1063v4 [gr-qc] 4 Dec 2009, accessed March 3, 2026, https://arxiv.org/pdf/0908.1063
22. Initial Data for Numerical Relativity - arXiv, accessed March 3, 2026, https://arxiv.org/pdf/gr-qc/0007085
23. Variational Principles in General Relativity, accessed March 3, 2026, https://arxiv.org/pdf/gr-qc/0205110
24. Numerical Brill-Lindquist initial data with a Schwarzschildean end at spatial infinity - MPG.PuRe, accessed March 3, 2026, https://pure.mpg.de/rest/items/item_2081443/component/file_2191907/content
25. [0908.1063] Bowen-York trumpet data and black-hole simulations - arXiv, accessed March 3, 2026, https://arxiv.org/abs/0908.1063
26. [1410.8607] Puncture Initial Data for Black-Hole Binaries with High Spins and High Boosts, accessed March 3, 2026, https://arxiv.org/abs/1410.8607
27. [0805.4192] Binary-black-hole initial data with nearly-extremal spins - arXiv, accessed March 3, 2026, https://arxiv.org/abs/0805.4192
28. The Cauchy problem and the initial data problem in effective theories of gravity - University of Cambridge, accessed March 3, 2026, https://www.repository.cam.ac.uk/bitstreams/0f7a3edd-df6c-4fde-beb7-bdab89e2789e/download
29. 3+1 Numerical Relativity - Institute for Advanced Study, accessed March 3, 2026, https://www.ias.edu/sites/default/files/NR_notes.pdf
30. Numerical evolutions of Einstein's equations, accessed March 3, 2026, https://sfb.aei.mpg.de/School04/Talks/Denis1.pdf
31. Constraint damping for the Z4c formulation of general relativity, accessed March 3, 2026, https://air.unipr.it/retrieve/handle/11381/2783920/20526/1107.5539.pdf
32. Compact binary evolutions with the Z4c formulation - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/233886024_Compact_binary_evolutions_with_the_Z4c_formulation

33. [1107.5539] Constraint damping for the Z4c formulation of general relativity - arXiv.org, accessed March 3, 2026, https://arxiv.org/abs/1107.5539
34. [1212.2901] Compact binary evolutions with the Z4c formulation - arXiv, accessed March 3, 2026, https://arxiv.org/abs/1212.2901
35. Spinning black hole in the puncture method: Numerical experiments - Semantic Scholar, accessed March 3, 2026, https://www.semanticscholar.org/paper/a7928cbe277e2fb27e23d776496249ffd3f647a3
36. Bowen-York Solutions of the Einstein Constraints with Asymptotically Flat, Hyperboloidal or Cylindrical Ends - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/325058963_Bowen-York_Solutions_of_the_Einstein_Constraints_with_Asymptotically_Flat_Hyperboloidal_or_Cylindrical_Ends
37. ED352 TOMMASO DE LORENZO - Theses.fr, accessed March 3, 2026, https://theses.fr/2018AIXM0264.pdf
38. Bondi Accretion in Trumpet Geometries - arXiv, accessed March 3, 2026, https://arxiv.org/pdf/1607.03047
39. Trumpet slices of the Schwarzschild-Tangherlini spacetime - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/47546599_Trumpet_slices_of_the_Schwarzschild-Tangherlini_spacetime
40. Critical Phenomena in Gravitational Collapse - PMC - NIH, accessed March 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5253925/
41. Critical Phenomena in Gravitational Collapse - arXiv, accessed March 3, 2026, https://arxiv.org/html/2507.07636v1
42. Critical collapse of a massive scalar field in semi-classical loop quantum gravity - arXiv.org, accessed March 3, 2026, https://arxiv.org/pdf/2601.18191
43. Critical Phenomena in Gravitational Collapse - PMC - NIH, accessed March 3, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5256106/
44. Black ring - Scholarpedia, accessed March 3, 2026, http://www.scholarpedia.org/article/Black_ring
45. Black Rings - ResearchGate, accessed March 3, 2026, https://www.researchgate.net/publication/265461670_Black_Rings
46. Numerical Relativity in Higher Dimensional Spacetimes - University of Cambridge, accessed March 3, 2026, https://www.repository.cam.ac.uk/bitstreams/8a821ac2-2341-4cbf-b957-578da5780773/download
47. Curriculum Vitae – Summary - ICREA, accessed March 3, 2026, https://www.icrea.cat/cvs/17415/roberto-emparan-garcia/
48. [1503.00721] The Black Ring is Unstable - arXiv.org, accessed March 3, 2026, https://arxiv.org/abs/1503.00721
49. [PDF] Compact binary evolutions with the Z4c formulation - Semantic Scholar, accessed March 3, 2026, https://www.semanticscholar.org/paper/Compact-binary-evolutions-with-the-Z4c-formulation-Hilditch-Bernuzzi/9a469b04a69b690ef8caf999f628f584199902ea