The Empirical Foundations of Quantum Mechanics: A Technical Analysis of Canonical Experiments and Their Conceptual Consequences

The transition from classical to quantum physics was not a sudden theoretical realization but a century-long empirical struggle, where laboratory results repeatedly violated the established frameworks of Newtonian mechanics and Maxwellian electrodynamics. This paradigm shift was characterized by a reluctant, and often grudging, abandonment of classical continuity in favor of quantization, wave-particle duality, intrinsic spin, and non-local entanglement.[1] Each of these concepts emerged as a necessary resolution to experimental "anomalies" that refused to fit the classical picture, moving from the thermodynamic puzzles of the late nineteenth century to the high-precision tests of quantum electrodynamics and satellite-based entanglement distribution in the modern era.[1]

The Old Quantum Theory: Quantization of Energy and Matter

The early twentieth century was defined by the realization that energy and physical properties are not continuous but are exchanged in discrete units. This phase of development, known as the "Old Quantum Theory," was driven by the failure of classical statistics to describe black-body radiation and the photoelectric effect.[1]

Black-body Radiation and the Planck Postulate

The formal genesis of quantum mechanics is found in the precision spectral measurements conducted at the Physikalisch-Technische Reichsanstalt (PTR) in Berlin.[1] In the late 1890s, physicists were tasked with creating standardized light sources for the burgeoning electrical industry, necessitating a deep understanding of the radiation emitted by a "black body"—an idealized object that absorbs and emits all frequencies of electromagnetic radiation in thermal equilibrium.[4]

Classical theories, specifically the Rayleigh-Jeans law, predicted that the energy density of emitted radiation would increase indefinitely with frequency, leading to the "ultraviolet catastrophe".[1] Conversely, Wien's law accurately described high-frequency radiation but failed at low frequencies.[4] Max Planck's derivation of his radiation law in 1900 served as a mathematical interpolation between these two regimes, driven by the data collected by Otto Lummer, Ernst Pringsheim, Heinrich Rubens, and Ferdinand Kurlbaum.[4]

Planck's critical insight involved the entropy of oscillators in the cavity walls. He found that a successful derivation of the observed spectrum required the assumption that these oscillators could only exchange energy in discrete increments, $E = h\nu$, where ν is the frequency and h is the newly introduced "Planck's constant".[1] Initially, Planck viewed this quantization as a "mathematical fix" rather than a physical reality, yet the accuracy of his formula—matching measurements from the infrared to the ultraviolet—forced a reconsideration of the nature of energy exchange.[4]

Percentile of Total Radiated Energy	**Wavelength at Sun Temperature (5778K)**	**Wavelength at Planet Temperature (288K)**
1%	0.296 μm	5.03 μm
10%	0.415 μm	7.62 μm
50%	0.727 μm	14.3 μm
90%	1.666 μm	32.6 μm
99.9%	8.208 μm	179 μm

The data provided in the table above illustrates the shift in spectral radiance peaks as a function of temperature, a phenomenon described by Wien's displacement law ($\lambda_{max} T = 2.898 \times 10^{-3}$ m·K) and accurately encompassed by Planck's Law.[5] The determination of h as a fundamental constant of nature set the stage for Einstein's subsequent leap into the quantization of light itself.[1]

The Photoelectric Effect: Millikan's Skeptical Confirmation

While Planck quantized the *exchange* of energy, Albert Einstein's 1905 paper proposed that light itself is composed of discrete "quanta," later called photons.[1] This hypothesis explained the photoelectric effect, where light incident on a metal surface ejects electrons.[9] Heinrich Hertz had noted in 1887 that ultraviolet light facilitates electrical discharge, and Philipp Lenard subsequently showed that the kinetic energy of the ejected electrons depended on the light's frequency, not its intensity—a result entirely inconsistent with the classical wave theory of light.[1]

Robert Millikan, a prominent experimentalist who viewed the light-quanta hypothesis as a "rash, not to say foolish hypothesis," spent ten years (1906–1916) attempting to disprove Einstein's photoelectric equation.[9] Millikan's experimental setup at the University of Chicago was a masterpiece of precision engineering; he worked with alkali metals (sodium, lithium, and potassium) inside high-vacuum chambers to prevent surface oxidation, which had plagued earlier measurements.[10]

Millikan measured the "stopping potential" V_{stop}, the negative voltage required to halt the flow of ejected electrons. According to Einstein's model, the maximum kinetic energy of an ejected electron is given by $K_{max} = h\nu - W$, where W is the work function (the energy required to remove an electron from the metal).[9] Millikan plotted the stopping potential against frequency and found a perfectly linear relationship, where the slope of the line, when multiplied by the elementary charge e, yielded Planck's constant h.[10]

Metal Target	Threshold Frequency (v0)	Calculated Work Function (W)
Sodium (Na)	$\approx 4.39 \times 10^{14}$ Hz	≈ 1.82 eV
Lithium (Li)	$\approx 5.50 \times 10^{14}$ Hz	≈ 2.28 eV

Millikan's 1916 paper, "A Direct Photoelectric Determination of Planck's h," reported a value of $h = 6.57 \times 10^{-27}$ erg-sec with an uncertainty of 0.5%.[9] Paradoxically, while Millikan provided the most rigorous experimental validation of Einstein's equation to date, he remained philosophically resistant to the concept of photons, asserting that the "semi-corpuscular"

theory of light was un-reconcilable with the well-established phenomena of diffraction and interference.[9] His integrity as an experimentalist forced him to confirm the very theory he sought to dismantle, eventually leading to his Nobel Prize in 1923.[9]

Millikan's Oil-Drop Experiment and the Elementary Charge

Concurrent with his work on the photoelectric effect, Millikan and his graduate student Harvey Fletcher performed the definitive measurement of the elementary electric charge e in 1909.[1] Prior to this, the existence of a discrete, fundamental unit of electricity was suspected but not universally accepted; J.J. Thomson had identified "corpuscles" (electrons) in cathode rays, but their charge remained a statistical average.[14]

Millikan and Fletcher used an atomized spray of oil droplets introduced into a chamber containing two parallel metal plates.[14] By ionizing the air with radiation (such as X-rays), electrons would attach to the falling droplets.[15] The researchers first measured the terminal velocity v_1 of a falling drop with the electric field turned off. At this velocity, the downward gravitational force F_g (minus buoyancy F_b) is balanced by the upward air drag F_d.[14] Applying Stokes' law: $F_d = 6\pi r\eta v_1$ where r is the radius and η is the viscosity of air.[14]

When the electric field E was turned on, the droplet's motion was altered. By adjusting the voltage V (where $E = V/d$), the researchers could suspend a droplet in mid-air or cause it to rise with a new terminal velocity v_2.[14] The charge q on the droplet was then calculated by relating the forces: $qE = m_{eff}g + 6\pi r\eta v_2$ By repeating this measurement for thousands of droplets, Millikan and Fletcher demonstrated that the charge q was always an integer multiple of a base value, which they calculated as $1.5924(17) \times 10^{-19}$ C.[14]

This value was slightly lower than the modern accepted value of $1.602176634 \times 10^{-19}$ C, a discrepancy later attributed to Millikan's use of an inaccurate value for the viscosity of air.[11] The experiment also faced historical scrutiny regarding "data selection"; physicist Gerald Holton noted in 1978 that Millikan’s notebooks contained more measurements than he published.[14] However, defenses by Allan Franklin and David Goodstein argue that Millikan only excluded measurements that were clearly compromised by technical failures or non-equilibrium conditions, and that including all data would have only increased the statistical uncertainty from 0.5% to approximately 2% without altering the central conclusion of charge quantization.[14]

Rutherford Scattering: The Discovery of the Atomic Nucleus

The internal structure of the atom was revealed through high-energy scattering experiments directed by Ernest Rutherford and performed by Hans Geiger and Ernest Marsden between 1908 and 1913.[17] At the time, J.J. Thomson's "plum-pudding" model posited that electrons were embedded in a diffuse sphere of positive charge.[17] Rutherford predicted that heavy, fast-moving alpha particles (He^{2+} ions) would pass through a thin gold foil with negligible deflection.[1]

Geiger and Marsden used a radium source to fire alpha particles at a gold foil only 0.4 μm thick (approximately 1000 atoms deep) and observed the scattering using a zinc-sulfide scintillation screen.[18] To their astonishment, while most particles passed straight through, a small fraction (about 1 in 8000) was deflected by angles greater than 90 degrees, and some bounced almost directly back toward the source.[1]

Rutherford famously remarked that this result was "as if you fired a 15-inch shell at a piece of tissue paper and it came back and hit you".[17] He concluded in 1911 that the atom's positive charge and nearly all its mass must be concentrated in a tiny, dense core—the nucleus—surrounded by mostly empty space where electrons reside.[1] Rutherford developed a mathematical scattering formula based on the Coulomb force, which predicted the number of scintillations s at an angle Φ to be proportional to $\csc^4(\Phi/2)$.[18]

Component	Rutherford Model (1911)	Thomson Model (1904)
Positive Charge	Concentrated in a central nucleus ($r \approx$ m).[19]	Uniformly distributed across the atom ($r \approx$ m).[19]
Mass Distribution	Concentrated in the nucleus.[17]	Evenly dispersed.[19]
Alpha Deflection	Occasional large-angle scattering.[18]	Only very small-angle deflections.[17]

This experimental refutation of the plum-pudding model set the stage for Niels Bohr to apply Planck's quantization to atomic orbits, creating the first quantum model of the atom in 1913.[1]

The Franck-Hertz Experiment: Atomic Energy Quantization

In 1914, James Franck and Gustav Hertz provided the first non-optical evidence of quantized energy levels within the atom.[21] They designed a vacuum tube containing mercury vapor at

approximately 100 pascals, through which they accelerated electrons from a cathode toward a grid.[21] A small retarding potential was placed between the grid and a collecting anode, ensuring that only electrons with sufficient kinetic energy could be measured as a current.[22]

As the accelerating voltage V was increased, the current initially rose. However, when V reached 4.9 V, the current dropped sharply.[21] This drop occurred because the electrons attained just enough energy to excite the mercury atoms through inelastic collisions.[22] In these collisions, the electron transfers its kinetic energy to the atom's internal degree of freedom, slowing down and becoming unable to overcome the retarding potential.[21]

As the voltage was further increased, the current rose again until it reached 9.8 V (2 x 4.9 V), where another drop occurred, signifying that a single electron had undergone two successive inelastic collisions.[22] Franck and Hertz further verified this by measuring the light emitted by the mercury vapor; the atoms, upon returning to the ground state, emitted ultraviolet light at exactly 254 nm, which corresponds to an energy of 4.9 eV according to the quantum relation $E = hc/\lambda$.[21] This "non-optical" confirmation of Bohr's energy levels was so profound that it earned Franck and Hertz the Nobel Prize in 1925.[22]

The Einstein-de Haas Effect and the Discovery of the Gyromagnetic Anomaly

In 1915, Albert Einstein and Wander de Haas conducted the only experimental work Einstein ever published, aimed at confirming Ampère's hypothesis that magnetism is caused by "molecular currents" (circulating charges).[1] They suspended a soft-iron cylinder inside a solenoid using a thin quartz fiber.[27] By reversing the magnetic field, they changed the magnetization of the iron and observed a mechanical rotation of the cylinder—a macroscopic demonstration of the conservation of angular momentum.[29]

According to classical theory, the ratio of the magnetic moment μ to the angular momentum J, known as the gyromagnetic ratio γ, should be $\gamma = e/2m$ for an orbiting electron.[28] Einstein and de Haas reported results that were "very close" to this classical prediction, seemingly confirming the orbital model of magnetism.[27] However, their reported consistency was the result of significant experimental uncertainty (quoted at 10% but later found to be much higher) and a likely expectation bias toward the classical result.[27]

Subsequent experiments by Barnett and others revealed that the actual gyromagnetic ratio for ferromagnetic materials like iron was approximately twice the classical value ($\gamma \approx e/m$).[27] This "gyromagnetic anomaly" remained unexplained for a decade until the discovery of electron spin in 1925.[27] It is now understood that in iron, 96% of the magnetization is due to the alignment of electron spins, which have a g-factor of approximately 2, while only 4% comes

from orbital motion.[27] The Einstein-de Haas effect remains a foundational observation that spin possesses real, mechanical angular momentum that can be transferred to macroscopic objects.[29]

The Stern-Gerlach Experiment: Initial Confusion and the Reality of Spin

The Stern-Gerlach experiment of 1922 is perhaps the most famous example of an experiment that provided the right result for what was initially the wrong reason.[32] Otto Stern and Walther Gerlach passed a beam of neutral silver atoms through an inhomogeneous magnetic field.[1] Classical physics predicted that the magnetic moments of the atoms would be randomly oriented, causing the beam to spread into a continuous, diffuse smear on the detector plate.[32]

Instead, the beam split into exactly two discrete components.[1] At the time, Stern and Gerlach interpreted this as a triumph for "space quantization"—the idea that the orientation of an atom's orbital angular momentum is quantized.[33] Specifically, they believed they were seeing the quantization of a Bohr-Sommerfeld orbital with $L = 1$.[32]

However, the silver atom is in a $^2S_{1/2}$ ground state, meaning its orbital angular momentum L is zero.[32] According to the theory Stern and Gerlach were testing, an $L = 0$ state should not have split at all.[32] It was only in 1925, when George Uhlenbeck and Samuel Goudsmit proposed the existence of intrinsic electron spin, that the result was correctly understood: the splitting was caused by the magnetic moment of the single unpaired electron's spin ($S = 1/2$).[1] Despite the initial theoretical confusion, the Stern-Gerlach experiment unequivocally ruled out classical trajectories and remains the archetype of quantum measurement and state preparation.[33]

Compton Scattering and the Momentum of Light

By the early 1920s, the particle-like nature of light was still contested. In 1923, Arthur Compton provided the decisive evidence by scattering X-rays off graphite.[1] Compton observed that the scattered X-rays had a longer wavelength than the incident beam, and that this wavelength shift $\Delta\lambda$ depended solely on the scattering angle θ.[37]

The classical wave theory of light could not explain this shift; it predicted that the scattered light should maintain the same frequency as the incident wave.[37] Compton explained the result by treating the interaction as an elastic collision between a photon and a "free" electron.[37] He assigned the photon a momentum $p = h/\lambda$ and applied the laws of conservation of energy and momentum from special relativity.[39]

The resulting "Compton shift" formula matched his experimental data perfectly:

$\Delta\lambda = \lambda' - \lambda = \frac{h}{m_e c}(1 - \cos\theta)$ The constant term $h/m_e c$ is known as the Compton wavelength of the electron, approximately 0.0242 Å.[38] This was the first experiment to demonstrate that light quanta carry momentum as well as energy, making the photon a "real" particle in the eyes of the physics community.[1]

Wave-Particle Duality and the Coherence of Matter

Following the success of de Broglie's 1924 hypothesis that matter, like light, should exhibit wave properties, several landmark experiments confirmed the reality of matter waves through diffraction and interference.[1]

The Davisson-Germer Experiment: Electron Diffraction

In 1927, Clinton Davisson and Lester Germer at Bell Labs were studying the scattering of low-energy electrons from a nickel surface.[1] An accidental explosion in their vacuum system led them to heat the nickel to remove oxidation, which unintentionally caused the polycrystalline surface to form large single-crystal regions.[42]

Upon resuming their measurements, they discovered sharp peaks in the intensity of reflected electrons at specific angles.[42] Specifically, they observed a strong peak at an accelerating voltage of 54 V and a scattering angle of 50 degrees.[43] Using the de Broglie wavelength for a 54 eV electron ($\lambda = h/p \approx 0.167$ nm) and applying the Bragg diffraction law to the known lattice spacing of nickel (111) planes, they found near-perfect agreement.[43] This provided the first direct confirmation that massive particles like electrons possess wave properties, a discovery for which Davisson shared the 1937 Nobel Prize with G.P. Thomson.[1]

Tonomura's Single-Electron Double-Slit Experiment

Perhaps the most visually compelling evidence for wave-particle duality was achieved by Akira Tonomura and his team at Hitachi in 1989.[1] While electron interference had been demonstrated by Jönsson in 1961 using multiple slits, Tonomura's experiment used a field-emission electron microscope and an electron biprism to perform the "single-electron" version—a thought experiment famously described by Richard Feynman as "impossible, absolutely impossible" to perform.[46]

Tonomura's apparatus was so sensitive that it could detect individual electrons with nearly 100% efficiency.[47] The emission rate was kept so low (approximately 10 electrons per second) that only one electron was present in the meter-long microscope at any given time.[47] At the start of the experiment, individual electrons appeared as random spots on the detector—behaving like particles.[47] However, as the detection events were accumulated over 20 minutes, a clear interference pattern of fringes emerged.[47] This demonstrated that even a single electron somehow "passes through both paths" of the biprism and interferes with itself,

with the final pattern reflecting the probability distribution of the wavefunction.[46]

The COW Experiment: Quantum Mechanics in a Gravitational Field

In 1975, Roberto Colella, Albert Overhauser, and Samuel Werner (COW) performed the first experiment to combine quantum mechanics and gravity.[1] They used a silicon-crystal neutron interferometer to split a beam of thermal neutrons into two paths.[50] By tilting the interferometer, one path was raised to a higher elevation in Earth's gravitational field than the other.[51]

The gravitational potential difference mgh between the two paths induced a phase shift in the neutron wavefunction.[51] The team observed a sinusoidal variation in the intensity of the emerging neutron beams as a function of the tilt angle, directly measuring the gravitationally-induced phase shift.[50] This confirmed that the Schrödinger equation accurately describes the behavior of matter waves in a non-inertial frame of reference, fundamentally linking quantum phase to the global structure of gravity.[52]

Precision Spectroscopy and the Development of QED

As experimental techniques reached higher levels of precision, the simple Bohr and Dirac models of the atom began to show discrepancies, leading to the development of Quantum Electrodynamics (QED).

Rabi Molecular Beam Resonance and the Birth of NMR

In 1938, Isidor Rabi developed the molecular beam magnetic resonance method, which allowed for the precise measurement of nuclear magnetic moments.[1] Rabi's setup passed a beam of molecules, such as lithium chloride (LiCl), through two inhomogeneous magnetic fields (A and B magnets) that would deflect the beam unless their effects were cancelled out.[54]

In a third, homogeneous field (C magnet), Rabi applied a perpendicular radiofrequency (RF) field.[54] When the RF frequency matched the Larmor precession frequency of the nuclei ($\nu = \mu H / hI$), the nuclei would flip their spin orientation.[54] This transition caused the molecules to be deflected differently by the second inhomogeneous field, resulting in a measurable drop in beam intensity at the detector.[54] Rabi's first success was with LiCl, where he reported distinct resonance peaks for both Lithium and Chlorine.[56] This technique provided the foundation for nuclear magnetic resonance (NMR), atomic clocks, and the extremely precise spectroscopy of the post-war era.[1]

The Lamb Shift and the Renormalization of the Vacuum

In 1947, Willis Lamb and Robert Retherford used microwave techniques—developed largely for radar during World War II—to investigate the fine structure of the hydrogen atom.[1] Dirac's

relativistic wave equation predicted that the $2S_{1/2}$ and $2P_{1/2}$ energy levels of hydrogen should be exactly degenerate.[60]

Lamb and Retherford's measurement revealed a small energy difference between these levels, equivalent to approximately 1058 MHz.[1] This "Lamb shift" could only be explained by the interaction of the electron with the "zero-point" fluctuations of the electromagnetic vacuum.[60] Hans Bethe soon performed the first non-relativistic calculation of this shift, which provided the stimulus for the development of QED.[60] The Lamb shift remains one of the most significant triumphs of quantum field theory, demonstrating that the "empty" vacuum is actually a dynamic medium that physically shifts the energy levels of atoms.[60]

The Anomalous Magnetic Moment and the $g-2$ Tradition

While Dirac's theory predicted that the electron's g-factor should be exactly 2, Kusch and Foley found in 1947 that it was slightly larger.[1] This deviation, known as the anomalous magnetic moment ($g-2$), is caused by the same vacuum interactions that produce the Lamb shift.[65]

Successive generations of experiments, moving from Penning-trap measurements of electrons by Hans Dehmelt to the massive superconducting storage rings used for muons at Brookhaven and Fermilab, have tested this value with increasing precision.[1] The Fermilab Muon $g-2$ experiment (E989) recently published a result of $a_\mu = 0.00116592040(54)$ with a precision of 460 parts per billion.[67] The continued disagreement (now at 4.2σ) between this experimental value and the Standard Model prediction remains one of the most exciting potential windows into "new physics" beyond the Standard Model.[65]

Gauge Topology and the Physicality of Potentials

The traditional view in classical physics was that electromagnetic potentials (A and Φ) were merely mathematical conveniences, and only the fields (E and B) were physical. This was overturned by the Aharonov-Bohm effect.

The Aharonov-Bohm Effect: Confirmation of Potential Reality

In 1959, Yakir Aharonov and David Bohm pointed out that a charged particle's phase is sensitive to the vector potential A, even if the particle never enters a region where the magnetic field B is non-zero.[1] They proposed an experiment where an electron beam is split and passed around a long solenoid.[69] Inside the solenoid, the magnetic field is strong, but outside it is zero; however, the vector potential A exists outside the solenoid and has different values on either

side.[69]

The resulting phase difference $\Delta\phi = (e/\hbar) \oint A \cdot dl$ is proportional to the magnetic flux enclosed by the paths.[70] After decades of controversy regarding flux leakage, Akira Tonomura provided the definitive proof in 1986.[1] He used a tiny, toroidal magnet made of a ferromagnetic material, encapsulated in a superconducting niobium layer.[69] The Meissner effect in the niobium completely shielded the magnetic field, ensuring that the electrons traveling inside the hole and outside the ring were in a region of exactly zero B field.[69] The observed interference shift of exactly half a fringe confirmed that the electrons "felt" the enclosed flux via the potential, proving that quantum mechanics is sensitive to the topology of the space.[1]

Nonlocality, Entanglement, and the Refutation of Realism

The most counter-intuitive aspect of quantum mechanics is "spooky action at a distance," or entanglement, which has been rigorously tested through Bell's theorem and its experimental implementations.[1]

The Evolution of Bell Tests: Closing the Loopholes

The 1935 EPR paper argued that quantum mechanics must be incomplete because it predicts correlations between distant measurements that seem to require non-local influences.[1] John Bell's 1964 theorem showed that this was a testable proposition: any local-realist theory must satisfy certain inequalities that quantum mechanics violates.[1]

Experiment	Year	Separation Distance	Loopholes Closed
Freedman-Clauser	1972	$\approx$ m	First violation.[1]
Aspect et al.	1982	$\approx$ m	Locality (via fast switching).[1]
Hanson (Delft)	2015	1.3 km	Locality and Detection simultaneously.[76]
Micius Satellite	2017	1203 km	Distance record; terrestrial fiber

			limit.[2]

The 2015 Delft experiment by Ronald Hanson used nitrogen-vacancy (NV) centers in diamonds separated by 1.3 km.[73] By using a technique called "entanglement swapping," where photons from the two diamonds are measured at a central station, the electron spins in the diamonds become entangled.[74] This setup closed the locality loophole by measuring the spins faster than light could travel between them, and the detection loophole by using high-efficiency readouts of the diamond spins.[74] The violation of Bell's inequality in this "loophole-free" experiment constitutes the definitive empirical refutation of local realism.[1]

Intercontinental Entanglement: The Micius Satellite

In 2017, Jian-Wei Pan's group used the Chinese Micius satellite to extend the range of Bell tests to over 1200 km.[1] The satellite contains a Sagnac-based source that generates polarization-entangled photon pairs at 810 nm.[3] These photons are transmitted to ground stations in Delingha and Lijiang using high-precision telescopes and multi-stage tracking systems with an accuracy of 0.41 μ rad.[78]

Despite the high channel loss (up to 82 dB) and the rotational effects of the satellite's motion, the team observed a Bell violation of $S = 2.37 \pm 0.09$.[78] This proved that entanglement can be distributed through the vacuum of space over distances where fiber-optic transmission is currently impossible due to absorption.[3] Subsequent experiments using Micius enabled an intercontinental quantum-secured video call between Beijing and Vienna, establishing the groundwork for a global quantum communication network.[1]

Hong-Ou-Mandel Interference and Two-Photon Coherence

In 1987, Chung Ki Hong, Zheyu Jeff Ou, and Leonard Mandel demonstrated a two-photon interference effect with no classical analogue.[1] When two identical (indistinguishable) photons arrive at the two input ports of a 50:50 beam splitter at the same time, they will always exit together from the same output port—never separately.[81]

This "bunching" effect occurs because the probability amplitudes for the two cases where both photons are reflected or both are transmitted cancel each other out.[81] The resulting "HOM dip" in coincidence counts is a critical tool in quantum optics for verifying the indistinguishability of single-photon sources and for implementing logic gates in optical quantum computing.[81]

The Delayed-Choice Quantum Eraser: Causality Challenged

The 1999 experiment by Kim et al. explored the deeper implications of the double-slit experiment using entangled photon pairs.[1] In this setup, an argon laser produces a photon that passes through a double-slit and hits a BBO crystal, where it is converted via spontaneous parametric down-conversion (SPDC) into two entangled photons: a "signal" photon and an

"idler" photon.[63]

The signal photon travels to detector D_0, while the idler photon travels through a series of beam splitters to detectors D_1 through D_4.[63] The detectors D_3 and D_4 provide "which-path" information, telling the experimenter which slit the photon pair came from.[63] Conversely, D_1 and D_2 are configured to "erase" this information.[63] Remarkably, the idler path is 2.5 meters longer, meaning the choice to preserve or erase path information is made approximately 8 nanoseconds *after* the signal photon has already hit its detector.[63]

The results showed that interference patterns only appeared at D_0 when correlated with the "erased" idler detections (D_1 and D_2), even though the detection at D_0 happened first.[63] This demonstrates that the mode of measurement—whether it preserves or erases information—determines the behavior of the particle (wave-like vs. particle-like) in a way that appears to transcend traditional linear causality.[63]

Macroscopic Quantum Phenomena and Collective States

The final canonical experiments prove that quantum mechanics is not limited to subatomic scales but can govern the behavior of macroscopic amounts of matter.

Superconductivity and the Josephson Effect

Superconductivity, discovered by Kamerlingh Onnes in 1911, was the first hint of macroscopic quantum behavior.[1] In 1962, Brian Josephson predicted that the collective wavefunctions of two superconductors could overlap across a thin insulating barrier, allowing Cooper pairs to tunnel through.[1] This was confirmed by Anderson and Rowell in 1963.[89] This effect is a "macroscopic manifestation" of quantum tunneling, where the entire superconducting fluid acts as a single quantum object, enabling current to flow at zero voltage.[88]

Bose-Einstein Condensation: The Superatom

Predicted in 1924, Bose-Einstein Condensation (BEC) was realized in 1995 by Eric Cornell and Carl Wieman using Rubidium-87.[1] By cooling a dilute gas of about 2000 atoms to 170 nanokelvin, they observed the atoms collapsing into the lowest energy state, characterized by a single, undulating quantum wave.[92]

Parameter	Cornell-Wieman BEC (1995)

Isotope	Rubidium-87 (^{87}Rb) [92]
Final Temperature	$\approx$ [94] nK (initially observed near 200 nK)
Number of Atoms	$\approx$ to 10^6 (in later refinements) [92]
Cooling Stages	Laser cooling $\rightarrow$ Magnetic trapping $\rightarrow$ Evaporative cooling [91]

The BEC took on an elliptical shape, confirming that it was governed by the harmonic trap potential rather than the random collisions of a classical gas.[91] Matter-wave condensation transformed from a theoretical curiosity into a laboratory reality, opening the field of ultracold atomic physics and providing a new platform for exploring fundamental quantum effects at the macroscopic scale.[1]

Conclusion

The trajectory of quantum mechanics in the laboratory has been one of increasing precision and widening scope. From Max Planck's initial attempt to explain the glow of a hot object to the distribution of entanglement across continents via satellite, each step has been forced by empirical data that defied classical explanation.[1] The consistent verification of quantization, wave-particle duality, spin, and nonlocality has established that the world is, at its foundations, irreducibly quantum.[1] These experiments have not only provided the foundation for modern technology—from MRI and transistors to superconducting qubits—but have also redefined our fundamental understanding of reality, potentials, and the nature of space and time.[55]

Works cited

1. qm_experiment_history.pdf
2. Micius Satellite for Quantum Science Experiments, accessed April 9, 2026, https://quantum.ustc.edu.cn/web/en/node/351
3. Chinese scientists set new distance record for quantum entanglement - 国务院新闻办公室, accessed April 9, 2026, http://www.scio.gov.cn/news_0/202209/t20220921_426314.html
4. max planck and the enigma of black-body radiation, accessed April 9, 2026, https://uni-tuebingen.de/fileadmin/Uni_Tuebingen/Fakultaeten/MathePhysik/Institute/IAP/Forschung/MOettel/Geburt_QM/mpiwg_notes_planck_blackbody.pdf
5. Planck's law - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Planck%27s_law
6. Introduction - The Information Philosopher, accessed April 9, 2026, https://informationphilosopher.com/solutions/scientists/planck/Planck_Kangro.pdf

7. Chronology of Milestone Events in Particle Physics - PLANCK 1900B - IHEP, accessed April 9, 2026, http://web.ihep.su/owa/dbserv/hw.part2?s_c=PLANCK+1900B
8. Planck's radiation law - PhysLab, accessed April 9, 2026, https://physlab.org/wp-content/uploads/2016/03/Phys_ref2.pdf
9. Millikan's struggle with theory - Europhysics News, accessed April 9, 2026, https://www.europhysicsnews.org/articles/epn/pdf/2000/03/epn00303.pdf
10. Robert Millikan Photoelectric Exercise - QuarkNet, accessed April 9, 2026, https://quarknet.org/sites/default/files/MillikanPhotoelectricAnalysis.pdf
11. Millikan, the First Physicist to See the Electron - OpenMind - RUA, accessed April 9, 2026, https://rua.ua.es/bitstreams/6b4614e9-8d31-47be-bcb8-501de2bfa8bb/download
12. Millikan, Einstein, and Planck: The Experiment io9 Forgot - Chad Orzel, accessed April 9, 2026, http://chadorzel.com/principles/2014/05/15/millikan-einstein-and-planck-the-experiment-io9-forgot/
13. Millikan Conducts His Oil-Drop Experiment | History | Research Starters - EBSCO, accessed April 9, 2026, https://www.ebsco.com/research-starters/history/millikan-conducts-his-oil-drop-experiment
14. Oil drop experiment - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Oil_drop_experiment
15. Millikan oil-drop experiment | Date, Summary, & Results - Britannica, accessed April 9, 2026, https://www.britannica.com/science/Millikan-oil-drop-experiment
16. Revisiting the Millikan Experiment - What's in an Oil Drop? - NaturPhilosophie, accessed April 9, 2026, https://naturphilosophie.co.uk/2013/10/18/revisiting-millikan-experiment-whats-oil-drop/
17. Geiger-Marsden Experiments - Encyclopedia.pub, accessed April 9, 2026, https://encyclopedia.pub/entry/30603
18. Rutherford scattering experiments - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Rutherford_scattering_experiments
19. Think Inside the Box: Recreating Rutherford's Scattering - arXiv, accessed April 9, 2026, https://arxiv.org/html/2507.21379v1
20. Experimental Evidence for the Structure of the Atom - Stanford University, accessed April 9, 2026, http://large.stanford.edu/courses/2017/ph241/sivulka2/
21. Franck–Hertz experiment - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Franck%E2%80%93Hertz_experiment
22. Franck-Hertz experiment with Hg-tube, accessed April 9, 2026, https://physics.iisc.ac.in/~ph212/Exp1_PH212(2019)_FrankHertz_Hgtube.pdf
23. C8 Expérience du Regel - EPFL, accessed April 9, 2026, https://www.epfl.ch/schools/sb/sph/wp-content/uploads/H4_Franck_hertz_en.pdf
24. Atomic and Nuclear Physics, accessed April 9, 2026, https://gritsan.pha.jhu.edu/2010.173.308/lab4/p6241_e.pdf
25. Franck, Hertz, and the Nature of ionization - csbsju, accessed April 9, 2026,

https://faculty.csbsju.edu/cgearhart/pubs/FH_APS_April%202013.pdf
26. accessed April 9, 2026, https://en.wikipedia.org/wiki/Franck%E2%80%93Hertz_experiment#:~:text=In%20a%20second%20paper%20presented,the%20flying%20electron%20had%20lost.
27. Einstein–de Haas effect - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Einstein%E2%80%93de_Haas_effect
28. Einstein-de Haas Effect - Wolfram Demonstrations Project, accessed April 9, 2026, https://demonstrations.wolfram.com/EinsteinDeHaasEffect/
29. Atomic spins set quantum fluid in motion - Science Tokyo, accessed April 9, 2026, https://www.isct.ac.jp/en/news/y4riu9qihv5b
30. Quantum Einstein-de Haas effect - PMC, accessed April 9, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC4855527/
31. How did gyromagnetic ratio come up before quantum mechanics, and who introduced it?, accessed April 9, 2026, https://hsm.stackexchange.com/questions/3145/how-did-gyromagnetic-ratio-come-up-before-quantum-mechanics-and-who-introduced
32. Appendix 5: Right Experiment, Wrong Theory: The Stern-Gerlach Experiment - Stanford Encyclopedia of Philosophy, accessed April 9, 2026, https://plato.stanford.edu/entries/physics-experiment/app5.html
33. Stern-Gerlach experiments: past, present, and future - BYU Physics ..., accessed April 9, 2026, https://physics.byu.edu/research/theory/docs/sgpaper.pdf
34. A Century Ago the Stern–Gerlach Experiment Ruled Unequivocally in Favor of Quantum Mechanics, accessed April 9, 2026, https://elearning.uniroma1.it/pluginfile.php/1335413/mod_resource/content/1/Israel%20Journal%20of%20Chemistry%20-%202023%20-%20Friedrich%20-%20A%20Century%20Ago%20the%20Stern%20Gerlach%20Experiment%20Ruled%20Unequivocally%20in%20Favor%20of.pdf
35. The Stern-Gerlach experiment re-examined by an experimenter - Europhysics News, accessed April 9, 2026, https://www.europhysicsnews.org/articles/epn/pdf/2019/03/epn2019503p15.pdf
36. Particle Spin and the Stern-Gerlach Experiment - Questions and Answers in MRI, accessed April 9, 2026, https://mriquestions.com/uploads/3/4/5/7/34572113/stern-gerlach.pdf
37. Compton Scattering | Definition, Equation & Formula - Study.com, accessed April 9, 2026, https://study.com/academy/lesson/compton-scattering-definition-equation-formula.html
38. 1.3 THE COMPTON EFFECT (1923) When a monochromatic beam of X-rays is scattered by an element of low atomic weight (for example c, accessed April 9, 2026, https://faculty.uobasrah.edu.iq/uploads/teaching/1632205528.pdf
39. 6.4: The Compton Effect - Physics LibreTexts, accessed April 9, 2026, https://phys.libretexts.org/Bookshelves/University_Physics/University_Physics_(OpenStax)/University_Physics_III_-_Optics_and_Modern_Physics_(OpenStax)/06%3A_Photons_and_Matter_Waves/6.04%3A_The_Compton_Effect
40. Compton scattering - Wikipedia, accessed April 9, 2026,

https://en.wikipedia.org/wiki/Compton_scattering
41. Compton Scattering | Ossila, accessed April 9, 2026, https://www.ossila.com/pages/compton-scattering
42. Davisson–Germer experiment - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Davisson%E2%80%93Germer_experiment
43. Davisson-Germer Experiment, accessed April 9, 2026, http://hyperphysics.phy-astr.gsu.edu/hbase/quantum/DavGer2.html
44. P HY S I C AL R. EV I EW, accessed April 9, 2026, https://www.ub.edu/hcub/hfq/sites/default/files/Davisson_Germer_1927%283%29.pdf
45. [1307.6049] A proper understanding of the Davisson and Germer experiments for undergraduate modern physics course - ar5iv, accessed April 9, 2026, https://ar5iv.labs.arxiv.org/html/1307.6049
46. tonomura-1989.pdf, accessed April 9, 2026, https://sites.ifi.unicamp.br/aguiar/files/2014/02/tonomura-1989.pdf
47. Double-slit experiment : Quantum Measurement : Research & Development - Hitachi Global, accessed April 9, 2026, https://www.hitachi.com/rd/research/materials/quantum/doubleslit/
48. The double-slit experiment - Physics World, accessed April 9, 2026, https://physicsworld.com/a/the-double-slit-experiment/
49. Double-slit experiment - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Double-slit_experiment
50. Quantum Phase Shifts Induced by Gravity and Rotation, accessed April 9, 2026, https://www.jps.or.jp/books/jpsjs/65A/jpsj.65sa.051.pdf
51. Neutron Interferometry: Lessons in Experimental Quantum Mechanics, Wave-Particle Duality, and Entanglement - Oxford Academic, accessed April 9, 2026, https://academic.oup.com/book/26204/book-pdf/56637552/9780191021251_web.pdf
52. Classical Underpinnings of Gravitationally Induced Quantum Interference - arXiv, accessed April 9, 2026, https://arxiv.org/pdf/gr-qc/9611037
53. Classical underpinnings of gravitationally induced quantum interference - OSTI, accessed April 9, 2026, https://www.osti.gov/biblio/567044
54. The Molecular Beam Resonance Method for Measuring Nuclear Magnetic Moments, accessed April 9, 2026, https://atomfizika.elte.hu/akos/orak/archive/atfsz/rabi/rabi.pdf
55. Discovery of Nuclear Magnetic Resonance: Rabi, Purcell, and Bloch - ACS Publications, accessed April 9, 2026, https://pubs.acs.org/doi/10.1021/bk-2020-1349.ch001
56. Discovery of NMR - Questions and Answers in MRI, accessed April 9, 2026, https://mriquestions.com/who-discovered-nmr.html
57. Rabi resonance method - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Rabi_resonance_method
58. Norman F. Ramsey - Nobel Lecture, accessed April 9, 2026, https://www.nobelprize.org/uploads/2018/06/ramsey-lecture.pdf

59. accessed April 9, 2026, https://mriquestions.com/who-discovered-nmr.html#:~:text=Rabi%20then%20passed%20a%20%22molecular,phenomenon%20%22nuclear%20magnetic%20resonance.%22
60. Lamb shift - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Lamb_shift
61. The Lamb Shift - HyperPhysics Concepts, accessed April 9, 2026, http://hyperphysics.phy-astr.gsu.edu/hbase/quantum/lamb.html
62. The Lamb shift at finite temperature and density - HawkWorks, accessed April 9, 2026, https://uhcl-ir.tdl.org/items/8568fdef-e0dc-4847-9f69-569282200331
63. Delayed-choice quantum eraser - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Delayed-choice_quantum_eraser
64. Revisiting Lamb Shift Theory through Brownian Motion of the Proton - arXiv, accessed April 9, 2026, https://arxiv.org/html/2504.05516v1
65. Muon g-2 experiment reinforces surprise result, setting up physics 'showdown', accessed April 9, 2026, https://news.uchicago.edu/story/muon-g-2-experiment-reinforces-surprise-result-setting-physics-showdown
66. Muon g-2 announces most precise measurement of the magnetic anomaly of the muon - Illinois Physics, accessed April 9, 2026, https://physics.illinois.edu/news/muon_g-2_final-result
67. The measurement of muon g – 2 at Fermilab, accessed April 9, 2026, https://lss.fnal.gov/archive/2023/conf/fermilab-conf-23-523-v.pdf
68. (PDF) The history of the muon (g-2) experiments - ResearchGate, accessed April 9, 2026, https://www.researchgate.net/publication/331240854_The_history_of_the_muon_g-2_experiments
69. Verification of the Aharonov-Bohm effect : Quantum Measurement - Hitachi Global, accessed April 9, 2026, https://www.hitachi.com/rd/research/materials/quantum/aharonov-bohm/index.html
70. Aharonov–Bohm effect - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Aharonov%E2%80%93Bohm_effect
71. The Aharonov-Bohm Effect and Tonomura et al. Experiments. Rigorous Results - arXiv, accessed April 9, 2026, https://arxiv.org/pdf/0903.2609
72. The Aharonov-Bohm effect and Tonomura experiments: Rigorous results - ResearchGate, accessed April 9, 2026, https://www.researchgate.net/publication/24164619_The_Aharonov-Bohm_effect_and_Tonomura_experiments_Rigorous_results
73. Loophole-free Bell test TU Delft crowns 80-years-old debate on nature of reality: Einsteins “spooky action” is real, accessed April 9, 2026, https://www.tudelft.nl/en/dossiers/loophole-free-bell-test-tu-delft-crowns-80-years-old-debate-on-nature-of-reality-einsteins-spooky-action-is-real
74. Physicists claim 'loophole-free' Bell-violation experiment - Physics World, accessed April 9, 2026,

https://physicsworld.com/a/physicists-claim-loophole-free-bell-violation-experiment/
75. Loophole-free Bell test at TU Delft crowns 80-years-old debate on... - Kavli Foundation, accessed April 9, 2026, https://www.kavlifoundation.org/news/loophole-free-bell-test-tu-delft-crowns-80-years-old-debate-nature-reality-einsteins
76. [2101.05370] Entanglement Swapping and Action at a Distance - arXiv, accessed April 9, 2026, https://arxiv.org/abs/2101.05370
77. Entanglement Swapping and Action at a Distance, accessed April 9, 2026, https://www.repository.cam.ac.uk/bitstreams/18e1dc86-d29e-4fa3-aca9-e923a85ca1b0/download
78. Quantum Experiments at Space Scale - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Micius_(satellite)
79. Quantum Experiments at Space Scale - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Quantum_Experiments_at_Space_Scale
80. 2017 World's Eye on China: Quantum breakthroughs with Micius satellite - YouTube, accessed April 9, 2026, https://www.youtube.com/watch?v=ywwrEAjRETE
81. Hong–Ou–Mandel effect - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Hong%E2%80%93Ou%E2%80%93Mandel_effect
82. Sample Experiments – 2-Photon-Interference by Hong, Ou & Mandel - qutools, accessed April 9, 2026, https://qutools.com/qued/qued-sample-experiments/sample-experiments-hong-ou-mandel/
83. Hong–Ou–Mandel effect - Physics Stack Exchange, accessed April 9, 2026, https://physics.stackexchange.com/questions/769820/hong-ou-mandel-effect
84. A Versatile Hong-Ou-Mandel Interference Experiment in Optical Fiber for the Undergraduate Laboratory - arXiv, accessed April 9, 2026, https://arxiv.org/html/2403.20272v1
85. (PDF) Delayed “Choice” Quantum Eraser - ResearchGate, accessed April 9, 2026, https://www.researchgate.net/publication/12307552_Delayed_Choice_Quantum_Eraser
86. Taming the Delayed Choice Quantum Eraser - PhilSci-Archive, accessed April 9, 2026, https://philsci-archive.pitt.edu/15095/1/Taming%20the%20Delayed%20Choice%20Quantum%20Eraser.pdf
87. Delayed-Choice Quantum Eraser Explained | PDF | Physics - Scribd, accessed April 9, 2026, https://www.scribd.com/document/849588079/Delayed-choice-Quantum-Eraser-Wikipedia
88. Josephson effect - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Josephson_effect
89. The Josephson Effect: Observations of Josephson's Effects - IEEE CSC, accessed April 9, 2026, https://ieeecsc.org/presentation/special/josephson-effect-observations-josephso

ns-effects
90. probable observation of the josephson superconducting tunneling effect - OSTI, accessed April 9, 2026, https://www.osti.gov/biblio/4730968
91. Appendix 3: The Discovery of Bose-Einstein Condensation: Confirmation After 70 Years, accessed April 9, 2026, https://plato.stanford.edu/archives/fall2025/entries/physics-experiment/app3.html
92. The Nobel Moment: Eric Cornell | NIST - National Institute of Standards and Technology, accessed April 9, 2026, https://www.nist.gov/nist-and-nobel/eric-cornell/nobel-moment-eric-cornell
93. REPORTS, accessed April 9, 2026, https://homepages.physik.uni-muenchen.de/~vondelft/Lehre/09qm/lec15-16-BoseEinsteinCondensation/Anderson1995.pdf
94. Bose–Einstein condensate - Wikipedia, accessed April 9, 2026, https://en.wikipedia.org/wiki/Bose%E2%80%93Einstein_condensate
95. Bose Einstein Condensates with Rubidium Atoms, accessed April 9, 2026, http://hyperphysics.phy-astr.gsu.edu/hbase/quantum/rubbec.html
96. Form Of Matter Discovered In 1995 Shows Ability To Collapse, Explode | CU Boulder Today, accessed April 9, 2026, https://www.colorado.edu/today/2001/07/17/form-matter-discovered-1995-shows-ability-collapse-explode