

Mass Spec Academy

Table of contents

Preface

The goal of this project is to create an online resource that provides comprehensive and accessible information on mass spectrometry methods used in geochronology, comprising:

- A detailed description and history of the method
- Best practices for sample preparation and analysis
- A guide to data and uncertainty analysis
- Applications, and case studies from peer-reviewed research
- Short exercises and worked solutions appropriate for graduate students.

Funding

This material is based upon work supported by the National Science Foundation under Grant Nos. EAR-2218547 and -2218544. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. Thanks to the AGeS program for its support.



Part I

Overview

What is a Mass Spectrometer?

A mass spectrometer separates atoms by their atomic mass. Scientists have long known how to separate different elements based on their chemical properties. Thousands of years ago, metallic copper was first smelted from copper ore, giving rise to the [Copper Age](#). Modern geochemical labs efficiently separate even very chemically similar elements, such as the rare earth elements, via techniques like anion exchange chromatography. These techniques separate one element from another, like separating samarium (Sm) from neodymium (Nd).

Recall that the number of protons in an atom determines the number of electrons that are needed to balance their charge. The number of electrons in an atom determines its chemical behavior — whether and how it makes chemical bonds with other atoms. But the nucleus of an atom doesn't just contain protons, it also contains neutrons. Atoms with the same number of protons but different numbers of neutrons are called isotopes. Different isotopes of the same element behave in a chemically similar manner: you can make CO_2 with ^{12}C , ^{13}C , or ^{14}C .

The relative abundances of the isotopes of an element are key to a broad array of geological processes, including the radioactivity used as a clock in geochronology, geochemical processes that fractionate radioactive parent isotopes from their radiogenic daughter products, and temperature- and environment-dependent kinetic reaction rates. To separate isotopes by their atomic mass, we need a mass spectrometer.

Separating atoms by their atomic mass is usually accomplished by first ionizing the atoms, for instance by stripping an electron off to create an ion with a +1 positive charge. The ions can then be separated according to their mass-to-charge ratio, often denoted m/z , using electric and magnetic fields.

Background Contents

1 Physics and Chemistry Refresher

1.1 Forget the Spectrometer, What is a Mass?

1.1.1 SI units for mass

The SI unit for mass is the kilogram, but samples analyzed for mass spectrometry are usually much smaller. The table below lists some typical sample sizes for geochemistry, both as SI terms with prefixes and as fractions of a gram. As we will soon learn, the number of atoms in a gram depends on the atomic mass of the atoms. The third column gives the number of atoms of that mass. It starts with atoms of mass 12 unified mass units (i.e., ^{12}C), but you can hover your slider over the blue atomic mass and drag left or right to increase or decrease its value.

Table 1.1: SI prefixes for small things.

Mass with Prefix	Mass in grams	Atoms of ^{12}C	Atoms of ^{238}U
kilogram	10^3 grams	6×10^{26}	3×10^{25}
gram	1 gram	6×10^{23}	3×10^{22}
milligram	10^{-3} grams	6×10^{20}	3×10^{19}
microgram	10^{-6} grams	6×10^{17}	3×10^{16}
nanogram	10^{-9} grams	6×10^{14}	3×10^{13}
picogram	10^{-12} grams	6×10^{11}	3×10^{10}
femtogram	10^{-15} grams	6×10^8	3×10^7
attogram	10^{-18} grams	602214	30357

1.1.2 Other units for mass

The familiar (and perhaps unfamiliar!) SI prefixes down to the attogram still don't reach a small enough value to easily compare the masses of single atoms, like ^{238}U and ^{235}U . For that, we'll need a new unit, the unified mass unit, also known as the Dalton (symbols: u or Da). The unified atomic mass unit is not in the SI, but it's commonly used in physics and chemistry for very small masses, like the mass of a single atom or molecule. It's defined as $\frac{1}{12}$ the mass of a ^{12}C atom. That's about 1.660539×10^{-27} kilograms. The equivalent unit Dalton is more widely used in the organic chemistry community.

What about the atomic mass unit, or amu? This very similar unit was used widely in the mid-twentieth century but was defined differently by physicists and chemists. It was formally abandoned in 1961, replaced by the unified atomic mass unit and the Dalton, and assigned unique unit abbreviations. However, many scientific communities still use amu to abbreviate the unified atomic mass unit. The inorganic mass spectrometry community is among them, and this textbook will use amu below.

1.1.3 Atomic masses of your favorite isotopes

The isotope ^{12}C is the only isotope with an integer mass (it has a mass of 12 amu). Other isotopes have non-integer masses, which are determined to high precision by nuclear physicists. Masses and 1σ uncertainty in parentheses are from @Wang_2021:

- ^1H has a mass of 1.007825031898(14) amu
- ^{86}Sr has a mass of 85.9092607309(91) amu
- ^{144}Nd has a mass of 143.9100873(25) amu
- ^{208}Pb has a mass of 207.9766521(13) amu
- ^{238}U has a mass of 238.0507882(20) amu

Isotopic masses aren't integers for several reasons. First, neutrons and protons don't have exactly the same mass. Neutrons are slightly heavier than protons (1.0087 vs. 1.0073 amu, respectively). But an atomic mass is different from the sum of the masses of its protons, neutrons, and much lighter electrons. The difference is the binding energy of the atom and specifically the nucleus, or the energy released by the formation of the nucleus from its constituent parts. This energy of fusion, which powers the sun and stars, can be converted to mass via Einstein's famous equation $e = mc^2$. So the combined mass of 6 protons + 6 neutrons + 6 electrons is 12.0989 amu, and the difference between that mass and the 12 amu mass of a ^{12}C atom is the energy released by putting the atom together.

The chemical energy released by forming a molecule out of atoms is small relative to the nuclear forces responsible for forming atoms, so the molecular mass of a molecule is very close to the sum of the atomic masses of its atoms. Note that two molecules with the same chemical formula might have two different molecular masses. For instance, $^{12}\text{C}^{16}\text{O}_2$ will have a different molecular mass than $^{13}\text{C}^{16}\text{O}_2$ will have a different mass than $^{12}\text{C}^{18}\text{O}^{16}\text{O}$. These three molecules, all with a different molecular mass, are called isotopologues.

Because each isotope has a slightly different mass, different atoms and/or molecules may have very similar masses. For instance, the mass of ^{40}Ar is 39.96238 amu, the mass of ^{40}Ca is 39.96259 amu, and the mass of ^{40}K is 39.96400 amu. Their proximity in mass makes these isotopes difficult (but not impossible) to separate with mass spectrometers. The more atoms a molecule has, the more opportunities isotopic substitution has to create near-overlaps. For instance, natural U is often measured by TIMS as UO_2^+ after adding a tracer containing synthetic U isotopes.




Figure 1.1: Uranium oxide (UO_2) isotopologues for a natural U sample with a ^{233}U - ^{236}U tracer added. The $^{238}\text{U}/^{236}\text{U}$ ratio is 0.5 for this sample, and the tracer $^{233}\text{U}/^{236}\text{U}$ is 1. Click to enlarge the figure.

1.2 Energy, Electricity, and Magnetism

To separate dissimilar objects like minerals or legos, one good strategy is to place them all together and then sort through and choose different elements from the pile. A chemical reaction might dissolve or precipitate one element and leave another behind. However, isotopes of the same element have nearly identical chemical behavior. Mass spectrometers don't inspect and sort a stack of static individual atoms like sorting legos, and they can't rely on chemical reactions to sort isotopes.

Instead, mass spectrometers move the atoms by first ionizing them and then manipulating the ions with electrical and magnetic forces. The resulting kinetic changes in the isotopes' motion depend on their atomic or molecular mass, which can be exploited to separate different isotopes. Once separated, the streams of ions in motion must be measured by sensitive electronic instruments. Here again, the ions' electrical properties are important.

1.2.1 Energy

$$KE = qV = \frac{1}{2}mv^2 \quad (1.1)$$

```
import numpy as np
import matplotlib.pyplot as plt

kg_per_amu = 1.66e-27
mass_amu = 238
mass_kg = mass_amu * kg_per_amu
velocity_meters_per_second = np.linspace(0, 2, 100)
kinetic_energy_joule = mass_kg * velocity_meters_per_second ** 2

fig, ax = plt.subplots()

ax.plot(velocity_meters_per_second, kinetic_energy_joule)
```



```
ax.set_xlabel('Veclocity (m/s)')
ax.set_ylabel('Kinetic Energy (J)')
ax.set_title('Stupid Plot')
fig.tight_layout()

plt.show()
```




Figure 1.2: Kinetic energy as a function of velocity for ^{238}U