

CROSS-DATASET GENERALISATION OF DEEP LEARNING FOR MALARIA CLASSIFICATION IN MICROSCOPY IMAGES

NOAH TJON SIEN KIE

TECHNICAL REPORT

CAPSTONE PROJECT, MSC DATA SCIENCE & SOCIETY, TILBURG UNIVERSITY

STUDENT NUMBER

2103939

SUPERVISOR

Dr. Sharon Ong

PANEL

to be confirmed

COURSE

Capstone Project, MSc Data Science & Society, Tilburg University

LOCATION

Tilburg University
School of Humanities and Digital Sciences
Department of Cognitive Science &
Artificial Intelligence
Tilburg, The Netherlands

DATE

11 November 2026 (Deliverable D4, Capstone Portfolio)

WORD COUNT

2,750

REPOSITORY

https://gitlab.uvt.nl/capstone/2026-fall/u927271

TECHNICAL REPORT

NOAH TJON SIEN KIE

ABSTRACT

Convolutional neural networks (CNNs) that classify red blood cells as malaria-infected score almost perfectly on the public NIH dataset but worse on images from other laboratories. This project measures that drop and tests two suspected causes: crop format and stain colour. ImageNet-pretrained VGG-16, ResNet-50 and MobileNetV2 were fine-tuned on NIH cells and tested unchanged on the external datasets BBBC041 and MP-IDB. Sensitivity fell by up to 63 percentage points, and the smallest network lost the most. Removing the background or matching stain colour at test time did not, on average, improve how well the models rank infected above uninfected cells (AUC). These two differences therefore do not explain the drop.

1 INTRODUCTION

Malaria caused an estimated 282 million cases and 610,000 deaths in 2024 (World Health Organization, 2025). Diagnosis still relies on a trained microscopist reading a stained blood smear (Rajaraman et al., 2018), and many remote areas have limited access to good-quality microscopy (World Health Organization, n.d.). Automated cell classification could help, if it stays accurate across laboratories.

The task is binary classification of single red-blood-cell images as parasitised or uninfected. Pretrained convolutional neural networks (CNNs) are trained on the NIH dataset and tested on two external datasets, BBBC041 and MP-IDB. The drop is the loss in sensitivity or specificity relative to held-out NIH patients. Stakeholders are analysed in the ELSA report.

MRQ. To what extent do pretrained CNNs trained to classify NIH single-cell images as parasitised or uninfected lose sensitivity on BBBC041 and MP-IDB, and specificity on BBBC041, relative to a patient-grouped NIH hold-out set, and which dataset differences explain this drop?

RQ1. How large is the drop from the patient-grouped NIH hold-out set, in percentage points, for MobileNetV2, ResNet-50 and VGG-16 applied without fine-tuning to BBBC041 and MP-IDB, and does the lighter architecture lose less than the heavier ones?

RQ2. What share of the RQ1 drop is recovered when test cells are matched to NIH at test time, first in crop format (background removed) and then in stain colour (Reinhard or histogram matching)?

RQ3. Does sensitivity on the pooled different-species MP-IDB cells (n = 140) differ from *P. falciparum* (n = 1,297) by more than the resampling interval, and which BBBC041 life stages account for the largest share of missed parasites?

Sections 2 to 5 cover the literature, method, results and conclusions.

2 LITERATURE REVIEW

On NIH itself, Rajaraman et al. (2018) reached 98.6% accuracy when cells were split at random but 95.9% when patients were kept apart, so random splits inflate the baseline. Zhao et al. (2020) tested an NIH-trained VGG-16 on BBBC041, where its AUC fell from 0.994 to 0.945. Loddo et al. (2022) found light and heavy CNNs similar within NIH, yet NIH-trained models averaged only 60.61% when separating rings, the earliest parasite stage, from later stages on MP-IDB. Hou et al. (2026), the state of the art and the protocol adopted here, split NIH by patient and tested on external datasets; their compact model kept 96.4% sensitivity on BBBC041 but reached only 18.0% specificity. Xu et al. (2025) showed that colour normalisation of malaria smears depends strongly on the method: accuracy at classifying parasite stages across imaging platforms was 29.34% after Reinhard colour transfer and 48.66% with BeerLaNet, their learned stain-normalisation network.

Three gaps follow. First, the drop has been reported with different metrics and decision thresholds and for different models, so its size and its dependence on architecture are unclear. Second, crop format (NIH cells cut out onto black, external cells cut as rectangles with background) and stain colour are named as suspected causes (Hou et al., 2026; Loddo et al.,

2022; Zhao et al., 2020), but no study changes one of them while keeping the model fixed. Third, none reports how much of the drop such a change recovers. This project addresses all three: three architectures under one protocol, and test images edited one difference at a time while the models stay fixed.

3 METHODOLOGY

Figure 1 shows the pipeline; sections of the Methodology Overview (MO) in the repository document each step.




Figure 1: Pipeline. F1 and F2 are the two RQ2 steps; each box names the script that implements it. Drawn by `scripts/make_mo_figures.py`.

3.1 *Data*

The training data are the NIH cell images, from the US National Library of Medicine (Rajaraman et al., 2018): 27,558 red blood cells from 200 patients in Chittagong, Bangladesh, half infected with the parasite *Plasmodium falciparum*, each cut out along its outline and placed on a black square. They were split 70/15/15 by patient, so no model is tested on a patient it was trained on; the 30 hold-out patients, never used in training, give the reference performance (MO §3.2).

Two external datasets are used for testing only. BBBC041 (Ljosa et al., 2012) contains *P. vivax* smears photographed in two batches, kept apart as site_a (79,569 cells, 2.7% parasitised) and site_b (5,917 cells, 5.1%). Ambiguous cells (446, marked "difficult") and white blood cells (103) were excluded. MP-IDB (Loddo et al., 2019) marks only parasites, so it has no uninfected cells. Each of its 1,437 parasites was cropped around the red cell containing it, like NIH's whole-cell crops, and grouped into *P. falciparum* (1,297) and three other species (140) (MO §3.1). Exploratory analysis (MO §4) found that a median 26% of each NIH crop is black frame, whereas test crops show background and neighbouring cells, and that BBBC041 cells carry three to four times more blue stain.

3.2 *Models and training*

Three CNNs pretrained on ImageNet, 1.2 million labelled photographs (Russakovsky et al., 2015), were used: MobileNetV2 (Sandler et al., 2018), ResNet-50 (He et al., 2016) and VGG-16 (Simonyan & Zisserman, 2015), with 2.2, 23.5 and 134.3 million parameters (learned weights). Each network's last layer was replaced by one output, the probability that the cell is parasitised, and all layers were fine-tuned (trained further on NIH), because features learned from everyday photographs do not fit stained cells. All three used the same settings (Appendix Table A4): flips and 90° rotations as augmentation, and early stopping, which ends training when the error on NIH validation patients stops falling and keeps the best model so far. Hyperparameters (training settings) were fixed, not tuned per model, so RQ1 compares architectures rather than tuning budgets. Random colour changes were left out on purpose: they teach a model to ignore colour, and RQ2 could then not show how much colour matters. Training is random, so each architecture was trained three times (seeds 0, 1 and 2; MO §5.1, §5.3).

3.3 *Test-time ablation (RQ2)*

RQ2 edits only the test images, so any change in the fixed models' performance comes from the edit. Step F1 removes the background with Otsu's method (Otsu, 1979), which picks the brightness threshold that best separates cell from surroundings: everything outside the cell becomes black, as in NIH crops. Its masks matched hand-drawn NIH outlines with a median Dice overlap of 0.90 (1 is perfect), while the neural model U²-Net found no cell in 21% of NIH crops (MO §5.2). Step F2 then matches stain colour to NIH in two separate variants: Reinhard transfer (Reinhard et al., 2001) matches the mean and spread of each cell's colour channels to NIH, and histogram matching maps each colour channel's whole distribution onto NIH's (reference: 1,000 NIH training cells; Figure 2).




Figure 2: The same test cells as the models see them: raw crop, background removed, and the two colour variants. Visible artefacts: Otsu keeps a touching neighbour cell (rows 1 and 3), very dark site_a pixels are taken for background (row 1), and histogram matching amplifies the noise of grey site_b cells (row 2). Drawn by `scripts/make_mo_figures.py`.

3.4 *Evaluation*

Sensitivity is the share of parasitised cells called parasitised (output above 0.5, a threshold fixed before testing), and specificity the share of uninfected cells called uninfected; drops are in percentage points (pp). They are the primary measures because BBBC041 is only 2.7 to 5.1% parasitised, so

calling every site_a cell uninfected would score 97.3% accuracy. AUC, the probability that a random parasitised cell scores above a random uninfected one, checks ranking without a threshold. F1 is secondary: it includes precision, the share of parasitised calls that are correct, which falls when infected cells are rare (49.2% of NIH hold-out cells, 2.7% of site_a), so an F1 drop would mix model failure with prevalence (MO §5.5).

Share recovered is specific to this project: the value after a step minus the raw value, divided by the raw drop. 1 means the step closes the drop, 0 that it changes nothing, and a negative value that it widens the drop. It is computed only when the raw drop's 95% interval lies above zero, since dividing by a drop near zero gives meaningless values. Because sensitivity also rises when a model simply calls more cells parasitised, every share is read with specificity and AUC.

Each value has a 95% interval from a bootstrap: the test set is redrawn 2,000 times with replacement and the middle 95% of the recomputed values kept. Whole photographs are redrawn, not cells, because cells from one photograph share its stain and focus; NIH is redrawn by patient. Training randomness is shown as the mean over three seeds with the lowest and highest seed in brackets. A finding counts as stable only if it holds in all three seeds. Models are compared on the same redrawn sets (a paired bootstrap), so luck in the draw cancels from their difference. With 24 such comparisons, about one would look real by chance at 95%, so a Bonferroni correction widens each interval to 99.8%. The external sets never inform training, early stopping, the colour reference or the threshold.

4 RESULTS, INTERPRETATION AND DISCUSSION

4.1 *RQ1: the drop and the architecture comparison*

Test set	Model	Sensitivity	Specificity	AUC
NIH hold-out	VGG-16	98.0 [97.6, 98.2]	97.0 [96.6, 97.5]	99.6 [99.6, 99.7]
	ResNet-50	97.3 [97.0, 97.6]	98.1 [97.7, 98.4]	99.6 [99.6, 99.7]
	MobileNetV2	97.2 [97.1, 97.5]	98.0 [97.8, 98.2]	99.6 [99.6, 99.7]
site_a	VGG-16	**2.1 [-1.1, 6.0]**	20.1 [13.2, 31.0]	**3.5 [2.5, 4.6]**
	ResNet-50	7.0 [-0.5, 17.9]	16.5 [11.0, 19.9]	5.6 [3.9, 7.6]
	MobileNetV2	21.8 [13.4, 31.9]	**5.9 [3.2, 8.1]**	5.7 [5.1, 6.2]
site_b	VGG-16	**7.1 [1.6, 11.2]**	1.3 [0.9, 1.5]	**2.2 [0.8, 3.5]**
	ResNet-50	30.4 [28.0, 33.9]	-0.6 [-1.3, 0.2]	2.9 [2.2, 3.4]
	MobileNetV2	59.4 [31.1, 80.2]	**-1.0 [-1.5, -0.7]**	8.0 [5.1, 12.3]
MP-IDB *P. falciparum*	VGG-16	**33.3 [28.6, 40.9]**	-	-
	ResNet-50	52.9 [44.6, 59.6]	-	-
	MobileNetV2	63.1 [42.4, 78.4]	-	-
MP-IDB other	VGG-16	**22.7 [18.2, 28.7]**	-	-
	ResNet-50	38.5 [35.6, 40.8]	-	-
	MobileNetV2	61.5 [54.6, 71.4]	-	-

Table 1: Model comparison: mean over seeds 0 to 2 [lowest, highest seed]. NIH hold-out rows give the value (%; AUC $\times$ 100); the other rows give the drop from that model's NIH value (pp). site_a and site_b are the two BBBC041 batches. Bold: smallest mean drop. Built by `scripts/make_report_tables.py`.

On the NIH hold-out the three models are close: sensitivity 97.2 to 98.0%, specificity 97.0 to 98.1% and AUC 0.996 (Table 1). Every external set lowers performance, but by how much depends on the model and the metric (Figure 3). VGG-16 loses the least sensitivity: 7.1 pp on site_b against 30.4 for ResNet-50 and 59.4 for MobileNetV2, and 33.3 and 22.7 pp on the two MP-IDB groups against 63.1 and 61.5 for MobileNetV2. MobileNetV2 never had the smallest sensitivity drop on any set in any seed, but it kept the most specificity on site_a in every seed (drop 5.9 pp against 20.1 for VGG-16). The answer to the second half of RQ1 is therefore no: the lighter architecture does not lose less. It calls fewer external cells parasitised, which trades missed parasites for fewer false alarms. This was not expected: a small network might fit less NIH-specific detail, but the larger networks kept their sensitivity better. On site_b the AUC drops (2.2 to 8.0 pp) are much smaller than the sensitivity drops, so much of that loss is a shift of the scores relative to the fixed threshold, with the ranking largely kept; MobileNetV2, with the largest AUC drop (8.0 pp), also ranks cells worse.




Figure 3: Drop against each model's own NIH hold-out value, per external set and metric (dot: mean of seeds 0 to 2; bar: lowest to highest seed). Built by `scripts/make_report_tables.py`.

4.2 *In-sample against out-of-sample performance*

Data	VGG-16	ResNet-50	MobileNetV2
Training (best epoch, augmented)	96.8 / 97.7	96.4 / 97.7	96.6 / 97.9
NIH validation (best epoch)	97.9 / 96.7	97.1 / 97.4	96.7 / 97.8
NIH hold-out	98.0 / 97.0	97.3 / 98.1	97.2 / 98.0
BBBC041 site_a	95.8 / 76.9	90.2 / 81.5	75.4 / 92.1
BBBC041 site_b	90.9 / 95.7	66.9 / 98.7	37.8 / 99.0
MP-IDB *P. falciparum*	64.7 / -	44.4 / -	34.1 / -
MP-IDB other species	75.2 / -	58.8 / -	35.7 / -

Table 2: Sensitivity / specificity (%) from the training data to the external sets, mean of seeds 0 to 2. Training values are measured on augmented batches at the epoch whose weights were kept. MP-IDB has no uninfected cells.

Training, validation and hold-out values differ by at most 1.2 pp for every architecture (Table 2), and early stopping kept the model after 5 to 12 epochs, i.e. passes over the training data (Appendix Table A3). The models did not overfit the NIH training patients; the failure appears only on other laboratories' images, where the same models fall to 34 to 75% sensitivity on MP-IDB. A patient-grouped hold-out set therefore says little about another laboratory.

4.3 *RQ2: matching crop format and stain colour*

Removing the background recovered part of the sensitivity drop for VGG-16 and ResNet-50, for example a share of 0.31 for ResNet-50 on MP-IDB *P. falciparum* and 0.43 on site_b (Appendix Table A1). It also lowered site_a specificity for all three models, and mean AUC improved for no model on either BBBC041 site (Figure 4). A likely reason is two known artefacts of the masks: Otsu sometimes keeps a touching neighbour with the cell, and very dark site_a pixels are taken for background (Figure 2), so some uninfected cells end up with dark, parasite-like fragments on black. Reinhard transfer then lowered site_a sensitivity by a further 34 to 43 pp, while histogram matching raised sensitivity to 85 to 100% by calling almost every cell parasitised: site_a specificity fell to 4.5 to 35.3%. Its apparent recovery of 85 to 106% of the MP-IDB drops is therefore not a gain. The answer to RQ2 is that little or none of the drop is recovered when ranking is the criterion: matching crop format and colour at test time moves scores across the threshold but does not help the models separate parasitised from uninfected cells.




Figure 4: Sensitivity against specificity on each BBBC041 site for each test input (mean of seeds 0 to 2). The star marks the NIH hold-out. Built by `scripts/make_report_tables.py`.

4.4 *RQ3: species and life stages*

P. falciparum, the training species, was detected less often than the other species in 8 of 9 runs (Appendix Table A2). The mean gap is −10.6 pp for VGG-16 and −14.4 pp for ResNet-50, while MobileNetV2's gap changes sign between seeds. Rings account for the largest share of missed parasites on site_b for all three models (54 to 74%) and on site_a for VGG-16 and ResNet-50; for MobileNetV2 on site_a, trophozoites (the next, larger stage) account for 53%. Rings were also missed at a higher rate than trophozoites on site_a in 9 of 9 runs and on site_b in 6 of 9. Rings are small and faint, and 95% of MP-IDB *P. falciparum* parasites are rings, which plausibly explains why the training species is detected worst. Grad-CAM (Selvaraju et al., 2017) shows which image regions drive a prediction; applied to the first training run of each model (Appendix Figure 5), it found that the parasite received 33 to 72% of the evidence while covering 11 to 27% of the crop, for missed cells as much as for detected ones. This suggests the models attend

to the parasite without recognising it, although MobileNetV2 gave empty maps for over half its MP-IDB misses.

4.5 *Comparison with the literature and across seeds*

Hou et al. (2026) found 96.4% sensitivity but 18.0% specificity on BBBC041; VGG-16 shows the same pattern in a milder form, with site_a specificity down to 65.5% in its worst seed while sensitivity stayed above 91%. Zhao et al. (2020) reported an AUC of 0.945 on BBBC041 for VGG-16; here VGG-16 reached 0.961 on site_a and 0.975 on site_b.

Seed variation is small on NIH and large on the external sets: the NIH values vary by at most 0.9 pp between seeds, while MobileNetV2's site_b sensitivity drop ranges from 31.1 to 80.2 pp. Of the 24 paired model comparisons, 18 point the same way in all three seeds, and 10 of those are also significant after Bonferroni correction in all three (`rq1_compare_seeds.csv`). The RQ1 ranking holds in every seed, but single-seed numbers do not: VGG-16's site_a specificity drop was 31.0 pp at seed 0 and 13.2 pp at seed 2. Reporting one training run, as most of the reviewed studies do, can misstate an external drop by a factor of two.

4.6 *Strengths and weaknesses*

The comparisons are fair by design: one split, identical settings, the same cells and threshold for every model and test input, paired intervals and three seeds. Every table can be rebuilt with one command, without a GPU. The weaknesses limit how far the conclusions reach. Corrections are applied only at test time, so the study cannot say whether training with colour augmentation, on masked cells or on labelled target images would remove the drop; these change the model, which RQ2 holds fixed. The fixed threshold of 0.5 mixes errors in ranking with errors in where the threshold sits, which the AUC only partly separates. The other-species group has only 140 cells, and the Grad-CAM check covers only the first training run and 100 cells per group. On BBBC041 the species and the laboratory change together, so its drop cannot be divided between them. MP-IDB has no uninfected cells, so a sensitivity gain there cannot be checked against false alarms. Colour was matched only on masked crops, so colour alone was

not tested. Training settings were fixed, so the architecture ranking holds for these settings only.

5 CONCLUSION, LIMITATIONS AND FUTURE WORK

CNNs trained on NIH single cells lose much of their sensitivity on BBBC041 and MP-IDB (up to 63 pp on average) and up to 20 pp of specificity on BBBC041, yet generalise well to unseen NIH patients. The lighter MobileNetV2 loses the most sensitivity, not the least. Matching crop format and stain colour at test time does not improve mean AUC, so these two differences, corrected this way, do not explain the drop. Rings form the largest share of misses in five of six model and site pairs, and *P. falciparum* is detected less often than other species. The contribution is a controlled test-time ablation of two suspected causes, and a share-recovered measure that exposes recovery driven by the threshold.

The results rest on one source dataset, three seeds and fixed training settings. A laboratory using MobileNetV2 at site_b would miss most infected cells (sensitivity 38%), so an NIH-trained model needs local validation before use (ELSA report). On site_b AUC falls far less than sensitivity, so the threshold could be re-set on a few labelled local cells, which the saved predictions can test without retraining. Test-time colour matching failed, so stain augmentation or a learned normaliser such as BeerLaNet (Xu et al., 2025) should be tried during training. Training data should come from several laboratories and hold more ring-stage cells.

REFERENCES

He, K., Zhang, X., Ren, S., & Sun, J. (2016). Deep residual learning for image recognition. *2016 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 770–778. https://doi.org/10.1109/CVPR.2016.90

Hou, M., He, G., Liu, Z., Huang, J., & Zou, H. (2026). MalariaNet: A microcontroller-deployable malaria-microscopy detector for point-of-care biosensing under leakage-free evaluation. *Biosensors*, *16*(7), 358. https://doi.org/10.3390/bios16070358

Ljosa, V., Sokolnicki, K. L., & Carpenter, A. E. (2012). Annotated high-throughput microscopy image sets for validation. *Nature Methods*, *9*(7), 637. https://doi.org/10.1038/nmeth.2083

Loddo, A., Di Ruberto, C., Kocher, M., & Prod'Hom, G. (2019). MP-IDB: The Malaria Parasite Image Database for image processing and analysis. *Processing and Analysis of Biomedical Information*, 57–65. https://doi.org/10.1007/978-3-030-13835-6_7

Loddo, A., Fadda, C., & Di Ruberto, C. (2022). An empirical evaluation of convolutional networks for malaria diagnosis. *Journal of Imaging*, *8*(3), 66. https://doi.org/10.3390/jimaging8030066

Otsu, N. (1979). A threshold selection method from gray-level histograms. *IEEE Transactions on Systems, Man, and Cybernetics*, *9*(1), 62–66. https://doi.org/10.1109/TSMC.1979.4310076

Rajaraman, S., Antani, S. K., Poostchi, M., Silamut, K., Hossain, M. A., Maude, R. J., Jaeger, S., & Thoma, G. R. (2018). Pre-trained convolutional neural networks as feature extractors toward improved malaria parasite detection in thin blood smear images. *PeerJ*, *6*, e4568. https://doi.org/10.7717/peerj.4568

Reinhard, E., Ashikhmin, M., Gooch, B., & Shirley, P. (2001). Color transfer between images. *IEEE Computer Graphics and Applications*, *21*(4), 34–41. https://doi.org/10.1109/38.946629

Russakovsky, O., Deng, J., Su, H., Krause, J., Satheesh, S., Ma, S., Huang, Z., Karpathy, A., Khosla, A., Bernstein, M., Berg, A. C., & Fei-Fei, L. (2015). ImageNet large scale visual recognition challenge. *International Journal of Computer Vision*, *115*(3), 211–252. https://doi.org/10.1007/s11263-015-0816-y

Sandler, M., Howard, A., Zhu, M., Zhmoginov, A., & Chen, L.-C. (2018). MobileNetV2: Inverted residuals and linear bottlenecks. *2018 IEEE/CVF Conference on Computer Vision and Pattern Recognition*, 4510–4520. https://doi.org/10.1109/CVPR.2018.00474

Selvaraju, R. R., Cogswell, M., Das, A., Vedantam, R., Parikh, D., & Batra, D. (2017). Grad-CAM: Visual explanations from deep networks via gradient-based localization. *2017 IEEE International Conference on Computer Vision (ICCV)*, 618–626. https://doi.org/10.1109/ICCV.2017.74

Simonyan, K., & Zisserman, A. (2015). Very deep convolutional networks for large-scale image recognition. *3rd International Conference on Learning Representations (ICLR 2015)*.

World Health Organization. (n.d.). *Rapid diagnostic tests for malaria*. Retrieved November 9, 2026, from https://www.who.int/teams/global-malaria-programme/case-management/diagnosis/rapid-diagnostic-tests

World Health Organization. (2025). *World malaria report 2025*. World Health Organization. Geneva. https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2025

Xu, T., Wu, Y., Tripathi, A. K., Ippolito, M. M., & Haeffele, B. D. (2025). Adaptive stain normalization for cross-domain medical histology. *Medical Image Computing and Computer Assisted Intervention – MICCAI 2025*, 24–33. https://doi.org/10.1007/978-3-032-04981-0_3

Zhao, O. S., Kolluri, N., Anand, A., Chu, N., Bhavaraju, R., Ojha, A., Tiku, S., Nguyen, D., Chen, R., Morales, A., Valliappan, D., Patel, J. P., & Nguyen, K. (2020). Convolutional neural networks to automate the screening of malaria in low-resource countries. *PeerJ*, *8*, e9674. https://doi.org/10.7717/peerj.9674

A EXTENDED RESULTS

Model	Test set	Metric	Raw	Masked	+ Reinhard	+ Hist. matching	Share recovered (seeds)
VGG-16	site_a	Sensitivity	2.1	-0.3	34.7	-1.7	0.62 (2/3)
		Specificity	20.1	32.7	13.3	89.2	-0.76 (3/3)
		AUC	3.5	3.9	14.2	8.4	-0.14 (3/3)
	site_b	Sensitivity	7.1	6.9	11.2	-1.6	0.06 (2/3)
		Specificity	1.3	5.6	11.4	94.5	n/a
		AUC	2.2	3.0	5.9	24.6	-0.38 (3/3)
	MP-IDB *P. falciparum*	Sensitivity	33.3	26.3	31.4	-1.3	0.19 (3/3)
	MP-IDB other	Sensitivity	22.7	20.3	34.1	-0.4	0.11 (3/3)
ResNet-50	site_a	Sensitivity	7.0	0.4	34.5	-2.0	1.07 (2/3)
		Specificity	16.5	40.7	20.1	93.5	-1.54 (3/3)
		AUC	5.6	5.9	19.9	21.7	-0.01 (3/3)
	site_b	Sensitivity	30.4	17.0	14.9	-0.7	0.43 (3/3)
		Specificity	-0.6	1.8	9.2	91.5	n/a
		AUC	2.9	3.3	6.5	33.9	-0.11 (3/3)
	MP-IDB *P. falciparum*	Sensitivity	52.9	36.0	42.6	-1.6	0.31 (3/3)
	MP-IDB other	Sensitivity	38.5	22.0	44.7	-2.2	0.43 (3/3)
MobileNetV2	site_a	Sensitivity	21.8	20.2	63.4	2.3	0.10 (3/3)
		Specificity	5.9	18.0	3.9	62.7	-2.20 (3/3)
		AUC	5.7	11.6	14.1	16.4	-1.02 (3/3)
	site_b	Sensitivity	59.4	50.1	41.4	12.1	0.15 (3/3)
		Specificity	-1.0	1.2	0.7	59.6	n/a
		AUC	8.0	12.5	9.3	26.8	-0.50 (3/3)
	MP-IDB *P. falciparum*	Sensitivity	63.1	63.1	59.3	4.1	0.01 (3/3)
	MP-IDB other	Sensitivity	61.5	60.1	63.4	9.8	0.03 (3/3)

Table A1: RQ2: drop (pp) per test input (Masked: background removed; the colour steps start from the masked crops), mean over seeds 0 to 2, and the share of the raw drop recovered by background removal (mean over the seeds in which it is defined). Seed ranges are in `outputs/tables/report_rq2.csv`, built by `scripts/make_report_tables.py`.

Measure	n cells	VGG-16	ResNet-50	MobileNetV2
Sensitivity, *P. falciparum* (%)	1,297	64.7	44.4	34.1
Sensitivity, other species (%)	140	75.2	58.8	35.7
Gap, *P. falciparum* minus other (pp)		-10.6	-14.4	-1.6

Measure	n cells	VGG-16	ResNet-50	MobileNetV2
... seeds where *P. falciparum* detected less		3/3	3/3	2/3
site_a rings, share of all misses (%)	353	43.6	48.7	19.8
site_a trophozoites, share of all misses (%)	1,473	28.2	41.3	53.0
site_a ring miss rate (%)	353	5.2	20.7	27.1
site_a trophozoite miss rate (%)	1,473	2.4	7.0	19.5
... seeds where rings missed more than trophozoites		3/3	3/3	3/3
site_b rings, share of all misses (%)	169	73.9	61.7	53.8
site_b trophozoites, share of all misses (%)	111	17.6	32.1	39.6
site_b ring miss rate (%)	169	11.6	36.7	60.4
site_b trophozoite miss rate (%)	111	5.1	28.8	67.6
... seeds where rings missed more than trophozoites		3/3	3/3	0/3

Table A2: RQ3: sensitivity by species group, and share of misses and miss rate by BBBC041 life stage, mean over seeds 0 to 2, with the number of seeds in which each finding holds. Seed ranges are in `outputs/tables/report_rq3.csv`.

Model	Parameters (M)	Epochs run (seeds 0, 1, 2)	Kept epoch (seeds 0, 1, 2)	Min per epoch	Training time (min)
VGG-16	134.3	11, 15, 13	8, 12, 10	3.2	42 [34, 49]
ResNet-50	23.5	9, 10, 8	6, 7, 5	1.8	16 [15, 17]
MobileNetV2	2.2	9, 10, 8	6, 7, 5	1.3	12 [11, 14]

Table A3: Model size and training effort per architecture (Colab T4 GPU).

Setting	Value
Learning rate (Adam optimiser: step size of each weight update)	0.0001
Batch size (images per weight update)	32
Maximum epochs (passes over the training data)	20
Early stopping: epochs without lower NIH validation error before training stops	3
Weight decay (penalty on large weights)	none
Input size (pixels, square)	224
Training augmentation	horizontal and vertical flips, rotations by 90°
Decision threshold (output above it = parasitised)	0.5
Started from ImageNet weights, all layers trained	yes

Table A4: Training settings, identical for all nine runs (read from the training logs).




Figure 5: Grad-CAM maps of the first training run (seed 0) of each model on example cells (Selvaraju et al., 2017): yellow marks the regions that most raise the predicted probability of parasitised. Drawn by `scripts/gradcam.py`.