

# Chapter 4

## Vectors

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus .....	2	■	■	■	■	■ ■ ■
4G ..... Geometric proofs .....	3	□	□	□	□	□ □ □

# Syllabus

### Geometric proofs using vectors (8 hours)

In this sub-topic, students will:

- ☐ prove the diagonals of a parallelogram meet at right angles if and only if it is a rhombus
- ☐ prove midpoints of the sides of a quadrilateral join to form a parallelogram
- ☐ prove the sum of the squares of the lengths of a parallelogram's diagonals is equal to the sum of the squares of the lengths of the sides
- ☐ prove an angle in a semicircle is a right angle
- ☐ prove geometric results in the plane and construct simple proofs in three dimensions.

[illegible]

## 4G Geometric proofs

Foundational Concepts												Vocabulary & Definitions											
Big Ideas & Concepts												Procedures											



**Solution:**









**Solution:**

### Example 4.7:

Prove the sum of the squares of the lengths of a parallelogram's diagonals is equal to the sum of the squares of the lengths of the sides.

**Solution:**

# Chapter 7

## Trigonometry

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus ....	12	■	■	■	■	■ ■ ■
<b>7.1 ..... Solving trigonometric equations ....</b>	13	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.1.1 ..... Solving trigonometric equations ....	13	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.1.2 ..... General solutions ....	19	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
<b>7.2 ..... Graphing and modelling with trigonometric functions ....</b>	22	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.2.1 ..... Graphs of sine and cosine functions ....	22	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.2.2 ..... Graphs of the tangent function ....	28	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.2.3 ..... Modelling with trigonometric functions ....	31	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
<b>7.3 ..... The reciprocal trigonometric functions ....</b>	33	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.3.1 ..... Calculating values of reciprocal functions ....	33	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.3.2 ..... Graphs of reciprocal trigonometric functions ....	36	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
<b>7.4 ..... The Pythagorean trigonometric identities ....</b>	40	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
<b>7.5 ..... Compound and double-angle identities ....</b>	45	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.5.1 ..... Sums and differences ....	45	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.5.2 ..... Double-angle identities ....	48	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
7.5.3 ..... Changing products to sums and differences ....	52	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
<b>7.6 ..... Transformations of trigonometric expressions ....</b>	54	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

## Syllabus

### The basic trigonometric functions (2 hours)

In this sub-topic, students will:

- ☐ find all solutions of  $f(a(x - b))$  where  $f(\theta)$  is one of  $\sin(\theta)$ ,  $\cos(\theta)$  or  $\tan(\theta)$
- ☐ sketch and graph functions with rules of the form  $y = f(a(x - b))$  where  $f(\theta)$  is one of  $\sin(\theta)$ ,  $\cos(\theta)$  or  $\tan(\theta)$ .

### The reciprocal trigonometric functions, secant, cosecant and cotangent (3 hours)

In this sub-topic, students will:

- ☐ define the reciprocal trigonometric functions, sketch their graphs, and graph simple transformations of them.

### Trigonometric identities (8 hours)

In this sub-topic, students will:

- ☐ prove and apply the Pythagorean identities
- ☐ prove and apply the angle sum, difference and double-angle identities for sines and cosines
- ☐ prove and apply the identities for products of sines and cosines expressed as sums and differences
- ☐ convert sums  $a \cos(x) + b \sin(x)$  to  $R \cos(x \pm a)$  or  $R \sin(x \pm a)$  and apply these to sketch graphs, solve equations of the form  $a \cos(x) + b \sin(x) = c$  and solve real-world problems
- ☐ use the binomial theorem to prove and apply multi-angle trigonometric identities up to  $\sin(4x)$  and  $\cos(4x)$ .

### Applications of trigonometric functions to model periodic phenomena (5 hours)

In this sub-topic, students will:

- ☐ model periodic motion using sine and cosine functions and understand the relevance of the period and amplitude of these functions in the model.

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

## 7.1 Solving trigonometric equations

### 7.1.1 Solving trigonometric equations

Foundational Concepts	Vocabulary & Definitions
	

### Example 7.1 – Solving an equation of the type $\sin(x) = c$ :

Solve the following equations for the given domain.

(a)  $\sin(x) = \frac{1}{2}, 0 \leq x \leq 2\pi$

**(b)**  $\sin(x) = -\frac{\sqrt{2}}{2}, 0 \leq x \leq 2\pi$

**Solution:**

[illegible]

### Example 7.2 – Solving an equation of the type $\cos(x) = c$ :

Solve the equation  $2 \cos(x) + \sqrt{3} = 0$ ,  $0 \leq x \leq 2\pi$ .

**Solution:**

[illegible]

**Example 7.3** – Solving an equation of the type  $\tan(x) = c$ :

Solve the equation  $\tan(x) - 1 = 0$ ,  $0^\circ \leq x \leq 360^\circ$ .

**Solution:**

**Example 7.4** – Solving equations with different domains:

Solve the following equations for the given domain.

(a)  $\tan(x) = -\sqrt{3}, -\pi \leq x \leq \pi$

**(b)**  $\sin(x) = \frac{1}{2}, -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$

**Solution:**

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

**Example 7.5** – Solving equations of the form  $\sin(ax) = c$ ,  $\cos(ax) = c$  or  $\tan(ax) = c$ :

Solve the following equations.

(a)  $\cos(2x) = \frac{1}{2}, 0 \leq x \leq 2\pi$

**(b)**  $\sin\left(\frac{x}{3}\right) = 0.6, 0 \leq x \leq 6\pi$

**Solution:**



**Example 7.6** – Solving equations of the form  $\sin(x - b) = c$ ,  $\cos(x - b) = c$  or  $\tan(x - b) = c$ :

Solve the following equations.

(a)  $\cos(x - 30^\circ) = \frac{\sqrt{2}}{2}, 0^\circ \leq x \leq 360^\circ$

**(b)**  $\sin\left(x + \frac{5\pi}{3}\right) = \frac{1}{2}, 0 \leq x \leq 2\pi$

**Solution:**

### Example 7.7 – Solving more complex equations:

Solve the equation  $\tan\left(3\left(x - \frac{\pi}{4}\right)\right) = -1$ ,  $0 \leq x \leq 2\pi$ .

**Solution:**

### 7.1.2 General solutions

Foundational Concepts												Vocabulary & Definitions											
Big Ideas & Concepts												Procedures											

**Example 7.8** – Solving equations with an unrestricted domain:

Solve these equations, leaving your answer in general form.

(a)  $\sin(x) = \frac{\sqrt{3}}{2}$

(b)  $\tan(x) = \frac{1}{\sqrt{2}}$

**Solution:**



**Example 7.9** – Solving equations with an unrestricted domain along the vertical axis:

Solve  $\cos(2x) = 0$ .

**Solution:**

## 7.2 Graphing and modelling with trigonometric functions

### 7.2.1 Graphs of sine and cosine functions

Foundational Concepts	Vocabulary & Definitions
	



**Example 7.11** – Graphing functions of the form  $f(x) = A \sin(ax)$  for  $A < 0$ :

Consider the function  $f(x) = -3 \sin(2x)$ ,  $0 \leq x \leq 2\pi$

- Determine the period, amplitude and range of the function.
- Determine the coordinates of the  $x$ -intercepts for the graph of  $y = f(x)$ .
- Determine the equation of the median line.
- Sketch the graph of the function, labelling all major points in coordinate form, including the maximum and minimum points and showing the median line.

**Solution:**



**Example 7.12** – Graphing a trigonometric function with a vertical translation:

Consider the function  $f(x) = 3 \sin(2x) + 4$ ,  $0 \leq x \leq 2\pi$ .

- Determine the amplitude, period and range of the function.
- Calculate the intercepts of  $f(x)$  with the median line.
- Sketch the graph of the function, labelling all key points in coordinate form and showing the median line.

**Solution:**



### Example 7.14 – Determining transformations of trigonometric functions:

Determine the transformations that have taken place to transform either  $f(x) = \sin(x)$  or  $f(x) = \cos(x)$  into the given function. Also list the period and amplitude of each function.


(a)  $f(x) = \cos\left(3x + \frac{\pi}{6}\right)$

**(b)**  $f(x) = -\sin(\pi x) + 2$

(c)  $f(x) = 6 \cos\left(\frac{1}{2}x - \frac{\pi}{4}\right)$

**Solution:**

### 7.2.2 Graphs of the tangent function

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures

**Example 7.15** – Graphing functions of the form  $f(x) = \tan(ax)$ :

Consider the function  $f(x) = \tan(2x)$ ,  $0 \leq x \leq 2\pi$ .

- Determine the period, equations of the asymptotes and  $x$ -intercepts of the function.
- Sketch the graph of  $y = f(x)$ .
- List the transformations that have taken place to transform  $y = \tan(x)$  into  $y = \tan(2x)$ .

**Solution:**

**Example 7.16** – Graphing functions of the form  $f(x) = \tan(a(x - b))$ :

Consider  $h(x) = \tan\left(2\left(x - \frac{\pi}{3}\right)\right)$ ,  $0 \leq x \leq \pi$ .

- (a) Sketch the graph of  $y = h(x)$ , showing the asymptotes and the coordinates of key points.
- (b) List the transformations in sequential order to produce the graph of  $y = h(x)$  from the graph of  $f(x) = \tan(x)$ .

**Solution:**








## 7.3 The reciprocal trigonometric functions

### 7.3.1 Calculating values of reciprocal functions

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures

**Example 7.18** – Calculating reciprocal trigonometric functions for exact values:

Calculate the exact value of each of the following.

(a)  $\sec(\pi)$

(b)  $\csc\left(\frac{\pi}{2}\right)$


(c)  $\text{csc}(\pi)$

(d)  $\cot\left(\frac{\pi}{3}\right)$

**Solution:**



### 7.3.2 Graphs of reciprocal trigonometric functions

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures



**Example 7.21** – Sketching reciprocal trigonometric functions of the form  $Af(x - b)$ :

Sketch the graph of  $f(x) = 2 \cot \left( x - \frac{\pi}{4} \right)$  for  $0 \leq x \leq 2\pi$ , stating the domain, range and period. Also show the asymptotes and any intercepts with the axes.

**Solution:**

**Example 7.22** – Sketching reciprocal functions of the form  $f(a(x - b)) + c$ :

Sketch the graph of  $f(x) = \csc\left(x + \frac{\pi}{4}\right) + 1$  for  $0 \leq x \leq 2\pi$ , stating the domain, range and period. Also show the asymptotes and any intercepts with the axes.

**Solution:**

## 7.4 The Pythagorean trigonometric identities

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures



**Example 7.23** – Calculating the values of trigonometric expressions using identities:

Calculate the values of the following.

(a)  $\sin(\theta)$  for  $\frac{\pi}{2} \leq \theta \leq \pi$  if  $\cos(\theta) = -\frac{7}{25}$ .

(b)  $\cos(\theta)$  for  $\pi \leq \theta \leq 2\pi$  if  $\tan(\theta) = 3$ .

**Solution:**

### Example 7.24 – Simplifying trigonometric expressions:

Simplify the following.

(a)  $\cos^2(A) + \sin^2(A) + \frac{1}{\cot^2(A)}$ .

(b)  $\frac{\sin(\theta)}{1 - \cos(\theta)} + \frac{\sin(\theta)}{1 + \cos(\theta)}$ , where  $\cos(\theta) \neq \pm 1$ .

**Solution:**

**Example 7.25** – Using trigonometric identities to solve equations:

Solve the following equations in the given domain.

(a)  $\cos^2(A) - \sin^2(A) = \frac{1}{2}, 0 \leq A \leq 2\pi.$

(b)  $\csc^2(x) - \sqrt{3} \cot(x) = 1, 0 \leq x \leq \pi.$

**Solution:**

**Example 7.26** – Proving trigonometric identities:

Prove that  $\frac{\csc(\theta)}{\tan(\theta) + \cot(\theta)} = \cos(\theta)$ .

**Solution:**

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

### Example 7.27 – Proving a more complex trigonometric identity:

Prove that  $\frac{\cos(x)}{1 - \sin(x)} = \sec(x) + \tan(x)$ .

**Solution:**

## 7.5 Compound and double-angle identities

### 7.5.1 Sums and differences

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 7.28** – Using identities to simplify trigonometric expressions:

Demonstrate that  $\sin\left(\frac{3\pi}{2} + x\right) = -\cos(x)$

**Solution:**

[illegible]

**Example 7.29** – Using a compound angle identity to determine a trigonometric expression:

**Example 7.29** Using a compound angle identity to determine a trigonometric expression.  
If  $x$  and  $y$  are acute angles such that  $\sin(x) = \frac{1}{4}$  and  $\cos(y) = \frac{3}{4}$ , determine  $\sin(x + y)$  without calculating  $x$  or  $y$ .

**Solution:**

[illegible]

**Example 7.30** – Determining the exact value of a trigonometric expression using a compound angle identity:

Determine the exact value of  $\cos(15^\circ)$ .

**Solution:**

This image shows a full page of white paper with horizontal dotted lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


**Example 7.31** – Simplifying trigonometric expressions with compound angles:

Simplify  $\sin\left(\frac{a+b}{2}\right)\cos\left(\frac{a-b}{2}\right) + \cos\left(\frac{a+b}{2}\right)\sin\left(\frac{a-b}{2}\right)$ .

**Solution:**

This image shows a full page of primary-ruled paper. It features ten sets of horizontal dotted lines, each set consisting of three dots. These lines are evenly spaced across the entire page, providing a guide for handwriting practice. The background is white, and there are no margins or other markings present.

### 7.5.2 Double-angle identities

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures



**Example 7.32** – Simplifying trigonometric expressions using double-angle identities:

Simplify  $(\sin(3x) - \cos(3x))^2$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 7.33** – Using a double-angle formula to calculate the value of trigonometric expressions:

Given that  $\tan\left(\frac{\pi}{6}\right) = \frac{1}{\sqrt{3}}$ , use a double-angle identity to demonstrate that  $\tan\left(\frac{\pi}{3}\right) = \sqrt{3}$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 7.34 – Using half-angle identities:**

Complete the following.

(a) Use the identity  $\cos(2x) = 1 - 2\sin^2(x)$ , demonstrate that  $\sin\left(\frac{A}{2}\right) = \pm\sqrt{\frac{1 - \cos(A)}{2}}$ .

(b) If  $\cos(A) = \frac{3}{5}$  for  $\frac{3\pi}{2} \leq A \leq 2\pi$ , determine the exact value of  $\sin\left(\frac{A}{2}\right)$  without calculating  $A$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 7.35 – Using double-angle identities in a proof:**

Complete the following.

(a) Prove that  $(\cos(\theta) - \sin(\theta))^2 = 1 - \sin(2\theta)$ .

(b) Prove that  $\cos(3x) = 4\cos^3(x) - 3\cos(x)$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 7.36** – Solving equations using double-angle identities:

Complete the following.

- (a) Use the expansion found in Worked Example 35 for  $\cos(3x)$  to solve the equation  $4 \cos^3(x) - 3 \cos(x) = \frac{1}{2}$  for  $0 \leq x \leq 2\pi$ .
- (b) Determine the value of  $\theta$  such that  $\sqrt{2} \sin(2\theta) = 2 \cos(\theta)$ ,  $0 \leq \theta \leq 2\pi$ .

**Solution:**

### 7.5.3 Changing products to sums and differences

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures



## 7.6 Transformations of trigonometric expressions

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 7.38** – Expressing a sum or difference in the form  $R \sin(x \pm \alpha)$ :

Express  $\sin(x) - \cos(x)$  in the form  $R \sin(x - \alpha)$ , where  $R$  is a positive constant and  $0 \leq \alpha \leq \frac{\pi}{2}$

**Solution:**

**Example 7.39** – Expressing a sum or difference in the form  $R \cos(x \pm \alpha)$  to solve an equation:

Consider the expression  $\sqrt{3}\cos(x) - \sin(x)$ .

- (a) Express  $\sqrt{3}\cos(x) - \sin(x)$  in the form  $R\cos(x + \alpha)$  where  $R > 0$  and  $0 \leq \alpha \leq \frac{\pi}{2}$ .
- (b) Solve the equation  $\sqrt{3}\cos(x) - \sin(x) = 1$ ,  $0 \leq x \leq 2\pi$ .

**Solution:**



**Example 7.40** – Expressing a sum or difference in the form  $R \sin(x \pm \alpha)$ :

In Worked Example 38, the expression  $\sin(x) - \cos(x)$  was found to be equal to  $\sqrt{2} \sin\left(x - \frac{\pi}{4}\right)$ .

Use this relationship to sketch the graph of  $y = \sin(x) - \cos(x)$ .

**Solution:**

**Example 7.41** – Solving a problem using the transformation of an expression to the form  $R \sin(x + \alpha)$ :

A rectangle  $OABC$  is formed with one vertex from the origin  $O$ , point  $A$  at  $(0, 3\cos(\theta))$  and point  $C$  at  $(4\sin(\theta), 0)$ .

- Express  $4 \sin(\theta) + 3 \cos(\theta)$  in the form  $R \sin(\theta + \alpha)$  where  $R > 0$  and  $0 \leq \theta \leq \frac{\pi}{2}$ .
- State the coordinates of  $B$  in terms of  $\theta$ .
- Express the perimeter of the rectangle in terms of  $f(\theta)$ .
- Calculate the maximum value of the perimeter of the rectangle.

**Solution:**

# Chapter 6

## Graphing functions

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus ....	60	■	■	■	■	■ ■ ■
<b>6.1 .....</b>	<b>Absolute value functions ....</b>	61	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.1.1 .....	Absolute values functions ....	61	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.1.2 .....	Graphs of the form $y =  f(x) $ and $y = f( x )$ ....	65	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b>6.2 .....</b>	<b>Reciprocal functions ....</b>	68	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
<b>6.3 .....</b>	<b>Rational functions ....</b>	72	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.3.1 .....	Sketching rational functions with linear denominators ....	72	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.3.2 .....	Sketching rational functions with quadratic denominators ....	75	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6.3.3 .....	Sketching rational functions with an oblique asymptote ....	78	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

# Syllabus


## Sketching graphs (5 hours)

In this sub-topic, students will:

- ☐ use and apply the notation  $|x|$  for the absolute value for the real number  $x$  and the graph of  $y = |x|$
- ☐ examine the relationship between the graph of  $y = f(x)$  and the graphs of  $y = \frac{1}{f(x)}$ ,  $y = |f(x)|$  and  $y = f(|x|)$
- ☐ sketch the graphs of simple rational functions where the numerator and denominator are polynomials to degree 3 without technology.

## 6.1 Absolute value functions

### 6.1.1 Absolute values functions

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures




**Example 6.3** – Sketching graphs of simple transformations of  $y = |x|$ :

Sketch the graph of the function  $y = -2|x - 3| + 4$ , identifying the coordinates of the vertex, and any intercepts with the axes.

**Solution:**

**Example 6.4** – Determining the equation of a simple absolute graph:

Determine the equation of the function drawn below.




**Solution:**

[illegible]



### 6.1.2 Graphs of the form $y = |f(x)|$ and $y = f(|x|)$


Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures

**Example 6.5** – Sketching graphs of the form  $y = |f(x)|$  and  $y = f(|x|)$ :

For the function  $f(x)$  graphed on the right, sketch the following graphs, clearly indicate the location of all intercepts and turning points.

(a)  $y = |f(x)|$

(b)  $y = f(|x|)$



**Solution:**

[illegible]






**Example 6.7** – Sketching reciprocal functions from coordinates:

A section of the graph of a function  $f(x)$  is shown with the coordinates of several points displayed.

Replicate this diagram.

Then determine the reciprocal of the  $y$ -coordinate of each point given.


Hence, sketch the corresponding section of  $y = \frac{1}{f(x)}$  on the same set of axes.



**Solution:**

[illegible]

**Example 6.8** – Sketching  $y = \frac{1}{f(x)}$  from the graph of  $y = f(x)$  when  $f(x)$  is linear:




The graph of the function  $f(x)$  is shown on the right. Sketch the graph of  $y = \frac{1}{f(x)}$  on the same set of axes, clearly identifying all asymptotes and intercepts.

**Solution:**

[illegible]

**Example 6.9** – Sketching  $y = \frac{1}{f(x)}$  from the graph of  $y = f(x)$  when  $f(x)$  contains local extrema:

The graph of the function  $f(x)$  is shown on the right. Sketch the graph of  $y = \frac{1}{f(x)}$  on the same set of axes, clearly identifying all asymptotes, intercepts, and local extrema.




**Solution:**

[illegible]

## 6.3 Rational functions

### 6.3.1 Sketching rational functions with linear denominators

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures






**Example 6.11** – Sketching functions of the form  $y = \frac{ax + b}{cx + d}$ :

Sketch the function  $y = \frac{6-3x}{2x+2}$ , clearly indicating all intercepts and asymptotes.

**Solution:**

### 6.3.2 Sketching rational functions with quadratic denominators

Foundational Concepts	Vocabulary & Definitions
	
	Procedures

Big Ideas & Concepts




**Example 6.13** – Sketching functions of the form  $y = \frac{ax^2 + bx + c}{dx^2 + ex + f}$ :

Sketch the function  $y = \frac{2x^2 + 2}{x^2 + 2x - 3}$ , clearly indicating all intercepts and asymptotes.

**Solution:**

### 6.3.3 Sketching rational functions with an oblique asymptote

Foundational Concepts	Vocabulary & Definitions
	
	Procedures

Big Ideas & Concepts

Procedures









# Chapter 11

## Techniques of integration

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus ....	82	■	■	■	■	■ ■ ■
11A Finding definite integrals and using the modulus function ....	83	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11B ..... Derivatives of inverse trigonometric functions ....	88	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11C Anti-derivatives involving inverse trigonometric functions ....	90	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11D ..... Integration by substitution ....	93	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11E ..... Definite integrals by substitution ....	98	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11F ..... Using trigonometric identities for integration ...	101	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11G ..... Partial fractions ...	104	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11H ..... Integration by parts ...	111	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
11I ..... Further techniques and miscellaneous exercises ...	115	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>



## 11A Finding definite integrals and using the modulus function

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures










## 11B Derivatives of inverse trigonometric functions

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures





## 11C Anti-derivatives involving inverse trigonometric functions

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures





## 11D Integration by substitution

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 11.8:**

Differentiating each of the following with respect to  $x$ :

(a)  $(2x^2 + 1)^5$

(b)  $\cos^3 x$

(c)  $e^{3x^2}$

we have:

(a) Let  $y = (2x^2 + 1)^5$  and  $u = 2x^2 + 1$ .

Then  $y = u^5$ ,  $\frac{dy}{du} = 5u^4$  and  $\frac{du}{dx} = 4x$ .

Then:

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= 5u^4 \cdot 4x \\ &= 20u^4 x \\ &= 20x (2x^2 + 1)^4\end{aligned}$$

(b) Let  $y = \cos^3 x$  and  $u = \cos x$ .

Then  $y = u^3$ ,  $\frac{dy}{du} = 3u^2$  and  $\frac{du}{dx} = -\sin x$ .

Then:

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= 3u^2 \cdot (-\sin x) \\ &= 3\cos^2 x \cdot (-\sin x) \\ &= -3\cos^2 x \sin x\end{aligned}$$

(c) Let  $y = e^{3x^2}$  and  $u = 3x^2$ .

Then  $y = e^u$ ,  $\frac{dy}{du} = e^u$  and  $\frac{du}{dx} = 6x$ .

Then:

$$\begin{aligned}\frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} = e^u \cdot 6x \\ &= 6xe^{3x^2}\end{aligned}$$

Note:

(a)  $\int 20x (2x^2 + 1)^4 dx = (2x^2 + 1)^5 + c$ , then letting  $h(x) = x^2 + 1$ , we have:

$$\int 5h'(x) (h(x))^4 dx = (h(x))^5 + c$$

(b)  $\int -3\cos^2 x \sin x dx = \cos^3 x + c$ , then letting  $h(x) = \cos x$ , we have:

$$\int 3h'(x) (h(x))^2 dx = (h(x))^3 + c$$

(c)  $\int 6xe^{3x^2} dx = e^{3x^2} + c$ , then letting  $h(x) = 3x^2$ , we have:

$$\int h'(x)e^{h(x)} dx = e^{h(x)} + c$$









## 11E Definite integrals by substitution

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures





## 11F Using trigonometric identities for integration

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures





## 11G Partial fractions

Foundational Concepts												Vocabulary & Definitions											
Big Ideas & Concepts												Procedures											









**Example 11.19:**

Evaluate  $\int \frac{3x + 5}{(x - 1)(x + 3)} dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 11.20:**

Evaluate  $\int \frac{x^5 + 2}{x^2 - 1} dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



**Solution:**

## 11H Integration by parts

Foundational Concepts												Vocabulary & Definitions											
Big Ideas & Concepts												Procedures											







**Example 11.25:**

Evaluate  $\int e^x \cos x \, dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 11.26:**

Evaluate  $\int_1^2 \ln x \, dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

## 11I Further techniques and miscellaneous exercises

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures



# Chapter 12

## Applications of integration

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus...	118	■	■	■	■	■ ■ ■
12A.....The fundamental theorem of calculus...	119	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
12B ..... Area of a region between two curves...	123	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
12C ..... Integration using a graphics calculator...	128	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
12D ..... Volumes of solids of revolution...	131	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
12E.....The exponential probability distribution...	135	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
12F ..... Simpson's rule...	138	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>




## 12A The fundamental theorem of calculus

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 12.1:**

The graph of  $y = \frac{1}{\sqrt{4-x^2}}$  is shown. Determine the area of the shaded region.



**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.2:**

Determine the area under the graph of  $y = \frac{6}{4+x^2}$  between  $x = -2$  and  $x = 2$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



**Example 12.3:**

Sketch the graph of  $f(x) = \sin^{-1}(2x)$ ,  $x \in [-\frac{1}{2}, \frac{1}{2}]$ . Shade the region defined by the inequalities  $0 \leq x \leq \frac{1}{2}$  and  $0 \leq y \leq f(x)$ . Determine the area of this region.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.4:**

Sketch the graph of  $y = \frac{1}{4-x^2}$ . Shade the region for the area determined by  $\int_{-1}^1 \frac{1}{4-x^2} dx$  and calculate this area.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



## 12B Area of a region between two curves

Foundational Concepts	Vocabulary & Definitions
	

**Example 12.6:**

Calculate the area of the region bounded by the parabola  $y = x^2$  and the line  $y = 2x$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.7:**

Calculate the area of the region enclosed by the curves with equations  $y = x^2 + 1$  and  $y = 4 - x^2$  and the lines  $x = -1$  and  $x = 1$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



### Example 12.9:


Calculate the area of the shaded region.



**Solution:**


### Example 12.10:

Calculate the area of the shaded region.



**Solution:**

## 12C Integration using a graphics calculator

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures







## 12D Volumes of solids of revolution

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 12.15:**

Calculate the volume of the solid of revolution formed by rotating the curve  $y = x^3$  about:

(a) the  $x$ -axis for  $0 \leq x \leq 1$

(b) the  $y$ -axis for  $0 \leq y \leq 1$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.16:**

Determine the volume of the solid of revolution when the region bounded by the graphs of  $y = 2e^{2x}$ ,  $y = 1$ ,  $x = 0$  and  $x = 1$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

**Example 12.17:**

The shaded region is rotated around the  $x$ -axis.  
Calculate the volume of the resulting solid.



**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.18:**

A solid is formed when the region bounded by the  $x$ -axis and the graph of  $y = 3 \sin(2x)$ ,  $0 \leq x \leq \frac{\pi}{2}$  is rotated around the  $x$ -axis. Determine the volume of this solid.

**Solution:**

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

**Solution:**

## 12E The exponential probability distribution

Foundational Concepts	Vocabulary & Definitions
	
Big Ideas & Concepts	Procedures

**Example 12.20:**

The time,  $X$  minutes, that a shop assistant waits before the next customer arrives is known to be exponentially distributed, with probability density function given by

$$f(x) = \begin{cases} 0.2e^{-0.2x} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

Calculate the probability that he will wait more than 8 minutes for the next customer to arrive.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.21:**

For the situation in Example 20, calculate the mean and standard deviation of the time that the shop assistant waits for a customer.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



**Example 12.22:**

The time,  $T$  minutes, that it takes a librarian to locate a book is exponentially distributed with a mean of 3 minutes. Determine:

- (a) the probability density function of  $T$
- (b) the probability that it takes her less than 2 minutes to find a book.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.23:**

For the situation in Example 20, determine the cumulative distribution function of  $X$ , and hence calculate the probability that the shop assistant waits between 5 and 10 minutes for a customer.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 12F Simpson's rule

Foundational Concepts												Vocabulary & Definitions											
Big Ideas & Concepts												Procedures											

**Example 12.24:**

Use Simpson's rule to estimate the integral  $\int_2^5 x^3 dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 12.25:**

Use Simpson's rule to estimate the integral  $\int_0^1 e^x dx$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



# Chapter 8

## Vector calculus

Section .....	Page	Notes	Worked Examples	Exercise Questions	Study Notes	Revision
..... Syllabus ...	142	■	■	■	■	■ ■ ■
8A ..... Summary of differentiation and anti-differentiation ...	143	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8B ..... Position vectors as a function of time ...	147	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8C ..... Vector calculus ...	150	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8D ..... Velocity and acceleration for motion along a curve ...	155	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8E ..... Motion in a straight line ...	160	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8F ..... Projectile motion ...	163	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
8G ..... Circular motion ...	165	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>

# Syllabus

**Vector calculus (5 hours)**

In this sub-topic, students will:

- ☐ consider position of vectors as a function of time
- ☐ derive the Cartesian equation of a path given as a vector equation in two dimensions, including circles, ellipses and hyperbolas
- ☐ differentiate and integrate a vector function with respect to time
- ☐ determine equations of motion of a particle travelling in a straight line with both constant and variable acceleration
- ☐ apply vector calculus to motion in a plane, including projectile and circular motion.

[illegible]

## 8A Summary of differentiation and anti-differentiation

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 8.1:**

Differentiate each of the following with respect to  $x$ :

(a)  $\sqrt{x} \sin x$

(b)  $\frac{x^2}{\sin x}$

(c)  $\cos(x^2 + 1)$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.2:**

Differentiate each of the following w.r.t.  $x$ :

(a)  $\tan(5x^2 + 3)$

(b)  $\tan^3 x$

(c)  $\sec^2(3x)$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



**Example 8.3:**

Determine the second derivative of each fo the following w.r.t.  $x$ :

(a)  $f(x) = 6x^4 - 4x^3 + 4x$

(b)  $y = e^x \sin x$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.4:**

Determine an anti-derivative of each of the following:

(a)  $\sin\left(3x - \frac{\pi}{4}\right)$

(b)  $e^{3x+4}$

(c)  $6x^3 - \frac{2}{x^2}$

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

### Example 8.5:

Determine  $f(x)$  for each of the following.

(a)  $f'(x) = 3x^2 + 1$ , given  $f(0) = 1$ .

(b)  $f''(x) = -9.8$ , given  $f(0) = 0$  and  $f'(0) = 3$ .

**Solution:**

## 8B Position vectors as a function of time

Foundational Concepts

Vocabulary & Definitions

Big Ideas & Concepts

Procedures

### Example 8.6:

Each of the following vector functions describes the motion of a particle by giving its position vector,  $\mathbf{r}(t)$ , at time  $t$ . For each vector function:

- i. Determine the Cartesian relation that represents the path of the particle, determine its domain and range, and sketch its graph.
- ii. Give the starting point of the particle's motion, the direction of motion and the period of motion (if applicable).

(a)  $\mathbf{r}(t) = -5 \cos\left(\frac{\pi t}{2}\right) \mathbf{i} + 5 \sin\left(\frac{\pi t}{2}\right) \mathbf{j}, t \geq 0$

**(b)**  $\mathbf{r}(t) = -5 \cos\left(\frac{\pi t}{2}\right) \mathbf{i} + 12 \sin\left(\frac{\pi t}{2}\right) \mathbf{j}, t \geq 0$

(c)  $\mathbf{r}(t) = 4 \tan\left(\frac{\pi t}{2}\right) \mathbf{i} - 3 \sec\left(\frac{\pi t}{2}\right) \mathbf{j}, t \in (1, 3)$

**Solution:**

[illegible]



## 8C Vector calculus

Foundational Concepts	Vocabulary & Definitions
Big Ideas & Concepts	Procedures

**Example 8.7:**

Let  $\mathbf{r}(t) = 20t\mathbf{i} + (15t - 5t^2)\mathbf{j}$ . Determine  $\dot{\mathbf{r}}(t)$  and  $\ddot{\mathbf{r}}(t)$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.8:**

Let  $\mathbf{r}(t) = \cos t\mathbf{i} - \sin t\mathbf{j} + 5t\mathbf{k}$ . Determine  $\dot{\mathbf{r}}(t)$  and  $\ddot{\mathbf{r}}(t)$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.9:**

Let  $\mathbf{r}(t) = t\mathbf{i} + ((t-1)^3 + 1)\mathbf{j}$ . Determine  $\dot{\mathbf{r}}(\alpha)$  and  $\ddot{\mathbf{r}}(\alpha)$  where  $\mathbf{r}(\alpha) = \mathbf{i} + \mathbf{j}$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.10:**

Let  $\mathbf{r}(t) = e^t\mathbf{i} + ((e^t - 1)^3 + 1)\mathbf{j}$ . Determine  $\dot{\mathbf{r}}(\alpha)$  and  $\ddot{\mathbf{r}}(\alpha)$  where  $\mathbf{r}(\alpha) = \mathbf{i} + \mathbf{j}$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



**Example 8.11:**

A curve is described by the vector equation  $\mathbf{r}(t) = 2 \cos t \mathbf{i} + 2 \sin t \mathbf{j}$ .

- (a) Determine  $\dot{\mathbf{r}}(t)$  and  $\ddot{\mathbf{r}}(t)$ .
- (b) Determine the gradient of the curve at the point  $(x, y)$ , where  $x = 2 \cos t$  and  $y = 3 \sin t$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.12:**

A curve is described by the vector equation  $\mathbf{r}(t) = \sec(t) \mathbf{i} + \tan(t) \mathbf{j}$ , with  $t \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ .

- (a) Determine the gradient of the curve at the point  $(x, y)$ , where  $x = \sec(t)$  and  $y = \tan(t)$ .
- (b) Determine the gradient of the curve where  $t = \frac{\pi}{4}$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.13:**

Let  $\ddot{\mathbf{r}}(t) = 10\mathbf{i} - 12\mathbf{k}$ .

- (a) Determine  $\dot{\mathbf{r}}(t)$  if  $\dot{\mathbf{r}}(0) = 30\mathbf{i} - 20\mathbf{j} + 10\mathbf{k}$ .
- (b) Determine  $\ddot{\mathbf{r}}(t)$  if additionally  $\mathbf{r}(0) = 0\mathbf{i} + 0\mathbf{j} + 2\mathbf{k}$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.14:**

Given  $\ddot{\mathbf{r}}(t) = -9.8\mathbf{j}$  with  $\mathbf{r}(0) = \mathbf{0}$  and  $\dot{\mathbf{r}} = 30\mathbf{i} + 40\mathbf{j}$ , determine  $\mathbf{r}(t)$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 8D Velocity and acceleration for motion along a curve

Foundational Concepts

Vocabulary & Definitions

Big Ideas & Concepts

Procedures

**Example 8.15:**

The position of an object is  $\mathbf{r}(t)$  metres at time  $t$  seconds, where  $\mathbf{r}(t) = e^t \mathbf{i} + \frac{2}{9}e^{2t} \mathbf{j}, t \geq 0$ .

- (a) Determine the velocity vector for time  $t$ .
- (b) Determine the acceleration vector for time  $t$ .
- (c) Determine the speed at time  $t$ .

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.16:**

The position vector of a particle at time  $t$  is given by  $\mathbf{r}(t) = (2t - t^2) \mathbf{i} + (t^2 - 3t) \mathbf{j} + 2t \mathbf{k}$ , where  $t \geq 0$ .

- (a) Determine the velocity of the particle at time  $t$ .
- (b) Determine the speed of the particle at time  $t$ .
- (c) Determine the minimum speed of the particle.

**Solution:**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Example 8.17:**

The position of a projectile at time  $t$  is given by  $\mathbf{r}(t) = 400t\mathbf{i} + (500t - 5t^2)\mathbf{j}$ , for  $t \geq 0$ , where  $\mathbf{i}$  is the unit vector in a horizontal direction and  $\mathbf{j}$  is a unit vector vertically up. The projectile is fired from a point on the ground.

- Determine the time taken to reach the ground again.
- Determine the speed at which the projectile hits the ground.
- Determine the maximum height of the projectile.
- Determine the initial speed of the projectile.

**Solution:**

[illegible]

**Example 8.18:**

The position vector of a particle at time  $t$  is given by  $\mathbf{r}(t) = 2 \sin(2t) \mathbf{i} + \cos(2t) \mathbf{j} + 2t \mathbf{k}$ , where  $t \geq 0$ .

- Determine the velocity at time  $t$ .
- Determine the speed of the particle at time  $t$ .
- Determine the maximum speed.
- Determine the minimum speed.

**Solution:**

**Example 8.19:**

The position vectors at time  $t \geq 0$ , of particles  $A$  and  $B$  are given by

$$\mathbf{r}_A(t) = (t^3 - 9t + 8) \mathbf{i} + t^2 \mathbf{j}$$

$$\mathbf{r}_B(t) = (2 - t^2) \mathbf{i} + (3t - 2) \mathbf{j}$$

Prove that  $A$  and  $B$  collide while travelling at the same speed but at right angles to each other.

**Solution:**

## 8E Motion in a straight line

Foundational Concepts

Vocabulary & Definitions

Big Ideas & Concepts

Procedures



**Example 8.20:**

A particle moves along a straight line such that its position vector,  $\mathbf{r}(t)$  cm, at time  $t$  seconds is given by  $\mathbf{r}(t) = (3t - t^3) \mathbf{i}$ , for  $t \geq 0$ .

- (a) Determine its initial position.
- (b) Determine its position when  $t = 2$ .
- (c) Determine its initial velocity.
- (d) Determine its velocity when  $t = 2$ .
- (e) Determine its speed when  $t = 2$ .
- (f) Determine when and where the velocity is zero.

**Solution:**

**Example 8.21:**

A cricket ball is projected vertically upwards from ground level with an initial speed of 15 m/s. Take the origin  $O$  to be the point of projection and take  $\mathbf{j}$  to be the unit vector vertically up, where the unit of distance is metres.

Relative to this frame of reference, let  $\mathbf{r}(t)$  m be the position of the ball at time  $t$  seconds. Then  $\ddot{\mathbf{r}} = -g\mathbf{j}$ , where  $g$  m/s<sup>2</sup> is the magnitude of acceleration due to gravity ( $g \approx 9.8$ ).

- (a) Determine an expression for  $\dot{\mathbf{r}}(t)$ . (c) Determine the max height reached by the ball.
- (b) Determine an expression for  $\mathbf{r}(t)$ . (d) Determine when the ball returns to ground level.

**Solution:**



**Example 8.22:**

A particle is projected from a point on horizontal ground with speed 50 m/s at an angle of  $30^\circ$  to the horizontal. Let  $\mathbf{i}$  and  $\mathbf{j}$  be unit vectors in the horizontal ( $x$ ) and vertical ( $y$ ) directions respectively. Neglect air resistance.

- (a) Determine the initial velocity vector. (c) Determine the position vector at time  $t$  seconds.
- (b) Determine the velocity vector at time  $t$  seconds. (d) Determine the Cartesian equation of the path.

**Solution:**

## 8G Circular motion

Foundational Concepts

Vocabulary & Definitions

Big Ideas & Concepts

Procedures

**Example 8.23:**

A particle is moving around a circle of radius 3 m with a constant speed of 2 m/s. It is known that  $\theta = 0$  at time  $t = 0$ .

- Determine the angular velocity of the particle.
- Determine the position of the particle at time  $t = \pi$  seconds.
- Determine the velocity of the particle at time  $t = \pi$  seconds.
- Determine the acceleration of the particle at time  $t = \pi$  seconds.

**Solution:**

**Example 8.24:**

A particle moves at a constant speed of 8 m/s around a circle with radius of 4 m. Assume that  $\theta = 0$  when  $t = 0$ .

- Determine the position of the particle, relative to the centre of the circle, at time  $t$  seconds.
- Determine the velocity of the particle at time  $t$  seconds.
- Determine the acceleration of the particle at time  $t$  seconds.

**Solution:**

