

ESTIMATION OF NEURAL CONNECTIONS FROM MULTIPLE SPIKE TRAINS

GRAPH STRUCTURE INFERENCE WITH NUISANCE INPUTS

Noboru Murata

July 30, 2021

Waseda University

1. Introduction
2. Problem Formulation
3. Numerical Examples
4. Concluding Remarks

INTRODUCTION

estimating neural connections:


- understand functions of biological systems
- investigate learning/adaptation mechanisms

typical methods for measuring brain activities:

- fMRI (functional magnetic resonance imaging)
- MEG (magnetoencephalography)
- EEG (electroencephalography)
- TPE (two-photon excitation microscopy)
- multi-electrode recording

different resolutions in:


- time (oxygen consumption - neuron firing)
- space (brain mapping - synaptic connections)



by courtesy of Dr. Tatsuno
at University of Lethbridge

activities of individual neurons:


- multiple neurons
(tens - hundreds)
- long term measurement
(several hours - several days)



rearranged as binary sequence
indicating states of neurons:


- 0: resting
- 1: firing

multi-variate binary time series
contains information of neural
interactions



mathematical representation -
directed graph

- node: neuron
- edge: synaptic connection



objective

estimate weights of edges
from binary time series at nodes


typical methods for analysis:

- pair-wise:
 - cross-correlation (e.g. Wilson and McNaughton 1994)
 - joint peri-stimulus time histogram (e.g. Ito and Tsuji 2000)
- graph-based:
 - sparse inverse covariance matrix (e.g. Friedman, Hastie, and Tibshirani 2008)
 - digraph Laplacian (e.g. Noda et al. 2014)
- higher-order:
 - information geometric measure (e.g. Nakahara and Amari 2002; Tatsuno, Fellous, and Amari 2009)
 - Granger causality (e.g. Kim et al. 2011)

- pseudo correlation caused by higher-order effects
- influence from unobserved neurons
- directed excitatory/inhibitory connections

correlation coefficient:

statistics for analyzing relation of two random variables




connections

- no direct relation exists
- two nodes are connected with the same node


correlation coefficient:

statistics for analyzing relation of two random variables



connections

- no direct relation exists
- two nodes are connected with the same node




pseudo-correlation

- spurious relation appears


correlation coefficient:

statistics for analyzing relation of two random variables



connections

- no direct relation exists
- two nodes are connected with the same node




pseudo-correlation

- spurious relation appears

pseudo correlation



a common problem in complex network analysis

delayed correlation coefficient:
statistics for analyzing time series / dynamical systems



- appropriate intervals have to be considered
- information propagates multiple paths
- spurious relation appears


delayed correlation coefficient:
statistics for analyzing time series / dynamical systems




- appropriate intervals have to be considered
- information propagates multiple paths
- spurious relation appears

- consider short intervals?


delayed correlation coefficient:
statistics for analyzing time series / dynamical systems



- appropriate intervals have to be considered
- information propagates multiple paths
- spurious relation appears



- consider short intervals?



- spike intervals are random

a mathematical framework for treating:

- pseudo correlation caused by higher-order effects
- influence from unobserved neurons
- directed excitatory/inhibitory connections

a mathematical framework for treating:

- pseudo correlation caused by higher-order effects
- influence from unobserved neurons
- directed excitatory/inhibitory connections

main contribution

solve those problems with simple mathematical tricks

PROBLEM FORMULATION

indices:

- $i \in \{1, 2, \dots, N\}$: index of neurons
- $t \in \{1, 2, \dots, T\}$: discrete time of measurement
- $t_\Delta = [t - \Delta, \dots, t - 1]$: interval for delayed correlation

states:

- $X_i(t) \in \{0, 1\}$: state of neuron i at time t
- $X_i[t_\Delta] \in \{0, 1\}$: state of neuron i in interval t_Δ
- $U_i(t) \in \mathbb{R}$: internal state of neuron i at time t

connections:


- $w_{ij} \in \mathbb{R}$: synaptic connection from neuron j to neuron i
- $\lambda_{ij} \in \mathbb{R}$: pseudo connection from neuron j to neuron i

weighted sum of inputs from unobserved/observed neurons:

$$U_i(t) = B_i(t) + \sum_{j=1}^N \lambda_{ij} X_j[t_\Delta],$$

$B_i(t)$: nuisance inputs from unobserved neurons

λ_{ij} : **pseudo connection** including undirect paths



remarks

- signal from neuron j has several paths
- λ_{ij} includes direct and undirect connections

stochastic dependency on internal state:


$$\Pr(X_i(t) = 1) = \Phi_{\sigma^2}(U_i(t)),$$

Φ_{σ^2} : cdf of $\mathcal{N}(0, \sigma^2)$.

model assumption

- we assume a probit model
- Φ_{σ^2} is the integral of

$$\phi_{\sigma^2}(z) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{z^2}{2\sigma^2}\right)$$



internal state:

$$U_i(t) = B_i(t) + \sum_{j=1}^N \lambda_{ij} X_j[t_\Delta],$$

$B_i(t)$: nuisance inputs,

λ_{ij} : pseudo connection.

neuron firing:

$$\Pr(X_i(t) = 1) = \Phi_{\sigma^2}(U_i(t)),$$


$$\phi_{\sigma^2}(z) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{z^2}{2\sigma^2}\right),$$

Φ_{σ^2} : cdf of $\mathcal{N}(0, \sigma^2)$, integral of ϕ_{σ^2} .

first step

remove nuisance input B and estimate pseudo connection λ

$$U_i(t) = B_i(t) + \sum_{j=1}^N \lambda_{ij} X_j[t_\Delta].$$



Theorem

Let X and Y be independent random variables. For any function g , we have

$$\mathbb{E}[g(X + Y)] = \mathbb{E}[h(X + \mathbb{E}[Y])],$$

where f_Y is the pdf of Y and

$$f_Y^-(x) = f_Y(\mathbb{E}[Y] - x),$$


$$h = g * f_Y^-.$$

A special case is discussed in Hyvärinen 2002.

Corollary

If the function g is Φ_{σ^2} and random variable X is constant value x , and probability density function f_Y is Gaussian with mean $\mathbb{E}[Y]$ and variance τ^2 , we have

$$\mathbb{E}[\Phi_{\sigma^2}(x + Y)] = \Phi_{\sigma^2 + \tau^2}(x + \mathbb{E}[Y]).$$



consider the case of $X_j[t_\Delta] = 1$:

$$\begin{aligned} U_i(t \mid X_j[t_\Delta] = 1) &= B_i(t) + \lambda_{ij}X_j[t_\Delta] + \sum_{k \neq j} \lambda_{ik}X_k[t_\Delta] \\ &= \lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta] = 1). \end{aligned}$$

consider the case of $X_j[t_\Delta] = 1$:

$$\begin{aligned} U_i(t \mid X_j[t_\Delta] = 1) &= B_i(t) + \lambda_{ij}X_j[t_\Delta] + \sum_{k \neq j} \lambda_{ik}X_k[t_\Delta] \\ &= \lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta] = 1). \end{aligned}$$

apply the corollary for calculating conditional expectation:

$$\begin{aligned} \mathbb{E}[X_i(t) \mid X_j[t_\Delta] = 1] &= \mathbb{E}[\Phi_{\sigma^2}(U_i(t \mid X_j[t_\Delta] = 1))] \\ &= \mathbb{E}[\Phi_{\sigma^2}(\lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta] = 1))] \\ &= \Phi_{\rho^2}(\lambda_{ij} + \bar{C}_{ij}), \end{aligned}$$

where we assume $C_{ij} \sim \mathcal{N}(\bar{C}_{ij}, \tau^2)$ and $\rho^2 = \sigma^2 + \tau^2$.

for binary random variables,

$$\mathbb{E}[X_i(t) \mid X_j[t_\Delta]=1] = \Pr(X_i(t)=1 \mid X_j[t_\Delta]=1).$$

holds, therefore, obtain:

$$\begin{aligned} \Phi_{\rho^2}(\lambda_{ij} + \bar{C}_{ij}) &= \Pr(X_i(t) = 1 \mid X_j[t_\Delta]=1), \\ \Leftrightarrow \lambda_{ij} + \bar{C}_{ij} &= \rho \cdot \Phi_1^{-1}(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=1)). \end{aligned}$$

consider the both cases of $X_j[t_\Delta] = 1$ and $X_j[t_\Delta] = 0$:

$$U_i(t \mid X_j[t_\Delta]=1) = \lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta]=1),$$

$$U_i(t \mid X_j[t_\Delta]=0) = C_{ij}(t \mid X_j[t_\Delta]=0).$$

consider the both cases of $X_j[t_\Delta] = 1$ and $X_j[t_\Delta] = 0$:

$$U_i(t \mid X_j[t_\Delta] = 1) = \lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta] = 1),$$

$$U_i(t \mid X_j[t_\Delta] = 0) = C_{ij}(t \mid X_j[t_\Delta] = 0).$$

assumption

$$C_{ij}(t \mid X_j[t_\Delta] = 1), C_{ij}(t \mid X_j[t_\Delta] = 0) \sim \mathcal{N}(\bar{C}_{ij}, \tau^2)$$

consider the both cases of $X_j[t_\Delta] = 1$ and $X_j[t_\Delta] = 0$:

$$U_i(t \mid X_j[t_\Delta]=1) = \lambda_{ij} + C_{ij}(t \mid X_j[t_\Delta]=1),$$

$$U_i(t \mid X_j[t_\Delta]=0) = C_{ij}(t \mid X_j[t_\Delta]=0).$$

assumption

$$C_{ij}(t \mid X_j[t_\Delta]=1), C_{ij}(t \mid X_j[t_\Delta]=0) \sim \mathcal{N}(\bar{C}_{ij}, \tau^2)$$

then obtain:

$$\lambda_{ij} + \bar{C}_{ij} = \rho \cdot \Phi_1^{-1}(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=1)),$$

$$\bar{C}_{ij} = \rho \cdot \Phi_1^{-1}(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=0)).$$

estimator of pseudo connection:

$$\lambda_{ij} = \rho \left\{ \Phi_1^{-1} \left(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=1) \right) - \Phi_1^{-1} \left(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=0) \right) \right\}.$$

empirical estimates of conditional probability:

$$\Pr(X_i(t)=1 \mid X_j[t_\Delta]=1) = \frac{1}{Z} \sum_t X_i(t \mid X_j[t_\Delta]=1),$$

$$\Pr(X_i(t)=1 \mid X_j[t_\Delta]=0) = \frac{1}{Z'} \sum_t X_i(t \mid X_j[t_\Delta]=0).$$


second step

decompose pseudo connections λ with direct connections w :

$$\lambda_{ij} = \begin{array}{c} \text{---} j \text{---} i \end{array} + \begin{array}{c} \text{---} j \text{---} k \text{---} i \end{array} + \begin{array}{c} \text{---} j \text{---} l \text{---} k \text{---} i \end{array} + \begin{array}{c} \text{---} j \text{---} m \text{---} l \text{---} k \text{---} i \end{array} + \dots$$

second step

decompose pseudo connections λ with direct connections w :



consider an expansion with appropriate δ, δ' (delay time)

$$\begin{aligned} \lambda_{ij} &= w_{ij} \\ &+ \sum_k w_{ik} \Pr(X_k(t-\delta) = 1 \mid X_j(t-\delta') = 1) \\ &+ (\text{higher order terms}). \end{aligned}$$

introducing a virtual probability with an appropriate interval t_δ

$$\theta_{ij} = \Pr(X_i(t)=1 \mid X_j[t_\delta]=1),$$

obtain an expansion of λ as:

$$\lambda_{ij} = w_{ij} + \sum_k w_{ik} \theta_{kj} + \sum_{k,l} w_{ik} \theta_{kl} \theta_{lj} + \sum_{k,l,m} w_{ik} \theta_{kl} \theta_{lm} \theta_{mj} + \dots$$

introducing a virtual probability with an appropriate interval t_δ

$$\theta_{ij} = \Pr(X_i(t)=1 \mid X_j[t_\delta]=1),$$

obtain an expansion of λ as:

$$\lambda_{ij} = w_{ij} + \sum_k w_{ik} \theta_{kj} + \sum_{k,l} w_{ik} \theta_{kl} \theta_{lj} + \sum_{k,l,m} w_{ik} \theta_{kl} \theta_{lm} \theta_{mj} + \dots$$

this expression gives a simple matrix form:

$$\begin{aligned} \Lambda &= W(I + \Theta + \Theta^2 + \Theta^3 + \dots) \\ &= W(I - \Theta)^{-1}, \end{aligned}$$

▷ Neumann series

where $W = (w_{ij})$ and $\Theta = (\theta_{ij})$.

relation between θ and w :

$$\begin{aligned}\theta_{ij} &= \Pr(X_i(t)=1 \mid X_j[t_\delta]=1) \\ &= \mathbb{E}[\Phi_{\sigma^2}(w_{ij} + C'_{ij})] \\ &= \Phi_{\rho^2}(w_{ij} + \mathbb{E}[C'_{ij}])\end{aligned}$$

- ▷ use expectation form
- ▷ t_δ is small enough
- ▷ by the corollary

relation between θ and w :

$$\begin{aligned}\theta_{ij} &= \Pr(X_i(t)=1 \mid X_j[t_\delta]=1) \\ &= \mathbb{E}[\Phi_{\sigma^2}(w_{ij} + C'_{ij})] \\ &= \Phi_{\rho^2}(w_{ij} + \mathbb{E}[C'_{ij}])\end{aligned}$$

- ▷ use expectation form
- ▷ t_δ is small enough
- ▷ by the corollary

assumption

$$C'_{ij} \sim \mathcal{N}(\bar{C}_{ij}, \tau^2)$$

relation between θ and w :

$$\begin{aligned}\theta_{ij} &= \Pr(X_i(t)=1 \mid X_j[t_\delta]=1) \\ &= \mathbb{E}[\Phi_{\sigma^2}(w_{ij} + C'_{ij})] \\ &= \Phi_{\rho^2}(w_{ij} + \mathbb{E}[C'_{ij}])\end{aligned}$$

- ▷ use expectation form
- ▷ t_δ is small enough
- ▷ by the corollary

assumption

$$C'_{ij} \sim \mathcal{N}(\bar{C}_{ij}, \tau^2)$$

calculate θ by using w as:

$$\begin{aligned}\theta_{ij} &= \Phi_{\rho^2}(w_{ij} + \bar{C}_{ij}), \\ \bar{C}_{ij} &= \rho \cdot \Phi_1^{-1}(\Pr(X_i(t)=1 \mid X_j[t_\Delta]=0)).\end{aligned}$$

third step

estimate types of neurons consistent with data:

- excitatory neurons - positive connections only
- inhibitory neurons - negative connections only

third step

estimate types of neurons consistent with data:

- excitatory neurons - positive connections only
- inhibitory neurons - negative connections only


treated as hidden variables $\mathbf{z} \in \{0, 1\}^N$:

$$\Pr(\text{Data} \mid W, \mathbf{z}) \Leftrightarrow \Pr(\mathbf{z} \mid \text{Data}, W)$$

third step

estimate types of neurons consistent with data:

- excitatory neurons - positive connections only
- inhibitory neurons - negative connections only



treated as hidden variables $\mathbf{z} \in \{0, 1\}^N$:

$$\Pr(\text{Data} \mid W, \mathbf{z}) \Leftrightarrow \Pr(\mathbf{z} \mid \text{Data}, W)$$

use em algorithm (Amari 1995)


with approximations:

- factorial model in data manifold
- Gibbs sampling

Proposed Algorithm:


```
1: Input:  $\Lambda, \bar{C}, \mathbf{z}$ 
2: function ESTIMATEW( $\Lambda, \bar{C}, \mathbf{z}$ )
3:   Initialization:  $\Theta^{(1)} \leftarrow [0, 1]^{N \times N}, \Lambda^{(1)} \leftarrow \Lambda$ 
4:   for  $\tau \leftarrow 1, T$  do
5:      $W^{(\tau+1)} \leftarrow \Lambda^{(\tau)}(I - \Theta^{(\tau)})$ 
6:     for  $i \leftarrow 1, N$  do
7:       for  $j \leftarrow 1, N$  do
8:          $[\hat{W}(\mathbf{z})^{(\tau+1)}]_{ij} \leftarrow \begin{cases} z_j [W^{(\tau+1)}]_{ij}, & [W^{(\tau+1)}]_{ij} > 0 \\ (1 - z_j) [W^{(\tau+1)}]_{ij}, & [W^{(\tau+1)}]_{ij} < 0 \end{cases}$ 
9:       end for
10:    end for
11:     $[\Theta^{(\tau+1)}]_{ij} \leftarrow \Phi_1([\hat{W}(\mathbf{z})^{(\tau+1)}]_{ij} + \bar{C}_{ij})$ 
12:     $\text{diag}(\Theta^{(\tau+1)}) \leftarrow 0$  ▷ update diagonal elements
13:     $\Lambda^{(\tau+1)} \leftarrow \Lambda^{(\tau)}$ 
14:     $\text{diag}(\Lambda^{(\tau+1)}) \leftarrow \text{diag}(\Lambda^{(\tau)} \Theta^{(\tau+1)})$  ▷ update diagonal elements
15:  end for
16: end function
17: Output:  $\hat{W}(\mathbf{z})$ 
```



NUMERICAL EXAMPLES



Izhikevich's neuron model
(Izhikevich 2003)

- $N = 33$ out of 100 neurons
- excitatory:inhibitory = 80%:20%
- $w_{ij} \sim \text{Unif}[-10, 10]$
- $\#\{w_{ij}\} \leq 10$







remarks

- estimation is scale indeterminate
- inhibitory connections are difficult to estimate

sensitivity





Kendall coefficient



remarks

- estimation accuracy gets better if neuron types are given
- order of weights is estimated with sufficient accuracy




remark

- sensitivity is affected by choice of correlation interval


memory trace replay: (Wilson and McNaughton 1994; Tatsuno, Lipa, and McNaughton 2006)

- purpose: examine the hypothesis “the replay of activity patterns during sleep plays an important role in the consolidation process of memory”
- measurements:
 - pre-task: activity of control
 - task: activity in learning stage
 - post-task: activity in non-REM stage


pre-task



task



post-task



remarks

- some connections changed at task period are retained at post-task period (e.g. 8,11,12,20)
- result should be discussed from the viewpoint of biology

CONCLUDING REMARKS







we consider an approach to solve the following problems:






- pseudo correlation caused by higher-order effect
- influence from unobserved neurons
- directional excitatory/inhibitory connections

possible extension would be:

- estimating the number of connections
- estimating activation functions of individual neurons
- applying other real-world data

REFERENCES

-  Amari, Shun-ichi (Jan. 1995). “The EM Algorithm and Information Geometry in Neural Network Learning.” In: *Neural Computation* 7.1, pp. 13–18. DOI: [10.1162/neco.1995.7.1.13](https://doi.org/10.1162/neco.1995.7.1.13).
-  Friedman, Jerome, Trevor Hastie, and Robert Tibshirani (July 2008). “Sparse inverse covariance estimation with the graphical lasso.” In: *Biostatistics* 9.3, pp. 432–441. DOI: [10.1093/biostatistics/kxm045](https://doi.org/10.1093/biostatistics/kxm045).
-  Hyvärinen, Aapo (Aug. 6, 2002). “Gaussian moments for noisy independent component analysis.” In: *IEEE Signal Processing Letters* 6 (6), pp. 145–147. DOI: [10.1109/97.763148](https://doi.org/10.1109/97.763148).
-  Ito, Hiroyuki and Satoshi Tsuji (Jan. 2000). “Model Dependence in Quantification of Spike Interdependence by Joint Peri-Stimulus Time Histogram.” In: *Neural Computation* 12.1, pp. 195–217. DOI: [10.1162/089976600300015952](https://doi.org/10.1162/089976600300015952).
-  Izhikevich, Eugene M. (Nov. 2003). “Simple model of spiking neurons.” In: *IEEE Transactions on Neural Networks* 14 (6), pp. 1569–1572. DOI: [10.1109/TNN.2003.820440](https://doi.org/10.1109/TNN.2003.820440).
-  Kim, Sanggyun et al. (Mar. 24, 2011). “A Granger Causality Measure for Point Process Models of Ensemble Neural Spiking Activity.” In: *PLoS Computational Biology* 7.3. DOI: [10.1371/journal.pcbi.1001110](https://doi.org/10.1371/journal.pcbi.1001110).

-  Nakahara, Hiroyuki and Shun-ichi Amari (Oct. 2002). “Information-Geometric Measure for Neural Spikes.” In: *Neural Computation* 14.10, pp. 2269–2316. DOI: [10.1162/08997660260293238](https://doi.org/10.1162/08997660260293238).
-  Noda, Atsushi et al. (July 2014). “Intrinsic Graph Structure Estimation Using Graph Laplacian.” In: *Neural Computation* 26.7, pp. 1455–1483. DOI: [10.1162/NECO_a_00603](https://doi.org/10.1162/NECO_a_00603).
-  Tatsuno, Masami, Jean-Marc Fellous, and Shun-ichi Amari (Aug. 2009). “Information-Geometric Measures as Robust Estimators of Connection Strengths and External Inputs.” In: *Neural Computation* 21.8, pp. 2309–2335. DOI: [10.1162/neco.2009.04-08-748](https://doi.org/10.1162/neco.2009.04-08-748).
-  Tatsuno, Masami, Peter Lipa, and Bruce L. McNaughton (Oct. 18, 2006). “Methodological Considerations on the Use of Template Matching to Study Long-Lasting Memory Trace Replay.” In: *The Journal of Neuroscience* 26 (42), pp. 10727–10742. DOI: [10.1523/JNEUROSCI.3317-06.2006](https://doi.org/10.1523/JNEUROSCI.3317-06.2006). PMID: [17050712](https://pubmed.ncbi.nlm.nih.gov/17050712/).
-  Wilson, Matthew A. and Bruce L. McNaughton (July 29, 1994). “Reactivation of Hippocampal Ensemble Memories During Sleep.” In: *Science* 265 (5172), pp. 676–679. DOI: [10.1126/science.8036517](https://doi.org/10.1126/science.8036517). PMID: [8036517](https://pubmed.ncbi.nlm.nih.gov/8036517/).