



(a) Table 3: basis and component count (98 type-1 beats)



(b) Table 6: diastolic-decline variants (type 3, $n = 3,378$)



Markers in (a), basis family: $\bullet$ = Skewed Gaussian $\alpha \in [0, 8]$ (frozen version); $\blacksquare$ = Gaussian (no skew); $\triangle$ = Skewed Gaussian $\alpha \in [-8, 8]$ (Basso form); $\blacklozenge$ = Gamma (frozen search range); ∇ = Gamma (wide search range). Numbers, component count (in parentheses, strata with the predicted sign / strata evaluated). Points of one family are joined in order of component count.

Markers in (b): open circle, add a term for the diastolic decline; filled circle, exclude or down-weight the diastolic decline; grey square, reference fits. Numbers: (0) Frozen; (2) Delay bound 0.01 s; (4) Reservoir term; (4c) Reservoir, $\tau \geq 0.15$ s; (3) Exponential decay; (7) Output convolution; (8) Two-stage deconvolution; (6b) Truncate at 0.65T; (6c) Truncate at 0.55T; (6d) Truncate at 0.75T; (6e) Truncate at 0.45 s; (9) Derivative domain. The fraction after a number, strata with the predicted sign / strata evaluated. Vermilion dotted line, tier-C value of the fiducial-point analysis (database-supplied). Grey dashed line, criterion 0.30.

(a) and (b) are different subject sets: (a) 98 type-1 beats (notch present) from 120 subjects, 4 strata (table 3); (b) the 3,378 type-3 subjects (inflection only), 6 strata (table 6). Both y axes are the median within-age-stratum Spearman $|\rho|$ (tier C).

Tier A, subjects accepted by the method itself; tier B, subjects accepted by every compared method; tier C, all subjects regardless of acceptance

Exploratory, post hoc analysis; not used for the prespecified decision. Using an improved fit as a primary analysis would require a new preregistration.