

Immunoglobulin Reaction Report

For patients receiving immunoglobulin at home - IVIG (into a vein) or SCIG (under the skin). Please complete this form if you notice any unexpected side effect or reaction linked to your infusion, even if it seemed mild or settled on its own. It goes to your Immunology Specialist Nurse and Consultant.

YOUR IMMUNOLOGY SPECIALIST NURSES

020 7794 0500, ext. 32232 or 32233

rf-tr.clinicalimmunology@nhs.net

CALL 999 NOW IF YOU HAVE ANY OF THESE

Stop your infusion, stop filling in this form, and call 999 or go to your nearest A&E if you have:

- Difficulty breathing, wheezing, or tightness in your throat
- Swelling of your face, lips, tongue, or throat
- A widespread rash together with feeling faint, dizzy, or unwell
- Chest pain or pressure
- Weakness or numbness down one side, a drooping face, slurred speech, or sudden loss of vision
- Pain, swelling, warmth, or redness in one leg or arm
- Collapse or loss of consciousness

Tell the paramedics or A&E staff that you receive immunoglobulin (IVIG or SCIG) at home, and bring your batch number if you can.

CONTACT THE IMMUNOLOGY TEAM TODAY IF YOU HAVE

These need to be looked at the same day. Call the Immunology Specialist Nurses on 020 7794 0500, ext. 32232 or 32233, or NHS 111 if the line is closed - then complete this form afterwards:

- A severe headache with a stiff neck, dislike of bright light, or vomiting
- Dark, cola-coloured, or red urine
- Yellowing of your skin or the whites of your eyes
- Unusual breathlessness or extreme tiredness that is new
- Passing much less urine than normal
- A fever with shivering or shaking (rigors)
- A rash that keeps spreading

If you are unsure how urgent your symptom is, always ring rather than wait.

How to fill in and send this form

- Type your answers straight into this PDF, then save it.
- Have your product box or vial label to hand - we need the batch number from it.
- Fill in as much as you can. If you are not sure of an answer, leave it blank rather than delay sending it.
- When you have finished, email the completed form to rf-tr.clinicalimmunology@nhs.net.



Prefer to fill this in on your phone?

Scan this code with your phone camera to open the same form online. You can type your answers, stop and come back to them later, and email the finished report straight to us.

Or type in: immunoglobulin-reaction-report.vercel.app

SECTION 1

About you

So we can find your records and get back to you quickly.

Patient name

Date of birth

NHS or hospital number

Today's date

Best phone number today

Who is filling in this form?

- ☐ The patient
- ☐ A parent or guardian
- ☐ A carer or family member - name:

SECTION 2

Your immunoglobulin

Details of the infusion the reaction followed. The batch number is the most important item on this page - it lets us trace the exact product you received.

How do you receive your immunoglobulin?

- ☐ **IVIG**
Into a vein, through a drip or line
- ☐ **SCIG**
Under the skin, using a pump or rapid push
- ☐ **Facilitated SCIG**
Under the skin with hyaluronidase first, e.g. HyQvia

Brand name of your product

For example Privigen, Octagam, Intratect, Hizentra, Cuvitru, HyQvia, Gammaplex, Subgam, Gamunex.

Batch or lot number

Copy this exactly from the vial, bottle, or syringe label - letters and numbers. If you used more than one vial, please list every batch number.

Dose given (grams)

How often you have it

Date of that infusion

Time it started

Number of infusion sites used (SCIG)

For example: 2 sites, abdomen and thigh. Leave blank if you have IVIG.

Was this a new batch number?

A batch number you had not used before.

- ☐ Yes
☐ No
☐ Not sure

Have you changed brand or product recently?

- ☐ Yes, this is a different brand from my usual one
☐ No, it is my usual brand
☐ Not sure

Was the infusion running faster than usual?

- ☐ Yes, faster than usual
☐ No, my usual rate
☐ It was slower than usual
☐ Not sure

Did you take your usual pre-medication before this infusion?

For example paracetamol, an antihistamine such as cetirizine, or a steroid.

- ☐ Yes - what I took:
☐ No, I missed it this time
☐ I am not prescribed any pre-medication

Were you unwell in the days before this infusion?

For example a cold, chest infection, urine infection, or a temperature.

- ☐ Yes - what was wrong:
☐ No, I was well

SECTION 3

The reaction

Help us understand what you experienced and when.

1. When did the reaction start?

- ☐ During the infusion
☐ Within 1 hour of finishing
☐ 1 to 6 hours after
☐ 6 to 24 hours after
☐ 1 to 3 days after
☐ More than 3 days after

Date it started

Time it started

2. Please describe what happened in your own words

3. How long did the reaction last?

- ☐ It is still going on
- ☐ Less than 1 hour
- ☐ 1 to 6 hours
- ☐ 6 to 24 hours
- ☐ 1 to 3 days
- ☐ More than 3 days

4. Has this happened with your immunoglobulin before?

- ☐ No, this is the first time
- ☐ Yes, with this same product
- ☐ Yes, but with a different product
- ☐ Not sure

SECTION 4

What you did

Help us understand how you managed the reaction so far.

5. Did you slow down or stop the infusion?

- ☐ I slowed it down
- ☐ I paused it, then restarted
- ☐ I stopped completely and did not restart
- ☐ I carried on at the same rate
- ☐ Not applicable - I had already finished

6. If you slowed or stopped, did the symptoms improve?

- ☐ Yes, they settled
- ☐ They improved a little
- ☐ No change
- ☐ They got worse
- ☐ Not applicable

7. Did you finish the full dose?

- ☐ Yes, I had the full dose
- ☐ I had part of it - roughly how much:
- ☐ None of it

8. Did you take any medication to treat these symptoms?

For example paracetamol, ibuprofen, or an antihistamine such as cetirizine.

- ☐ No
- ☐ Yes - what and when:

Email your completed form to rf-tr.clinicalimmunology@nhs.net. This form does not replace urgent medical care - if your symptoms get worse, contact the Immunology team, NHS 111, or 999.