

1. SYLLABUS REVIEW

- (1) Pictures + Computer are ok so long as they're used for note taking.
- (2) Expect for the tests to be at ends of the first third of the class, and the second third of the class.
- (3) Theoretically this is a graduate course, and will be switched to 852, rather than remaining as 452.

1.1. Day 1.

- (1) The idea of algebraic topology
- (2) Given topological spaces X and Y , how can we prove that X and Y are or aren't homeomorphic.
- (3) To prove $X \cong Y$, we simply exhibit a homeomorphism.
E.g. $(-1, 1) \cong \mathbb{R}$, using $f(x) = \frac{x}{1-x^2}$
E.g. $\square \cong \circ$
- (4) To prove $X \not\cong Y$, we'd find a topological invariant, (connected, compact, Hausdorff,...), that only one has.
E.g. $(0, 1) \not\cong [0, 1]$, here, the closed interval is compact, and the open interval is not.
E.g. $(0, 1) \not\cong [0, 1)$, because,

$$\begin{aligned}[0, 1) \setminus \{0\} &= (0, 1) \text{ which is connected, but} \\ (0, 1) \setminus \{\text{any point}\} &\text{ is disconnected}\end{aligned}$$

Note, with the following exercise, If $X \cong Y$ via a homeomorphism, $\psi : X \rightarrow Y$, then $X \setminus \{p\} \cong Y \setminus \{\psi(p)\}$

- (5) Show the following.

$$\mathbb{R} \not\cong \mathbb{R}^2$$

Here, we note that $\mathbb{R} \setminus \{0\}$ is disconnected.

Suppose towards contradiction that $\mathbb{R} \cong \mathbb{R}^2$, call the homeomorphism $\phi : \mathbb{R} \rightarrow \mathbb{R}^2$, because $\mathbb{R} \setminus \{0\}$, the exercise implies that $\mathbb{R} \setminus \{0\} \cong \mathbb{R}^2 \setminus \{\phi(0)\}$, and therefore $\mathbb{R}^2 \setminus \{\phi(0)\}$ is disconnected, but that's just wrong, because \mathbb{R}^2 without a single point is still connected, rigorously showing this should be done through working with path connectedness. Therefore these are not homeomorphic.

$$\mathbb{R}^2 \not\cong \mathbb{R}^3$$

This was a trick question, we don't actually have any topological properties that we can rely on. If we were to attempt to remove a line from \mathbb{R}^2 , we don't have enough information about what the line is homeomorphic to in \mathbb{R}^3 , which is the major stumbling block.

- (6) The Fundamental Group

- (7) The fundamental group is a way to associate a topological space X to a group $\pi_1(X)$ so that $X \cong Y \Rightarrow \pi_1(X) \cong \pi_1(Y)$.
- (8) We'll be able to use this to prove spaces aren't homeomorphic.
Ex: In this course we'll learn the following.

$$\begin{aligned}\pi_1(\mathbb{R}^2 \setminus \{(0,0)\}) &= \mathbb{Z} \\ \pi_2(\mathbb{R}^3 \setminus \{\text{any point}\}) &= \{1\} \\ \pi_1(\mathbb{R}^2 \setminus \{(0,0)\}) &\not\cong \pi_2(\mathbb{R}^3 \setminus \{\text{any point}\}) \\ \mathbb{R}^2 &\not\cong \mathbb{R}^3\end{aligned}$$

Using this, we can show that these things are not homeomorphic, which is why we do algebraic topology. More powerful tools allow for more results.

- (9) Note: It's not true that $\pi_1(X) \cong \pi_1(Y) \Rightarrow X \cong Y$
 More generally, algebraic topology is about associating the topological space X with the algebraic object $A(X)$, in such a way that $X \cong Y \Rightarrow A(X) \cong A(Y)$
 There's a spectrum though.
- (a) Easy to compute and says nothing, $A(x)$ is the same for all of X
 - (b) Hard to compute, but says everything, $A(X) \cong A(Y) \iff X \cong Y$

1.2. Day 2.

- (1) The Fundamental Group
- (2) Idea: $\pi_1(X) = \{\text{"loops" in } X\} / \sim$, where $L_1 \equiv L_2$ if L_1 can be "deformed" inside X into L_2
- (3) Ex: Last time it was claimed that $\pi_1(\mathbb{R}^2 \setminus \{(0,0)\}) = \mathbb{Z}$.
- (4) Paths and Homotopies
- (5) Let X be a topological space.
- (6) Def: A path in X is a continuous map $f : I \rightarrow X$, where $I = [0, 1] \subseteq \mathbb{R}$ (with the subspace topology from the Euclidean topology on \mathbb{R}).
 If $f(0) = p$ and $f(1) = q$, we say f is a path from p to q .
- (7) Ex:


$$\begin{aligned}X &= \mathbb{R}^2 \\ f : I &\rightarrow \mathbb{R}^2 \\ f(t) &= (1 - 2t, 0)\end{aligned}$$

f is a path in \mathbb{R}^2 from $(1,0)$ to $(-1,0)$.

- (8) Another path in \mathbb{R}^2 from $(1,0)$ to $(-1,0)$ is,

$$\begin{aligned}g : I &\rightarrow \mathbb{R}^2 \\ g(t) &= (\cos(\pi t), \sin(\pi t))\end{aligned}$$

(9) To make precise, “Deforming” one path into another:



(10) Def: Let f and g be paths in X from p to q . A path homotopy from f to g is a continuous function,

$$H : I \times I \rightarrow X$$

(note that elements of $I \times I$ resemble, (s, t)) Such that,

$$H(s, 0) = f(s), \forall s$$

$$H(s, 1) = g(s), \forall s$$

$$H(0, t) = p, \forall t$$

$$H(1, t) = q, \forall t$$


To make sense of this, define, $\forall t$,

$$h_t : I \rightarrow X$$

$$h_t(s) = H(s, t)$$

Then, $\forall t$,

$$h_t = \text{path in } X \text{ from } p \text{ to } q$$



This is continuous because H is continuous, and it goes from p to q , because $h_t(0) = H(0, t) = p$ and $h_t(1) = H(1, t) = q$. $h_0(s) = f$ because $h_0(s) = H(s, 0) = f(s)$, $\forall s$ and $h_1(s) = g$ because $h_1(s) = H(s, 1) = g(s)$, $\forall s$

- (11) Def: If \exists a path homotopy from f to g , we say f and g are path-homotopic, and $f \cong g$
Ex: $X = \mathbb{R}^2$, Let,

$$f(s) = (\cos(\pi s), \sin(\pi s))$$

$$g(s) = (\cos(\pi s), 2 \sin(\pi s))$$

Both are paths in \mathbb{R}^2 from $(1, 0)$ to $(-1, 0)$.
 Then,

$$H : I \times I \rightarrow \mathbb{R}^2$$

$$H(s, t) = (\cos(\pi s), (t + 1) \sin(\pi s))$$

H is a path homotopy from f to g , because,

$$H(s, 0) = (\cos(\pi s), \sin(\pi s)) = f(s)$$

$$H(s, 1) = (\cos(\pi s), 2 \sin(\pi s)) = g(s)$$

$$H(0, t) = (\cos(0), (t + 1) \sin(0)) = (1, 0) \forall t$$

$$H(1, t) = (\cos(\pi), (t + 1) \sin(\pi)) = (-1, 0) \forall t$$

- (12) Question: Find a path homotopy from \mathbb{R}^2 from $f(s) = (s, s)$, and $g(s) = (s, s^2)$

Answer(June): $H(s, t) = (s, s^{t+1})$

(see the notebook, there's a solution there. Keep in mind that you want to try to find p and q first, before you do anything else)

Answer(Dr. Clader): General Trick In \mathbb{R}^2 let f and g be any two paths from p to q , then

the straight line homotopy is as follows,

$$H : I \times I \rightarrow \mathbb{R}^2$$

$$H(s, t) = (1 - t) * f(s) + t * g(s)$$

Note that this resembles the stuff you've seen in optimization and advanced linear algebra. This is a pretty powerful tool, remember and fear it.

- (13) Ex: In the question above, $H(s, t) = (s, (1 - t)s + ts^2)$

1.3. Day 3.

- (1) Products of Paths

- (2) Last time: If f and g are any two paths in \mathbb{R}^2 from p to q , then $f \cong_p q$.

- (3) By contrast: In, $S' = \{(x, y) \in \mathbb{R}^2 | x^2 + y^2 = 1\}$ if

$$f(s) = (\cos(\pi s), \sin(\pi s))$$

$$g(s) = (\cos(\pi s), -\sin(\pi s))$$

Then $f \not\cong_p g$. (We'll prove this carefully later).

- (4) Fact: (HW) \cong_p is an equivalence relation on the set {paths in X from x to y } Thus we can consider the set,


$$\{\text{paths in } X \text{ from } x \text{ to } y\} / \cong_p$$

$$= \{\text{path-homotopy classes of paths in } X \text{ from } x \text{ to } y\} \ni [f]$$

E.g. in the S' example above, $[f] \neq [g]$

- (5) Def: Let the following be so,

X = topological space
 f = path in X from x to y
 g = path in X from y to z




Then the concatenation of f and g is the path $f * g$ from x to z given by,

$$f * g : I \rightarrow X$$
$$(f * g)(s) = \begin{cases} f(2s) & \text{if } 0 \leq s \leq \frac{1}{2} \\ g(2s) & \text{if } \frac{1}{2} \leq s \leq 1 \end{cases}$$

(6) Why is $f * g$ continuous?

(7) Gluing Lemma: Let the following be so,


$$\begin{aligned} X &= \text{topological space} \\ A, B &\subseteq X, \text{ closed subsets such that } X = A \cup B \\ Y &= \text{topological space} \end{aligned}$$



Let the following continuous functions be defined,

$$\begin{aligned} f &: A \rightarrow Y \\ g &: B \rightarrow Y \end{aligned}$$

such that $f(x) = g(x) \forall x \in A \cap B$.



Then the function,

$$\begin{aligned} h &: X \rightarrow Y \\ h(x) &= \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B \end{cases} \end{aligned}$$

is continuous. The proof is left as an exercise to the reader. Thanks. (Homework Problem 1)

Note: Applying the gluing lemma to $I = [0, \frac{1}{2}] \cup [\frac{1}{2}, 1]$ shows that $f * g$ is continuous.

(8) Question: Let the following be so,

$$\begin{aligned} X &= \mathbb{R}^2 \\ f(s) &= (s - 1, s) \\ g(s) &= (s, s + 1) \end{aligned}$$

What is $f * g$? Draw a picture.

(9) Answer:

$$f * g = \begin{cases} f(2s) & \text{if } 0 \leq s \leq \frac{1}{2} \\ g(2s - 1) & \text{if } \frac{1}{2} \leq s \leq 1 \end{cases}$$

Which is a straight line from $(-1, 0)$ to $(1, 2)$.

(10) Proposition: $*$ is well defined on path-homotopy classes of paths
I.e., if,

$$\begin{aligned} f_0 &\cong_p f_1 \\ g_0 &\cong_p g_1 \end{aligned}$$

then,


$$f_0 * g_0 \cong_p f_1 * g_1$$

This means that if $[f] = \{\text{path-homotopy equivalence class of } f\}$ then we can define,

$$[f] * [g] := [f * g]$$

as long as the end point of f is the starting point of g .
So, now $*$ is an operation.

$$\{\text{paths from } x \rightarrow y\}_{/\cong_p} * \{\text{paths } y \rightarrow z\}_{/\cong_p} \rightarrow \{\text{paths } x \rightarrow z\}_{/\cong_p}$$



(11) Idea of proof of proposition:
 Let,

$F : I \times I \rightarrow X$ be a path homotopy from f_0 to f_1
 $G : I \times I \rightarrow X$ be a path homotopy from g_0 to g_1

Then we can define,

$$H : I \times I \rightarrow X$$

$$H(s, t) = \begin{cases} F(2s, t) & \text{if } 0 \leq s \leq \frac{1}{2} \\ G(2s - 1, t) & \text{if } \frac{1}{2} \leq s \leq 1 \end{cases}$$

Then,

$$\begin{aligned} h_0 &= H(s, 0) = (f_0 * g_0)(s) \\ h_1 &= H(s, 1) = (f_1 * g_1)(s) \\ h_t &= H(s, t) = (f_t * g_t)(s) \text{ (some path between } x \text{ and } z \text{)} \end{aligned}$$

So, H is a path homotopy from $(f_0 * g_0)$ to $(f_1 * g_1)$.

1.4. Day 4.

- (1) Definition of Fundamental Group
- (2) Recall: If,

$$\begin{aligned} f &= \text{path in } X \text{ from } x \text{ to } y \\ g &= \text{path in } X \text{ from } y \text{ to } z \end{aligned}$$

Then,


$$[f] * [g] := [\text{concatenation } f * g \text{ of } f \text{ and } g]$$

- (3) Properties of $*$:
 - (a) $*$ is associative, or


$$[f] * ([g] * [h]) = ([f] * [g]) * [h]$$

The idea here is that we can adjust the time taken to travel on the path. These two paths are path-homotopic: interpolate between $f * (g * h)$ and $(f * g) * h$ by making f take less and less time and h take more and more time.

$$f * (g * h):$$



$$(f * g) * h:$$




- (b) $*$ has left/right identities.
Let

$$e_x : I \rightarrow X$$

$$e_x(s) = x, \forall s \in I, \text{ "constant path at } x \text{"}$$

Then, for all paths f from x to y , $[f] * [e_y] = [f]$, and $[e_x] * [f] = [f]$. The premise here is that e_x or e_y spend "half the time" sitting at either x or y .



These are path-homotopic: interpolate between $f * e_y$ and f by making f take longer and longer.

(c) $*$ has inverses.


Let f be a path from x to y , and let \bar{f} be the “reverse” path,

$$\bar{f}(s) = f(1 - s)$$

Then,

$$[f] * [\bar{f}] = [e_x]$$

$$[\bar{f}] * [f] = [e_y]$$



Idea: The verbal gist of this is that the path takes half the time to travel to its destination, and is concatenated with a path that spends half the time to travel to the origin of the original function.

These are path-homotopic: interpolate between $f * \bar{f}$ and e_x by doing less and less of f before turning around.

(d) Let,

X = topological space

$x \in X$

Definition: A loop in X based at $x \in X$ is a path,

$$f : I \rightarrow X$$

such that $f(0) = f(1)$



(e) Observation: If f and g are any two loops in X based at x , then $f * g$ is a loop.

(f) Definition: The fundamental group of the X with basepoint x is:

$$\pi(X, x) = \{\text{path-homotopy classes of loops in } X \text{ based at } x\}$$

This is a group with the operation $*$

- (i) e_x and e_y are loops.
- (ii) $f * \bar{f}$ and $\bar{f} * f$ are also loops.
- (iii) Good question Katy!
- (g) Note: The fact that $\pi_1(X, x)$ satisfies the axioms of a group, and follows from the properties of $*$ we just checked.
(E.g. the identity element is $[e_x]$)
- (h) Question: What is $\pi_1(\mathbb{R}^2, (0, 0))$?
Do you have a guess for $\pi_1(S', (1, 0))$?
Answer 1: $\pi_1(\mathbb{R}^2, (0, 0)) \cong \{1\}$
To prove this, it's enough to show that $\pi_1(\mathbb{R}^2, (0, 0))$ has just one element, i.e., any loop in \mathbb{R}^2 based at $(0, 0)$, is path-homotopic to any other. This is true via the straight line homotopy. Answer 2: $\pi_1(S', (1, 0)) \cong \mathbb{Z}$.

1.5. Day 5.

- (1) π_1 continued: To what extent does π_1 depend on x ?
- (2) Theorem: Let X be a path-connected topological space, and let $x_0, x_1 \in X$, then $\pi_1(X, x_0) \cong \pi_1(X, x_1)$. This section builds off the worksheet provided in class.
 - (a) Part 1: see drawing
 - (b) Part 2: Let f and g be in $\pi_1(X, x_1)$

$$\begin{aligned}\hat{\alpha}([f] * [g]) &= [\bar{\alpha}] * [f] * [g] * [\alpha] \\ &= [\bar{\alpha}] * [f] * [\alpha] * [\bar{\alpha}] * [g] * [\alpha] \\ &= \hat{\alpha}([f]) * \hat{\alpha}([g])\end{aligned}$$
 - (c) Part 3: Let $f \in \pi_1(X, x_1)$

$$\begin{aligned}\hat{\alpha}([f]) &= [\bar{\alpha}] * [f] * [\alpha] \\ \hat{\alpha}([\bar{\alpha}] * [f] * [\alpha]) &= [\alpha] * [\bar{\alpha}] * [f] * [\alpha] * [\bar{\alpha}] \\ &= [f]\end{aligned}$$
 - (d) Therefore this mfer is an isomorphism.
- (3) For which topological spaces X can we actually compute $\pi_1(X, x)$?
- (4) Definition: A topological space X is simply-connected if
 - (a) X is path connected
 - (b) $\pi_1(X, x) = 1 \ \forall x \in X$
(Because X is path connected, we only need to check this for one $x \in X$)
- (5) Ex: \mathbb{R}^2 is simply connected
- (6) Intuition: X is simply-connected if any loop in X if any loop in X can be “shrunk down” to a constant loop.
(for all loops f in X saying f can be “shrunk down” means $f \cong_p c_x$ where c_x is a constant path)
- (7) Next time: A convex subset of \mathbb{R}^n is simply connected.

1.6. Day 6.

- (1) Goal: Prove that $\pi_1(S^1, x) \cong \mathbb{Z}$

- (2) Idea: S^1 can be built by “wrapping \mathbb{R} around itself”.
: Concretely, this is

$$p : \mathbb{R} \rightarrow S^1$$

$$p(x) = (\cos(2\pi x), \sin(2\pi x))$$

We’ll try to “unwrap” loops in S^1 to get paths in \mathbb{R}

- (3) The above map p is an example of a “covering map”. The ultimate goal of today is to understand what it means to be a covering map, before we get to the definition of it.
- (4) Questions: Let the following be so,

$$u_1 = \{(x, y) \in S^1 | y > 0\}$$

$$u_2 = \{(x, y) \in S^1 | x > 0, y < 0\}$$

Include the drawings from class, really get sick wit it.

- (5) Observation: For any particular $n \in \mathbb{Z}$, the piece,

$$(n, n + \frac{1}{2}) \cong u_1$$

The homeomorphism in Dr. Clader’s mind is,

$$\phi : (n, n + \frac{1}{2}) \rightarrow u_1$$

$$\phi(x) = (\cos(2\pi x), \sin(2\pi x))$$

i.e. $\phi = p|_{(n, n + \frac{1}{2})}$

The inverse of ϕ is,

$$\phi^{-1} : u_1 \rightarrow (n, n + \frac{1}{2})$$

$$\phi^{-1} = \frac{\cos^{-1}(x)}{2\pi} + n$$

(Recall: by definition $\cos^{-1}(x) \in [0, \pi]$)

Similarly, for u_2 for any particular $n \in \mathbb{Z}$, $(n - \frac{1}{4}, n) \cong u_2$.

- (6) Definition: Let $p : E \rightarrow B$ be a function between two topological spaces. We say p is a covering map if p is,
- (a) p is continuous and surjective
 - (b) $\forall b \in B$ there exists a neighborhood u of b such that,

$$p^{-1}(u) = \cup_{\alpha} v_{\alpha}$$

where $v_{\alpha} \subseteq E$ are open, disjoint and,

$$p|_{v_{\alpha}} : v_{\alpha} \rightarrow u$$

is a homeomorphism for every α . Note that these open subsets with this property are called evenly covered

Note that b is one particular point or neighborhood, but there should be a neighborhood for every single point in B where all of this junk holds reasonably truish.

(7) Ex:

$$p : \mathbb{R} \rightarrow S^1$$

$$p(x) = (\cos(2\pi x), \sin(2\pi x))$$

p is a covering map. We just showed that u_1 is evenly covered:

$$p^{-1}(u_1) = \cup_{n \in \mathbb{Z}} (n, n + \frac{1}{2})$$

Note that in this case the $(n, n + \frac{1}{2})$ are the v_α from the definition of covering maps. u_2 is also evenly covered, but, $U = S^1$ is not evenly covered because, $p^{-1}(S^1) = \mathbb{R}$, and the only way to write \mathbb{R} as a union of disjoint open sets v_α , is to take $v_\alpha = \mathbb{R}$, but $\mathbb{R} \not\cong S^1$

(8) Ex:

$$B = \text{any space}$$

$$E = B \times \{1, 2, \dots, n\} = n \text{ discrete copies of } B$$

Where $\{1, 2, \dots, n\}$ is equipped with the discrete topology.

1.7. Day 7.

- (1) Guest lecturer: Mattias “i think your regular lecturer is more qualified for this” Beck
- (2) Recalling the definition of an evenly covered set. New notation was introduced, but L^AT_EX is behind the times. Let E and B be topological spaces

$$\phi : E \rightarrow B$$

$$\forall b \in B, \exists u \text{ a neighborhood of } b : p^{-1}(u) = \cup_{\alpha} v_{\alpha}$$

$$p|_{v_{\alpha}} : v_{\alpha} \rightarrow u$$

(3) Fun notation facts:

\rightarrow indicates a surjective function

\hookrightarrow indicates an injective function

Combining the two gives you a bijective function, but that symbol doesn't exist in latex apparently.

(4) Example covering:

$$E = \mathbb{R}$$

$$\phi(x) = (\cos(2\pi x), \sin(2\pi x))$$

$$B = S^1$$

- (5) Definition: Given a covering map from topological spaces E to B

$$p : E \rightarrow B$$

a path in our topological space B ,

$$f : I \rightarrow B$$

A lift of f is a path, $\tilde{f} : I \rightarrow E$, such that $f = p \circ \tilde{f}$

- (6) Theorem: Given covering map $p : E \rightarrow B$, $p(e) = b$, $f : I \rightarrow B$ path beginning at b , then there does not exist a left \tilde{f} , of f beginning at e Read Lemma 54.1 Munkres. (!!?!?)
- (7) The same theorem but reworded:
Let the following be so,

E be a topological space

B be a topological space

$p : E \rightarrow B$ a covering map

$f : I \rightarrow B$ path beginning at b

$e \in E$, s.t. $p(e) = b$

Then there exists a unique path, \tilde{f} in E such that $p \circ \tilde{f} = f$, and $\tilde{f}(0) = e$

1.8. Day 8.

- (1) Guest Lecturer: Matthias “you can have a hint, but you can’t quote me on it” Beck
- (2) ???????

1.9. Day 9.

- (1) Guest Lecturer: Anastasia the Assassin, Deadly David, and Killa Katy
- (2) Let p be a covering map.

$$p : E \rightarrow B$$

Let, $e \in E$, $b \in B$, such that $p(e) = b$.

Summary of what we know about this situation,

- (a) Any path f in B , beginning at b has a unique lift \tilde{f} to a path in E beginning at e .
- (b) If f and g are two paths in B , beginning at b , such that $f \cong_p g$, then $\tilde{f} \cong_p \tilde{g}$
- (c) If f is a loop in B based at b , then $\tilde{f} \in p^{-1}(b)$

1.10. Day 10.

- (1) $p_1(S^1)$, continued:
- (2) Recap:

$$p : \mathbb{R} \rightarrow S^1$$

$$p(x) = (\cos(2\pi x), \sin(2\pi x))$$

Then there exists a function,

$$\begin{aligned}\phi : \pi_1(S^1, b) &\rightarrow p^{-1}(b) \\ \phi([f]) &= \tilde{f}(1)\end{aligned}$$

Where \tilde{f} is the lift of f to \mathbb{R} starting at 0.
E.g., (draw that spiraleboye)

$$\begin{aligned}\phi([\text{loop once counterclockwise}]) &= 1 \\ \phi([\text{loop twice counterclockwise}]) &= 2 \\ \phi([\text{loop once clockwise}]) &= -1\end{aligned}$$

The fact that there exists a unique lift, \tilde{f} of any f is a feature of covering maps.
In fact,

$$p^{-1}(b) = \mathbb{Z}$$

and,

(3) Claim: $\phi : \pi_1(S^1, b) \rightarrow \mathbb{Z}$ is a bijection.

Proof. (a) Surjective: Given $c \in \mathbb{Z}$, choose a path, $\alpha : I \rightarrow \mathbb{R}$, from 0 to c in \mathbb{R} . Then let,
 $f : I \rightarrow S^1$ be $f = p \circ \alpha$
Then f is a loop in S^1 based at $b = (1, 0)$ because

$$\begin{aligned}f(0) &= p(\alpha(0)) = p(0) = (1, 0) \\ f(1) &= p(\alpha(1)) = p(c) = (1, 0)\end{aligned}$$

And, $\tilde{f} = \alpha$ because $p \circ \tilde{f} = p \circ \alpha = f$. Thus,

$$\phi([f]) = \tilde{f}(1) = \alpha(1) = c$$

(b) Injective: Suppose,

$$\begin{aligned}\phi([f]) &= \phi([g]) \\ \implies \tilde{f}(1) &= \tilde{g}(1)\end{aligned}$$

Then, \tilde{f} and \tilde{g} are two paths in \mathbb{R} , that both start at 0 and both end at the same point.
 \Rightarrow (courtesy of homework 2) $\tilde{f} \cong_p \tilde{g}$ (because \mathbb{R} is simply connected)
 $\Rightarrow p \circ \tilde{f}$ is a path homotopy from $p \circ \tilde{f}$ to $p \circ \tilde{g}$.
 $\Rightarrow f \cong_p g$
 $\Rightarrow [f] = [g] \in \pi_1(S^1, b)$

□

(4) Claim: ϕ is a group homomorphism (thus, an isomorphism).

Proof. Let $[f], [g] \in \pi\pi_1(S^1, b)$, we want to show that, $\phi([f] * [g]) = \phi([f]) + \phi([g])$
 By definition,

$$\phi([f] * [g]) = \phi([f * g]) = f * \tilde{g}(1)$$

What is $f * \tilde{g}$? By definition $f * \tilde{g}$ is the lift of $f * g$ starting at 0 and,

$$\begin{aligned}\tilde{f} &= \text{lift of } f \text{ starting at 0 ending at some } n \\ \tilde{g} &= \text{lift of } g \text{ starting at 0 ending at some } m\end{aligned}$$

So, $\tilde{f} * \tilde{g}$ doesn't make sense, but let:

$$\begin{aligned}\tilde{g}' &= \text{"shift } \tilde{g} \text{ by } n \text{"} \\ \text{i.e., } \tilde{g}' &= g(s) + n\end{aligned}$$

Now notice that $\tilde{f} * \tilde{g}'$ now makes sense, and \tilde{g}' is a lift of g , because:

$$\begin{aligned}(p \circ \tilde{g}')(s) &= p(\tilde{g}(s)) \\ &= p(\tilde{g}(s) + n) \\ &= p(\tilde{g}(s)) \\ \text{because } p(x + n) &= p(x), \forall n \in \mathbb{Z} \\ &= (p \circ \tilde{g})(s) \\ &= g(s)\end{aligned}$$

Thus, $\tilde{f} * \tilde{g}'$ is a lift of $f * g$ starting at 0

$$\begin{aligned}\implies \tilde{f} * \tilde{g}' &= f * \tilde{g} \\ f * \tilde{g}'(1) &= \tilde{f} * \tilde{g} \\ &= \text{endpoint of } \tilde{g}' \\ &= \tilde{g}'(1) + n \\ &= m + n\end{aligned}$$

This shows that

$$\begin{aligned}\phi([f] * [g]) &= m + n \\ &= \tilde{f}(1) + \tilde{g}(1) \\ &= \phi([f]) + \phi([g])\end{aligned}$$

□

(5) We want:

$$X \cong Y \implies \pi_1(X, x) \cong \pi_1(Y, y)$$

(X is homeomorphic to Y)

The big tool we'll use to do that is the tool from the second homework about maps between spaces being homomorphisms. That's for next time!

1.11. Day 11.

- (1) Note that this Friday, office hours will be at 3-4pm.
- (2) We want: If $X \cong Y$, then $\pi_1(X, x) \cong \pi_1(Y, y)$, or that, if two spaces are homeomorphic, then their fundamental groups are isomorphic. We will explore the tools used to show this in this lecture
- (3) Definition(HW2): Let $\varphi : X \rightarrow Y$, be a continuous map, then the homomorphism induced by φ is:

$$\begin{aligned}\varphi_* : \pi_1(X, x) &\rightarrow \pi_1(Y, y) \\ \varphi_*([f]) &= [\varphi \circ f]\end{aligned}$$

See the picture of the picture drawn on the board, make a drawyboye.

- (4) Lemma: (this is referred to lemma 1) If

$$X \xrightarrow{\varphi} Y \xrightarrow{\psi} Z$$

Where φ and ψ are both continuous, then,

$$(\psi \circ \varphi)_* = \psi_* \circ \varphi_*$$

Additionally, (This is referred to as lemma 2)

$$id_* = id$$

(or that given the $id : X \rightarrow Y$, the induced homomorphism, $\pi_1(X, x) \rightarrow \pi_1(Y, y)$ is the identity)

- (5) *Proof.* (a) Both sides are homomorphisms,

- (1) $\pi_1(X, x) \rightarrow \pi_1(Z, (\psi \circ \varphi)(x))$

- (2)

Given any $[f] \in \pi_1(X, x)$:

$$\begin{aligned}(\psi \circ \varphi)_*([f]) &= [(\psi \circ \varphi) \circ f] \\ &= [\psi \circ (\varphi \circ f)] \\ &= \psi_*[\varphi \circ f] \\ &= \psi_*(\varphi_*([f])) \\ &= (\psi_* \circ \varphi_*)([f])\end{aligned}$$

(b) Given any $[f] \in \pi_1(X, x)$:

$$\begin{aligned} id_*([f]) &= [id \circ f] \\ &= [f] \end{aligned}$$

□

(6) Theorem: if $\varphi : X \rightarrow Y$ is a homeomorphism, then $\varphi_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ is an isomorphism.

Proof. We already know that φ_* is a homomorphism, to prove that it's a bijection, we'll find an inverse to φ_* . Claim that,

$$(\varphi)_* : \pi_1(Y, \varphi(x)) \rightarrow \pi_1(X, x)$$

is the inverse to φ_* .

(Note that this is doable, because φ is a homeomorphism, $\varphi^{-1} : Y \rightarrow X$ exists, and is continuous)

To check this:

$$\begin{aligned} &\varphi_* \circ (\varphi^{-1})_* \\ &= (\varphi \circ \varphi^{-1})_*, \text{ by lemma 1 shown today} \\ &= id_*, \text{ by definition of } \varphi^{-1} \text{ (identity on } y) \\ &= id, \text{ by lemma 2 shown today (identity on } x) \\ &(\varphi^{-1})_* \circ \varphi_* = (\varphi^{-1} \circ \varphi)_* = id_* = id \end{aligned}$$

This by definition means φ_* and $(\varphi^{-1})_*$ are inverse functions. Additionally, this small red box has made it onto the board, for clarification.

$$\begin{aligned} id_x : X &\rightarrow Y \\ id_{\pi_1(X, x)} : \pi_1(X, x) &\rightarrow \pi_1(X, x) \\ \text{Lemma: } (id_x)_* id_{\pi_1(X, x)} & \end{aligned}$$

□

(7) This ends up proving that,

$$X \cong Y \implies \pi_1(X, x) \cong \pi_1(Y, \varphi(x))$$

But, non-homeomorphic spaces can have isomorphic π_1

Ex:

$$\begin{aligned} X &= . \\ Y &= \mathbb{R}^2 \end{aligned}$$

These are not homeomorphic, clearly X is compact and Y isn't, but their fundamental groups are isomorphic, since the fundamental group of X is just $\{1\}$, and clearly this is also true about \mathbb{R}^2

(8) So, given X and Y , how can we tell if $\pi_1(X) \cong \pi_1(Y)$?

(9) Homotopy of Maps:

Definition: Let $f : X \rightarrow Y$ and $g : X \rightarrow Y$ be continuous functions. Then a homotopy from f to g is a continuous function,

$$H : X \times I \rightarrow Y$$

such that,

$$H(x, 0) = f(x), \forall x \in X$$

$$H(x, 1) = g(x), \forall x \in X$$

Our goal is to make remark about the lower star versions of these maps, given their being homotopic.

1.12. **Day 12.**

(1) Homotopy of maps: Definition: Let $f : X \rightarrow Y$ be a continuous function. A homotopy from f to g is a continuous function,

$$H : X \times I \rightarrow Y$$

such that

$$H(x, 0) = f$$

$$H(x, 1) = g$$

We'll often write,

$$h_t : X \rightarrow Y$$

$$h_t(x) = H(x, t)$$

Then there's one h_t for each $t \in I$ and,

$$h_0 = f$$

$$h_1 = g$$

h_t = "A function interpolating between f and g "

(2) Terminology/Notation: If there exists a homotopy from f to g , we'll say that f is homotopic to g and write $f \cong g$.

(3) Ex:

$$f : S^1 \rightarrow \mathbb{R}^2$$

$$g : S^1 \rightarrow \mathbb{R}^2$$

$$f(x, y) = (x, y)$$

$$g(x, y) = (0, 0)$$

Then $f \cong g$. A homotopy from f to g is,

$$H : S^1 \times I \rightarrow \mathbb{R}^2$$

$$H((x, y), t) = ((1 - t)x, (1 - t)y)$$

Do the drawing from the board.

(4) Ex:

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$g : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = x$$

$$g(x) = x + 2$$

Then $f \cong g$. A homotopy from f to g is:

$$H : \mathbb{R} \times I \rightarrow \mathbb{R}$$

$$H(x, t) = x + 2t$$

Refer again to the picture from the board.

(5) Questions:

(a)

$$f : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$g : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$f(x) = (x, 0)$$

$$g(x) = (x, e^x)$$

(b)

$$f : \mathbb{R}^2 \setminus (0, 0) \rightarrow \mathbb{R}^2 \setminus (0, 0)$$

$$g : \mathbb{R}^2 \setminus (0, 0) \rightarrow \mathbb{R}^2 \setminus (0, 0)$$

$$f(x) = (x, y)$$

$$g(x) = \left(\frac{x}{\sqrt{x^2 + y^2}}, \frac{y}{\sqrt{x^2 + y^2}} \right)$$

$$f : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$g : \mathbb{R} \rightarrow \mathbb{R}^2$$

$$f(x) = (x, 0)$$

$$g(x) = (x, e^x)$$