

# Fitting Social Network Models to Data

---

Mark S. Handcock

handcock@ucla.edu

Statistical Analysis of Networks

October 22, 2024

# Fitting a General Model to Data

- Suppose we have network data and want to see how the models fit
- Initially focus on models where the  $Y_{ij}$  are independent (given covariates  $X$ ), so that

$$\text{logit}[P(Y_{ij} = 1|X = x, \beta)] = \beta_1 x_{1,ij} + \beta_2 x_{2,ij} \dots + \beta_K x_{K,ij}$$

for some choice of covariates  $x_{1,ij}, x_{2,ij}, \dots, x_{K,ij}$

# Estimation based on the Likelihood Function

- Develop a framework based on the likelihood function
- What is maximum likelihood estimation?  
Find  $\beta$  that makes the observed network most likely under the model with  $\beta$  as the parameter.

The log of the *likelihood* (as a function of  $\beta$ ) is

$$\ell(\beta|y, x) \equiv \log[P(Y = y|X = x, \beta)] \quad \beta \in \mathbf{R}^k, \text{ given } y \in \mathcal{Y}$$

- $Y_{ij}$  are independent and equally likely

$$P(Y_{ij} = 1 | X = x, \beta) = \frac{\exp(\beta)}{1 + \exp(\beta)} \quad \forall \ i, j = 1, \dots, n$$

for some  $-\infty < \beta < \infty$

# Example: Homogeneous Bernoulli model Rényi and Erdős model

- $Y_{ij}$  are independent and equally likely

$$P(Y_{ij} = 1 | X = x, \beta) = \frac{\exp(\beta)}{1 + \exp(\beta)} \quad \forall \ i, j = 1, \dots, n$$

for some  $-\infty < \beta < \infty$

- Equivalently

$$\log \text{odds}(Y_{ij} = 1 | X = x, \beta) = \beta \quad \forall \ i, j = 1, \dots, n$$

## Example: Homogeneous Bernoulli model Rényi and Erdős model

More abstractly:

$$P(Y = y|X = x, \beta) = \frac{e^{\beta g(y)}}{\kappa(\beta)} \quad y \in \mathcal{Y}$$

$$\kappa(\beta) = [1 + \exp(\beta)]^N$$


$$g(y) = \sum_{i,j} y_{ij} \quad \text{the number of ties}$$

The log of the *likelihood* (as a function of  $\beta$ ) is

$$\ell(\beta|y, x) \equiv \log[P(Y = y|X = x, \beta)] = \beta g(y) - N \log[1 + \exp(\beta)] - \infty$$

# Example: Effect on the number of Priorates on centrality

Let  $X$  be the number of priorates (seats on the civic council) held by the family between 1282- 1344.



# Joint specification

Recall

$$P(Y = y|X = x, \beta) = \frac{e^{\beta_1 g_1(y) + \beta_2 g_2(y) \dots + \beta_K g_K(y)}}{c(\beta)} \quad y \in \mathcal{Y}$$

where

$$g_k(y) = \sum_{i,j} x_{1,ij} y_{ij}, \quad k = 1, \dots, K$$

for the covariates  $x_{1,ij}, x_{2,ij}, \dots, x_{K,ij}$

The *log-likelihood* is

$$\begin{aligned} \ell(\beta; y, x) &\equiv \log [P(Y = y|X = x, \beta)] \\ &= \beta_1 g_1(y) + \beta_2 g_2(y) \dots + \beta_K g_K(y) \\ &\quad - \sum_{i,j} \log[1 + \exp(\sum_{k=1}^K \beta_k x_{k,ij})] \end{aligned}$$



# Maximum Likelihood Estimation

Our goal is to find the

**Maximum likelihood estimator (MLE):**

The maximizer  $\hat{\beta}$  of the loglikelihood function,  $\ell(\beta|y, x)$

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

- if the likelihood function is very concentrated around  $\hat{\beta}$

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

- if the likelihood function is very concentrated around  $\hat{\beta}$
- Based on a Central Limit Theorem, as  $n \rightarrow \infty$ , the distribution of  $\hat{\beta}$  approaches a Gaussian with:

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

- if the likelihood function is very concentrated around  $\hat{\beta}$
- Based on a Central Limit Theorem, as  $n \rightarrow \infty$ , the distribution of  $\hat{\beta}$  approaches a Gaussian with:

$$\mathbb{E}_Y(\hat{\beta}) \rightarrow \beta \quad \text{as } n \rightarrow \infty$$

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

- if the likelihood function is very concentrated around  $\hat{\beta}$
- Based on a Central Limit Theorem, as  $n \rightarrow \infty$ , the distribution of  $\hat{\beta}$  approaches a Gaussian with:

$$\mathbb{E}_Y(\hat{\beta}) \rightarrow \beta \quad \text{as } n \rightarrow \infty$$

$$\mathbb{V}_Y(\hat{\beta}) \rightarrow I^{-1}(\beta) \quad \text{as } n \rightarrow \infty$$

where

$$I(\beta) = \mathbb{E}_Y \left[ -\frac{\partial^2 \ell(\beta; y, x)}{\partial \beta^T \partial \beta}(\beta) \right] \quad (1)$$

is the *information matrix* for  $\beta$

## Computing standard errors for $\hat{\beta}$

- How accurate an estimate is  $\hat{\beta}$  of  $\beta$ ?

This depends on how “spiked” or “peaked” the likelihood function is.

- if the likelihood function is very concentrated around  $\hat{\beta}$
- Based on a Central Limit Theorem, as  $n \rightarrow \infty$ , the distribution of  $\hat{\beta}$  approaches a Gaussian with:

$$\mathbb{E}_Y(\hat{\beta}) \rightarrow \beta \quad \text{as } n \rightarrow \infty$$

$$\mathbb{V}_Y(\hat{\beta}) \rightarrow I^{-1}(\beta) \quad \text{as } n \rightarrow \infty$$

where

$$I(\beta) = \mathbb{E}_Y \left[ -\frac{\partial^2 \ell(\beta; y, x)}{\partial \beta^T \partial \beta}(\beta) \right] \quad (1)$$

is the *information matrix* for  $\beta$

- We can compute  $p$ -values and confidence intervals



## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?

## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?
- Compared to what?

## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?
- Compared to what?
- The model were all networks are equally likely?

## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?
- Compared to what?
- The model were all networks are equally likely?
  - For this model  $\beta = 0$
- Use the likelihood and compare  $\ell(\beta = \hat{\beta}|y, x)$  to  $\ell(\beta = 0|y, x)$ .

## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?
- Compared to what?
- The model were all networks are equally likely?
  - For this model  $\beta = 0$
- Use the likelihood and compare  $\ell(\beta = \hat{\beta}|y, x)$  to  $\ell(\beta = 0|y, x)$ .
- For historical reason, use the *deviance*:  
$$\text{deviance}(\beta) = -2\ell(\beta|y, x)$$

## How well does the model fit the network?

- How can we tell if the estimated model is “likely”?
- Compared to what?
- The model were all networks are equally likely?
  - For this model  $\beta = 0$
- Use the likelihood and compare  $\ell(\beta = \hat{\beta}|y, x)$  to  $\ell(\beta = 0|y, x)$ .
- For historical reason, use the *deviance*:  
$$\text{deviance}(\beta) = -2\ell(\beta|y, x)$$
- In particular  
$$\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$$

## Example: Effect on the number of Priorates on centrality

For  $k = 1, \dots, n$ :

$$x_{ij}^{\text{priorates}} = \text{Number of priorates for } i + \text{Number of priorates for } j$$

$$x_{ij}^{\text{density}} = 1/N$$

## Example: Effect on the number of Priorates on centrality

Formula: `flomarriage ~ density + nodecov("priorates")`

	Estimate	Std. Error	p-value
density	-307.82	64.181	<1e-04 ***
nodecov.priorates	0.01643	0.0073	0.0276 *

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05

Null	Deviance: 166.355	on 120	degrees of freedom
Residual	Deviance: 102.921	on 118	degrees of freedom
	Deviance: 63.435	on 2	degrees of freedom



Quantifying how well the model fits the network?

Refer to deviance( $\beta = 0$ ) as the *null deviance*

## Quantifying how well the model fits the network?

Refer to deviance( $\beta = 0$ ) as the *null deviance*

Refer to deviance( $\beta = \hat{\beta}$ ) as the *residual deviance*

## Quantifying how well the model fits the network?

Refer to  $\text{deviance}(\beta = 0)$  as the *null deviance*

Refer to  $\text{deviance}(\beta = \hat{\beta})$  as the *residual deviance*

Refer to  $\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$   
as (just) the *deviance*

## Quantifying how well the model fits the network?

Refer to  $\text{deviance}(\beta = 0)$  as the *null deviance*

Refer to  $\text{deviance}(\beta = \hat{\beta})$  as the *residual deviance*

Refer to  $\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$   
as (just) the *deviance*

When there are large numbers of actors, the sampling  
distribution of the deviance

$$\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$$

has been worked out.

## Quantifying how well the model fits the network?

Refer to  $\text{deviance}(\beta = 0)$  as the *null deviance*

Refer to  $\text{deviance}(\beta = \hat{\beta})$  as the *residual deviance*

Refer to  $\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$   
as (just) the *deviance*

When there are large numbers of actors, the sampling  
distribution of the deviance

$$\text{deviance}(\beta = 0) - \text{deviance}(\beta = \hat{\beta})$$

has been worked out.

It is approximately  $\chi^2$  on  $K$  degrees of freedom.

## Example: Effect on the number of Priorates on centrality

Formula: `flomarriage ~ density + nodecov("priorates")`

	Estimate	Std. Error	p-value
density	-307.82	64.181	<1e-04 ***
nodecov.priorates	0.01643	0.0073	0.0276 *

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05

Null	Deviance: 166.355	on 120	degrees of freedom
Residual	Deviance: 102.921	on 118	degrees of freedom
	Deviance: 63.435	on 2	degrees of freedom

Does the model fit perfectly?

Compare the *residual deviance* to a  $\chi^2$  on  $N - K$  degrees of freedom.

## Example: Effect on the number of Priorates on centrality

Formula: `flomarriage ~ density + nodecov("priorates")`

	Estimate	Std. Error	p-value
density	-307.82	64.181	<1e-04 ***
nodecov.priorates	0.01643	0.0073	0.0276 *

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05

Null	Deviance: 166.355	on 120	degrees of freedom
Residual	Deviance: 102.921	on 118	degrees of freedom
	Deviance: 63.435	on 2	degrees of freedom